

Increased $p\text{CO}_2$ changes the lipid production in important aquacultural feedstock algae *Isochrysis galbana*, but not in *Tetraselmis suecica*

Susan C. Fitzer^a, Julien Plancq^{b,*}, Cameron J. Floyd^b, Faith M. Kemp^b, Jaime L. Toney^b

^a Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK

^b School of Geographical and Earth Sciences, University of Glasgow, Glasgow G12 8QQ, UK

ARTICLE INFO

Keywords:

Ocean acidification
Algae
Lipids
Aquaculture
Feedstock

ABSTRACT

Increased anthropogenic CO_2 emissions are leading to an increase in CO_2 uptake by the world's oceans and seas, resulting in ocean acidification with a decrease in global ocean water pH by as much as 0.3–0.4 units by the year 2100. The direct effects of changing $p\text{CO}_2$ on important microalgal feedstocks are not as well understood. Few studies have focused on lipid composition changes in specific algal species in response to ocean acidification and yet microalgae are an indispensable food source for various marine species, including juvenile shellfish. *Isochrysis galbana* and *Tetraselmis suecica* are widely used in aquaculture as feeds for mussels and other shellfish. The total lipid contents and concentrations of *I. galbana* and *T. suecica* were investigated when grown under present day (400 ppm) and ocean acidification conditions (1000 ppm) to elucidate the impact of increasing $p\text{CO}_2$ on an important algae feedstock. Total lipids, long-chain alkenones (LCAs) and alkenoates decreased at 1000 ppm in *I. galbana*. *I. galbana* produces higher lipids than *T. suecica*, and is perhaps as a result more impacted by the change in carbon available for lipid production under higher $p\text{CO}_2$. *I. galbana* is an important feedstock, more easily assimilated for growth in juvenile shellfish and reductions in lipid composition may prove problematic for the growth of future shellfish aquaculture. Our findings suggest that higher $p\text{CO}_2$ impacts on algal lipid growth are species specific and warrant further study. It is therefore vital to examine the impact of high CO_2 on algal lipid production, especially those commercial shellfish feed varieties to predict future impacts on commercial aquaculture.

1. Introduction

Increased anthropogenic CO_2 emissions are leading to an increase in CO_2 uptake by the world's oceans and seas, resulting in a decrease in global ocean water pH by as much as 0.3–0.4 units by the year 2100 (IPCC, 2013). This decrease in ocean pH, termed ocean acidification, will have severe ramifications for marine organisms, particularly calcifying organisms such as shellfish, mussels and corals, whose calcium carbonate constituents dissolve more readily and grow more slowly in increasingly acidic waters (Berge, Bjerkeng, Pettersen, Schaanning, & Øxnevad, 2006; Tunnicliffe et al., 2009). It has been suggested that the current IPCC 2100 prediction could lead to a decrease in global calcification rates between 5% and 50% (Kleypas et al., 2005). The large variation in this estimate highlights the uncertainty surrounding the impacts of a pH decrease on marine calcifiers.

Studies on the impact of increased levels of partial pressure CO_2 ($p\text{CO}_2$) and ocean acidification on mussel and shellfish have been undertaken in recent years. Gazeau et al. (2007) led an early study on the

mussel *Mytilus edulis* and concluded that there was a 25% reduction in the calcification rate when it was exposed to a $p\text{CO}_2$ of 740 ppm, a level predicted by the IPCC in next century. Further work by Fitzer et al. (2015) found that both the newest and older calcite of *Mytilus edulis* grown under high $p\text{CO}_2$ (1000 ppm) for a period of 6 months were harder and more brittle and hence fractured more easily. These shells would be more susceptible to breakage from industry or predation (Fitzer et al., 2015). Thus, increasing ocean CO_2 has the potential to significantly reduce global mussel and shellfish production. This could have major socio-economic impacts across the globe as it is currently estimated that shellfish account for approximately 23.6% of a \$119.4 billion fish industry (Fitzer, Cusack, Phoenix, & Kamenos, 2014). Research has begun to investigate methods of mitigating against these impacts. It has been suggested by Melzner et al. (2011) that increasing the food supply available to mussels, can limit the potentially devastating effects of increasing ocean acidity. Indeed when mussels were reared in an environment with a high concentration of nutrients in the water column, inner shell corrosion was only visible in the specimens

* Corresponding author. School of Geographical and Earth Sciences, University of Glasgow, Glasgow G12 8QQ, UK

E-mail address: julien.plancq@glasgow.ac.uk (J. Plancq).

<https://doi.org/10.1016/j.aaf.2019.02.008>

Received 19 October 2018; Received in revised form 22 February 2019; Accepted 28 February 2019

Available online 16 March 2019

2468-550X/ © 2019 Shanghai Ocean University. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Table 1Seawater TA, salinity, temperature and $p\text{CO}_2$ were used to calculate other seawater parameters using CO2Sys (Riebesell et al., 2010).

Treatment (ppm)	Salinity	T (°C)	TA ($\mu\text{mol/kgSW}$)	$p\text{CO}_2$ (μatm)	HCO_3^- ($\mu\text{mol/kgSW}$)	CO_3^{2-} ($\mu\text{mol/kgSW}$)	ΩCa	ΩAr	DIC $\delta^{13}\text{C}$ VPDB (per mil)
400	35.5	19	1028 \pm 206	409 \pm 57	900	42	0.99	0.64	−2.02 \pm 0.56
400	35.6	19	1028 \pm 206	434 \pm 69	906	40	0.95	0.61	−1.98 \pm 0.38
1000	37.9	19	1139 \pm 262	1043 \pm 132	1067	24	0.56	0.37	−5.86 \pm 0.57
1000	37.0	19	1139 \pm 262	984 \pm 80	1065	25	0.59	0.38	−4.78 \pm 0.27

subjected to the very highest CO_2 concentrations far exceeding the IPCC predictions (2400 ppm and 4000 ppm). Similarly, Thomsen, Casties, Pansch, Körtzinger, and Melzner (2013) argued that when food supply is abundant, *Mytilus edulis* can withstand and thrive in higher $p\text{CO}_2$ conditions. These studies suggest that supplementary feeding may have positive effects on mussel shell growth when mussels are reared in acidic water conditions, alleviating the negative impacts described by Hettinger et al. (2013) and Fitzer et al. (2015).

While these studies measured the direct effect of changing $p\text{CO}_2$ on important edible shellfish, other important impacts, for instance, on the microalgal feedstocks are not as well understood. Microalgae are an indispensable food source for various marine species, including juvenile molluscs and crustaceans (Dörner, Carbonell, Pino, & Farias, 2014), providing key unsaturated fatty acids that most higher organisms cannot produce themselves. A few studies have focused on lipid composition changes in specific algal species in response to ocean acidification (Riebesell, Revill, Holdsworth, & Volkman, 2000; Rossoll et al., 2012; Tsuzuki, Ohnuma, Sato, Takaku, & Kawaguchi, 1990). For instance, the diatom *Thalassiosira pseudonana* has been shown to decrease the proportion of long-chain polyunsaturated fatty acids relative to saturated fatty acids in response to high $p\text{CO}_2$ (Rossoll et al., 2012). Fatty acids were also shown to be depleted in ^{13}C in the coccolithophore *Emiliania huxleyi* (Riebesell et al., 2000) and depleted fatty acids in cells of *Chlorella vulgaris* was observed in response to high $p\text{CO}_2$ (Tsuzuki et al., 1990). Similar results were found in other algae with respect to a shift to higher proportions of saturated fatty acids, but also accompanied by a shift from nano-plankton (i.e., chlorophytes and haptophytes) to favoring pico-plankton biomass (Bermúdez, Riebesell, Larsen, & Winder, 2016). Typically, aquaculture feedstocks contain a mix of algal species and sizes to satisfy both adult and juvenile molluscs (Beck & Neves, 2003; Gatenby, Parker, & Neves, 1997), hence it is important to understand changes in size classes induced by increasing $p\text{CO}_2$.

Algae size has been linked to improved juvenile growth, however algae in the size range of 2.8–8.5 μm is recommended for best results (Beck & Neves, 2003), although perhaps more importantly higher polyunsaturated fatty acids are required for newly metamorphosed juvenile mussels (Gatenby et al., 1997). A varied diet of mixed algae is thought to improve retention time and greater assimilation efficiency in bivalves, however, there is no evidence to relate particle size to assimilation of energy in *M. edulis* (Shumway, Cucci, Newell, & Yentsch, 1985; Wang & Fisher, 1996). It is suggested that cell wall structure and biochemical composition of algae cells may influence assimilation with a lower efficiency in bivalves from green algae cells due to the difficulty in breaking down sporopollenin (Wang & Fisher, 1996). Therefore there is potential for the juveniles to more easily digest a non-sporopollenin containing *Isochrysis* sp. chosen as a preferential food source for juveniles such as the Veligers of *Mercenaria mercenaria* (Shumway et al., 1985). *Isochrysis galbana* (~5 μm) and *Tetraselmis suecica* (~10 μm) are widely used in aquaculture as feeds for mussels and other calcifying organisms (Cripps, Lindeque, & Flynn, 2014; Fabregas, Herrero, Cabezas, & Abalde, 1985; Parker et al., 2012; Priyadarshani, Sahu, & Rath, 2012; Zittelli, Rodolfi, Biondi, & Tredici, 2006). Limited research has been done on the effects of $p\text{CO}_2$ change on the lipid content of these commercially important algae for shellfish aquaculture and the potential effect in mussel feeding. Here, the total lipid contents and

concentrations of *I. galbana* and *T. suecica* were investigated when grown under present day (400 ppm) and ocean acidification conditions (1000 ppm) to elucidate the impact of increasing $p\text{CO}_2$ on an important algae feedstock.

2. Methods

2.1. Algae cultures

I. galbana and *T. suecica* cultures were grown from stock algae obtained from the Culture Collection of Algae and Protozoa (CCAP) (Strains 927/1 and 66/4 CCAP, respectively), Oban. Both species were cultured at two different levels of $p\text{CO}_2$; 400 ppm representing present day conditions and 1000 ppm representing predicted levels by the year 2100 (IPCC, 2007; Doney, Fabry, Feely, & Kleypas, 2009). To create these mesocosms, CO_2 was bubbled into the samples from gas lines at the desired concentrations. CO_2 was mixed into air lines supplying all experimental cultures and gas concentrations were logged continuously using LI-COR® Li-820 CO_2 gas analysers at 400 and 1000 μatm $p\text{CO}_2$ (Table 1). The concentrations of $p\text{CO}_2$ were recorded daily and the temperature remained constant at 19 °C throughout the experimental period, controlled via air-conditioned laboratory. Seawater samples were collected at the start and end of the experimental culture in air tight rubber-septum glass exetainers and spiked with mercuric chloride (50 μl in 12 ml seawater) (Dickson, Sabine, & Christian, 2007). Total alkalinity (TA) was determined using standard semi-automated titration (Metrohm 848 Titrino plus) of 0.1 Molar (M) hydrochloric acid (HCl) in 0.6 Molar (M) sodium chloride (NaCl) (Dickson et al., 2007), combining the spectrometric analysis using bromocresol indicator (Yao & Byrne, 1998) and the absorbance read through a spectrometer (Hach DR 5000™ UV-Vis) at wavelengths of 444 and 616. Certified seawater references materials for oceanic CO_2 (Batch 156, Scripps Institution of Oceanography, University of California, San Diego) were used as standards to quantify the error of analysis (Measured $2186.39 \pm 4.56 \mu\text{mol/kg}$, CRM value $2234.07 \pm 0.39 \mu\text{mol/kg}$) (Dickson et al., 2007). Dissolved inorganic carbon (DIC) was measured in triplicate for seawater sampled at the start, end of the experimental culture, and reported as $\delta^{13}\text{C}$ VPDB (per mil) (Table 1). The DIC was analysed using an AP 2003 continuous flow automated carbonate system giving results as VPDB $\delta^{13}\text{C}$ (Vienna Pee Dee Belemnite) Marine Carbonate Standard obtained from a Cretaceous marine fossil, *Belemnite Americana* (Scottish Universities Environment Research Centre, SUERC). Seawater TA, salinity, temperature and $p\text{CO}_2$ were used to calculate other seawater parameters using CO2Sys (Riebesell, Fabry, Hansson, & Gattuso, 2010) (Table 1).

Six replicates were used for the different combinations of controlled variables (400 ppm and 1000 ppm $p\text{CO}_2$, *T. suecica* and *I. galbana* species); hence there were 24 samples in total. The samples were each placed in a 250 ml conical flask containing f/2 medium, connected to air supplies using 0.2 μm filters to prevent contamination. To replicate conditions in the field, a light source was set up above the samples and was placed on a timer from 8 a.m. to 6 pm to represent daylight cycles. In order to enable continuous growth in a closed system culture the algae samples were diluted with sterilized media every week, the dilution frequency was determined from growth rate during the first week. F/2 media was prepared and autoclaved (Priorclave TACTROL®2)

Table 2

Average concentrations of the main lipids identified in *Isochrysis galbana* and *Tetraselmis suecica* for each $p\text{CO}_2$ treatment. Values in bold and italic indicate that differences between the two $p\text{CO}_2$ treatments are statistically significant (t -test, $p < 0.05$).

	<i>I. galbana</i> ($\mu\text{g/g}$ algae)		<i>T. suecica</i> ($\mu\text{g/g}$ algae)	
	400 ppm	1000 ppm	400 ppm	1000 ppm
C_{14:0} Fatty acid	28.2	21.1	1.69	1.98
C_{16:0} Fatty acid	42.4	31.9	45.7	44.8
C_{18:1} Fatty acid	34.3	26.2	32.5	27.8
C_{18:0} Fatty acid	11.5	8.72	6.76	7.95
Brassicasterol	64.0	61.8	0.000	0.000
Campesterol	60.8	40.8	60.2	49.1
24-methyl-Cholesta-3,5-triene isomer	70.1	60.6	0.000	0.000
24-methyl-Cholesta-3,5-triene isomer	70.9	65.8	37.5	31.9
Methyl-Cholestatetraene	107.5	80.5	0.000	3.12
24-methyl-Cholesta-3,5-triene isomer	205.2	194.2	35.6	29.8
24-methyl-Cholesta-3,5-diene isomer	55.8	51.7	57.6	47.7
24-methyl-Cholesta-2,5-diene	47.1	21.9	62.4	49.9
Methyl-Cholestatriene	47.7	39.5	80.2	67.7
24-methyl-Cholesta-3,5-diene isomer	52.1	60.2	83.8	70.8
24-methyl-Cholesta-3,5-diene	0.000	0.000	283.9	249.0
24-ethyl-Cholesta-3,5-triene	67.3	53.8	0.000	0.000
Ethyl-Cholestatriene	133.9	94.2	0.000	0.000
24-ethyl-Cholesta-3,5-diene	167.5	149.6	0.000	0.000
C_{31:2} Alkene	225.9	152.2	–	–
C_{31:1} Alkene	115.8	95.2	–	–
C_{37:4} LCA	2246.8	1967.5	–	–
C_{37:3} LCA	10169.5	4439.3	–	–
C_{37:2} LCA	454.9	194.6	–	–
C_{38:4} LCA	995.4	862.8	–	–
C_{38:3} LCA	4172.9	1962.7	–	–
C_{38:2} LCA	849.3	364.8	–	–
C_{39:3} LCA	749.8	388.9	–	–
C_{40:3} LCA	264.2	147.6	–	–
C_{36:2} Alkenoate	154.2	88.4	–	–
C_{36:4} Alkenoate	304.4	354.3	–	–
C_{36:3} Alkenoate	876.0	518.6	–	–
C_{36:2} Alkenoate	1007.6	403.3	–	–
Total Fatty acids	116.3	87.9	86.7	82.6
Total Sterols	1150.1	974.8	701.2	599.1
Total LCAs	19902.8	10328.1	0.000	0.000
Total Alkenoates	2342.3	1364.7	0.000	0.000
Total lipids	24376.5	13414.5	1351.5	1168.9

for sterile preparation following the CCAP f/2 recipe (<https://www.ccap.ac.uk/media/documents/f2.pdf>). 50 ml media was used to replace 50 ml of concentrated algae, which was added to each culture per week in a sterile laminar flow cupboard (Laminar Air Flow Cabinet GELAIRE CLASS 100).

After the 9-week period of experimental culture, microscopic counts were undertaken in order to determine the cell concentrations. A Haussner haemocytometer was used and 4 counts were done for each algae sample. The algal cells in five boxes of the haemocytometer were counted and an average taken, this was then divided by 4×10^{-6} to obtain the concentration of cells/ml. This was repeated for the 4 counts and a mean concentration for each sample was noted.

2.2. Lipid extraction and analysis

Between 25 and 40 ml of each sample was filtered on a pre-weighed glass fibre filter paper and freeze-dried at -55°C and 0.006 Pa. The freeze-dried samples were extracted with dichloromethane (DCM):methanol (MeOH) (9:1, v:v) using a Thermoscientific Dionex ASE350 accelerated solvent extractor (ASE). Prior to analysis by gas chromatography, the total lipid extracts (TLEs) were derivatised with 30 μl of O-Bistrifluoroacetamide (BSTFA) and 40 μl of Pyridine at 80°C for 2 h. Lipids were detected and quantified using gas chromatography with a flame ionization detector (GC-FID). Analyses were performed on an Agilent 7890B Series GC system configured with a Restek RTX-1 fused silica column (60 m length, 0.25 mm i.d., 0.1 μm film thickness). Helium was used as the carrier gas at a flow pressure of 20.8 psi. The GC method used splitless injection, and the oven temperature was programmed from 60°C (hold for 2 min) to 120°C at $30^\circ\text{C}/\text{min}$, then to 330°C at $5^\circ\text{C}/\text{min}$ and held for 15 min. Samples were run with the same temperature program on an Agilent 7890B Series GC coupled with a 5977A GC-MS mass spectrometer (GC-MS) to confirm the identity of the lipids using the known ion chromatograms and by comparison of mass spectral data and GC retention times with published data. Lipids (fatty acids, sterols, alkenes, long-chain alkenones, alkenoates) were quantified from the TLEs using Eicosane ($n\text{-C}_{20}$ alkane) and Hexatriacontane ($n\text{-C}_{36}$ alkane) as external standards. Calibration curves had R^2 values of 0.99 and 0.96 for $n\text{-C}_{20}$ alkane and $n\text{-C}_{36}$ alkane, respectively.

To improve fatty acid identification, TLEs were separated into neutral and acid fractions by elution through a LC-NH₂ SPE column using DCM:isopropyl alcohol (1:1, v:v), followed by ether with 4% acetic acid (v:v). Acid samples were methylated with a Boron trifluoride (BF₃)-methanol solution (14% BF₃ in MeOH) at 70°C for 1 h. The methylated samples were then eluted through silica-gel columns with n -hexane to remove impurities from the methylation reaction, followed by DCM to collect the fatty acid methyl esters (FAMES). The hexane and DCM fractions were then analysed in GC-FID and GC-MS as described above. All the FAMES were contained in the DCM fractions.

3. Results

3.1. Algal concentrations

Average concentrations ranged from 7.69×10^{-6} cells/ml to 9.58×10^{-6} cells/ml for *I. galbana*, and from 2.88×10^{-5} cells/ml to 9.88×10^{-5} cells/ml for *T. suecica*. No significant differences were observed between the concentration of algae cultured at 400 ppm $p\text{CO}_2$ and 1000 ppm $p\text{CO}_2$.

3.2. Lipid composition and concentrations

The lipid profiles (from the TLEs) for each species of algae were not affected by the change of $p\text{CO}_2$ level, which means that the same lipids were observed at 400 ppm $p\text{CO}_2$ and 1000 ppm $p\text{CO}_2$ for *I. galbana* and *T. suecica*.

For *I. galbana*, the main lipids identified were the fatty acids C_{14:0} (myristic acid), C_{16:0} (palmitic acid), C_{18:0} (stearic acid), and C_{18:1} (oleic acid), the sterols brassicasterol and campesterol, and the 24-methyl-Cholesta-3,5-triene and 24-ethyl-Cholesta-3,5-diene. In addition, *I. galbana* presented C_{31:2} and C_{31:1} alkenes, long-chain alkenones (LCAs), and alkenoates. These latter lipids are specific to the Isochrysidale haptophyte clade to which it belongs (Marlowe et al., 1984). For *T. suecica*, the main lipids identified were also the fatty acids C_{14:0} (myristic acid), C_{16:0} (palmitic acid), C_{18:0} (stearic acid), and C_{18:1} (oleic acid). Only the sterol campesterol was identified, whereas the 24-methyl-Cholesta-3,5-diene prevailed. Additional fatty acids were identified in both *I. galbana* and *T. suecica* through acid/neutral separation including the C_{12:0}, C_{13:0}, C_{15:0}, C_{17:0}, C_{18:2}, C_{20:0} and C_{24:0} fatty acids,




Fig. 1. Variations in the content of total lipids and major lipid groups between the two $p\text{CO}_2$ treatments for *Isochrysis galbana* (A1-D1) and *Tetraselmis suecica* (A2-C2). Error bars represent standard deviation. Differences in concentrations that are statistically significant (t -test, $p < 0.05$) between the two $p\text{CO}_2$ treatments are indicated by the red asterisks. Note that the scales are different for each diagram. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

however these were too low in concentration to be quantified and therefore have not been compared between treatments.

The concentrations of the total lipid content (determined from the TLEs) and the major lipids for each species at the different $p\text{CO}_2$ levels are presented in Table 2 (averages of the six replicates for each species and each treatment). In general, *I. galbana* had higher lipid content than *T. suecica*, which is easily explained by the high concentrations of LCAs and alkenoates (Table 2). When comparing between the different $p\text{CO}_2$ levels, a decrease in concentrations is observed at 1000 ppm for each species for the total lipids, fatty acids, sterols, LCAs and alkenoates (Fig. 1). Since the concentrations showed large standard deviations (error bars in Fig. 1), a t -test was performed to evidence changes in concentrations that are statistically significant between 400 ppm and 1000 ppm. Only the concentrations of total lipids, LCAs and alkenoates for *I. galbana* proved to be significantly ($p < 0.05$) lower at 1000 ppm (Fig. 1; Table 1). When specifically looking at lipids, a decrease in concentrations is also observed at 1000 ppm for each species (Fig. 2). However, for *I. galbana*, only most of the LCAs and alkenoates showed a significant ($p < 0.05$) concentration decrease (Fig. 2; Table 2). For *T. suecica*, only the campesterol significantly decreased (Fig. 2; Table 2).

In summary, for *I. galbana*, a significant concentration decrease was observed at 1000 ppm for the total lipids, LCAs and alkenoates, while for *T. suecica*, only the campesterol significantly decreased.

3.3. Seawater chemistry

The seawater chemistry data outlines changes to the $p\text{CO}_2$ and resulting changes to the DIC and carbonate (CO_3^{2-}). The carbonate and

DIC is lighter in the seawater at 1000 ppm $p\text{CO}_2$ compared with the 400 ppm $p\text{CO}_2$ (Table 1).

4. Discussion

The nutritional value of algae is related to its fatty acid and lipid compositions, with the most important aspect being the proportion of polyunsaturated fatty acids (PUFAs) (Fidalgo, Cid, Torres, Sukenik, & Herrero, 1998). It has been stated in previous research that the profile and quantity of fatty acid algal lipids is crucial to the development and growth of certain marine fauna (Zhu, Lee, & Chao, 1997). While in the present study we identified fatty acids not previously reported (for example $\text{C}_{12:0}$, $\text{C}_{13:0}$, $\text{C}_{15:0}$, $\text{C}_{17:0}$, $\text{C}_{20:0}$ and $\text{C}_{24:0}$ fatty acids), other studies have reported more fatty acids and especially PUFAs (such as $\text{C}_{18:3}$, $\text{C}_{18:4}$, $\text{C}_{20:4}$, $\text{C}_{20:5}$ and $\text{C}_{22:6}$ PUFAs) for *I. galbana* and *T. suecica* (e.g., Custódio et al., 2014; Servel, Claire, Derrien, Coiffard, & De Roeck-Holtzhauer, 1994). These discrepancies in the fatty acid composition are certainly due to the study of different algal strains coming from different Culture Collections and/or different culture conditions. Fatty acid content and distributions can be strongly affected by environmental conditions such as differing nutrient, irradiance, salinity conditions, and growth stage as shown by culture studies (Dunstan, Volkman, Barrett, & Garland, 1993; Martinez-Roldan, Perales-Vela, Canizares-Villanueva, & Torzillo, 2014; Pal, Khozin-Goldberg, Cohen, & Boussiba, 2011). We note that Servel et al. (1994) studied *I. galbana* and *T. suecica* strains from an aquaculture centre of Loire Atlantique (France), whereas our strains are from the Culture Collection of Oban (Scotland). However, since we studied the same algae grown under




Fig. 2. Variations in the content of specific lipids between the two $p\text{CO}_2$ treatments for *Isochrysis galbana* (A1-C1) and *Tetraselmis suecica* (A2-B2). Error bars represent standard deviation. Differences in concentrations that are statistically significant (t -test, $p < 0.05$) between the two $p\text{CO}_2$ treatments are indicated by the red asterisks. Note that the scales are different for each diagram. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

different CO_2 levels, we do believe that these discrepancies with previous studies in the fatty acid composition do not affect the interpretations/conclusions of the present work.

The correlation between increased lipid content and growth rate in calcifiers has been documented in numerous studies. Parrish, Wells, Yang, and Dabinett (1999) found that there was a significant increase in growth of scallop juveniles, *Placopecten magellanicus*, when fed *I. galbana* with a higher lipid content with the addition of the algae *Chaetoceros muelleri* compared to *I. galbana* with a lower lipid content along with *C. muelleri*. The importance of PUFAs is also reiterated as the higher lipid *I. galbana* alone did not affect the growth rate of the juvenile scallops (Parrish et al., 1999). Parrish et al. (1999) suggest that because *C. muelleri* had much more of the essential PUFA $\text{C}_{20:4}$ than *I. galbana* this supplement was needed to reach the threshold required for growth.

The fact that concentrations of the total lipids, fatty acids, sterols, LCAs and alkenoates decreased at 1000 ppm in *I. galbana* contradicts biofuel studies examining lipid composition under experimental rapid growth in higher CO_2 (Nakanishi et al., 2014; Singh & Singh, 2014; Tanadul, Vanderghenst, Beckles, Powell, & Labavitch, 2014). The role of sterols and alkenones for bivalve growth are not well understood (Soudant et al., 1998). For the Scallop *Pecten maximus*, ingested sterols cholesterol and stigmasterol were preferentially incorporated by the larvae, potentially indicating the sterols as essential for growth (Soudant et al., 1998). Ocean acidification research into the diatom *T. pseudonana* has similarly observed a decrease in the proportion of long-chain polyunsaturated fatty acids relative to saturated fatty acids in

response to high $p\text{CO}_2$ (Rossoll et al., 2012), similarly to the depleted fatty acids in the *E. huxleyi* under high $p\text{CO}_2$ (Riebesell et al., 2000), and in agreement with the *I. galbana* lipids of this study. Examining the carbon isotopes of the seawater might explain why lower lipid production occurs when higher carbon-13 ($\delta^{13}\text{C}$) incorporation has been observed in monosaccharides and lipids of aquatic plants (van Dongen, Schouten, & Sinninghe Damsté, 2002), where the $\delta^{13}\text{C}$ is lighter as a result of higher $p\text{CO}_2$ at 1000 ppm, the algae are unable to incorporate the $\delta^{13}\text{C}$ for lipid production. When comparing the two studied species, *I. galbana* produces higher lipids than *T. suecica*, linked in part to the production of alkenones, and is perhaps as a result more impacted by the change in carbon available for lipid production under higher $p\text{CO}_2$. Higher lipid contents (as total lipid content as percentage of dry mass; TLDM) have been observed previously in *I. galbana* compared *T. suecica* (Servel et al., 1994). The two species of microalgae were chosen for this study as commercially important species for shellfish aquaculture, and are widely used in aquaculture as feeds for mussels and other calcifying organisms (Cripps et al., 2014; Fabregas et al., 1985; Parker et al., 2012; Priyadarshani et al., 2012; Zittelli et al., 2006). *I. galbana* is particularly important as it is more easily assimilated for growth in bivalves, particularly juveniles such as the clam Veligers of *M. mercenaria* (Shumway et al., 1985). We show in this study that higher $p\text{CO}_2$ reduces the lipid production, specifically total lipids, fatty acids, LCAs and alkenoates for *I. galbana*, which may affect shellfish growth in aquaculture. Our findings, and the findings of others (Rossoll et al., 2012) suggest that higher $p\text{CO}_2$ impacts on algal lipid growth are species specific and warrant further study. In commercial hatchery shellfish aquaculture

algae are mass produced often using high CO₂ to enhance growth rates. Additionally, in field shellfish aquaculture where climate change is increasing the CO₂ within seawater naturally available algae species may alter their lipid composition affecting shellfish growth. The findings of this study suggest that lipid production in *I. galbana* is significantly reduced under higher pCO₂. Further studies are therefore necessary to examine the impact of high CO₂ on algal lipid production, especially those commercial shellfish feed varieties which are used in 6 mix algal feeds for aquaculture (e.g. Shellfish Diet 1800® a mix of six marine microalgae - *Isochrysis*, *Pavlova*, *Tetraselmis*, *Chaetoceros calcitrans*, *Thalassiosira weissflogii* and *Thalassiosira pseudonana* http://www.reedmariculture.com/product_instant_algae_shellfish_diet_1800.php).

5. Conclusions

In this study, total lipids, LCAs and alkenoates decreased at 1000 ppm in *I. galbana*. *I. galbana* produces higher lipids than *T. suecica*, and is perhaps as a result more impacted by the change in carbon available for lipid production under higher pCO₂. *I. galbana* is an important feedstock, more easily assimilated for growth in juvenile shellfish and reductions in lipid composition may prove problematic for the growth of future shellfish aquaculture. Our findings suggest that higher pCO₂ impacts on algal lipid growth are species specific and warrant further study. It is therefore vital to examine the impact of high CO₂ on algal lipid production, especially those commercial shellfish feed varieties to predict future impacts on commercial aquaculture.

Acknowledgments

We thank Mohammad Ali Salik for laboratory and technical assistance. SF would like to acknowledge funding from NERC Independent Research Fellowship [NE/N01409X/1]. We thank two anonymous reviewers for their constructive comments that helped to improve the quality of the manuscript.

References

- Beck, K., & Neves, R. J. (2003). An evaluation of selective feeding by three age-groups of the rainbow mussel *Villosa iris*. *North American Journal of Aquaculture*, 65, 203–209. <https://doi.org/10.1577/C02-031>.
- Berge, J. A., Bjerkeng, B., Pettersen, O., Schaanning, M. T., & Øxnevad, S. (2006). Effects of increased seawater concentrations of CO₂ in growth of the bivalve, *Mytilus edulis* L. *Chemosphere*, 62, 681–687. <https://doi.org/10.1016/j.chemosphere.2005.04.111>.
- Bermúdez, J. R., Riebesell, U., Larsen, A., & Winder, M. (2016). Ocean acidification reduces transfer of essential biomolecules in a natural plankton community. *Scientific Reports*, 6, 27749. <https://doi.org/10.1038/srep27749>.
- Cripps, G., Lindeque, P., & Flynn, K. J. (2014). Have we been underestimating the effects of ocean acidification in zooplankton? *Global Change Biology*, 20, 3377–3385. <https://doi.org/10.1111/gcb.12582>.
- Custódio, L., Soares, F., Pereira, H., Barreira, L., Vizetto-Duarte, C., Rodrigues, M. J., et al. (2014). Fatty acid composition and biological activities of *Isochrysis galbana* T-ISO, *Tetraselmis* sp. and *scenedesmus* sp.: Possible application in the pharmaceutical and functional food industries. *Journal of Applied Phycology*, 26, 151–161. <https://doi.org/10.1007/s10811-013-0098-0>.
- Dickson, A. G., Sabine, C. L., & Christian, J. R. (2007). *Guide to best practises for ocean CO₂ measurements*. Sidney, British Columbia: North Pacific Marine Science Organization PICES Special Publication 3.
- Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). ocean acidification: The other CO₂ problem. *Annual Review of Marine Science*, 1, 169–192. <https://doi.org/10.1146/annurev.marine.010908.163834>.
- van Dongen, B. E., Schouten, S., & Sinninghe Damsté, J. S. (2002). Carbon isotope variability in monosaccharides and lipids of aquatic algae and terrestrial plants. *Marine Ecology Progress Series*, 232, 83–92. <https://doi.org/10.3354/meps232083>.
- Dörner, J., Carbonell, P., Pino, S., & Farias, A. (2014). Variation of fatty acids in *Isochrysis galbana* (T-Iso) and *Tetraselmis suecica*, cultured under different nitrate availabilities. *Journal of Fisheries and Aquaculture*, 5, 1–3.
- Dunstan, G. A., Volkman, J. K., Barrett, S. M., & Garland, C. D. (1993). Changes in the lipid composition and maximisation of the polyunsaturated fatty acid content of three microalgae grown in mass culture. *Journal of Applied Phycology*, 5, 71–83. <https://doi.org/10.1007/BF02182424>.
- Fabregas, J., Herrero, C., Cabezas, B., & Abalde, J. (1985). Mass culture and biochemical variability of the marine microalga *Tetraselmis suecica* Kylin (Butch) with high nutrient concentrations. *Aquaculture*, 49, 231–244. [https://doi.org/10.1016/0044-8486\(85\)90082-1](https://doi.org/10.1016/0044-8486(85)90082-1).
- Fidalgo, J. P., Cid, A., Torres, E., Sukenik, A., & Herrero, C. (1998). Effects of nitrogen source and growth phase on proximate biochemical composition, lipid classes and fatty acid profile of the marine microalga *Isochrysis galbana*. *Aquaculture*, 166, 105–116. [https://doi.org/10.1016/S0044-8486\(98\)00278-6](https://doi.org/10.1016/S0044-8486(98)00278-6).
- Fitzer, S. C., Cusack, M., Phoenix, V. R., & Kamenos, N. A. (2014). Ocean acidification reduces the crystallographic control in juvenile mussel shells. *Journal of Structural Biology*, 188, 39–45. <https://doi.org/10.1016/j.jsb.2014.08.007>.
- Fitzer, S. C., Zhu, W., Tanner, K. E., Kamenos, N. A., Phoenix, V. R., & Cusack, M. (2015). Ocean acidification alters the material properties of *Mytilus edulis* shells. *Journal of The Royal Society Interface*, 12, 20141227. <https://doi.org/10.1098/rsif.2014.1227>.
- Gatenby, C. M., Parker, B. C., & Neves, R. J. (1997). Growth and survival of juvenile rainbow mussels, *Villosa iris* (Bivalvia: Unionidae) reared on algal diets and sediment. *American Malacological Bulletin*, 14, 57–66.
- Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J.-P., Middelburg, J. J., & Heip, C. H. R. (2007). Impact of elevated CO₂ on shellfish calcification. *Geophysical Research Letters*, 34, L07603. <https://doi.org/10.1029/2006GL028554>.
- Hettinger, A., Sanford, E., Hill, T. M., Hosfelt, J. D., Russell, A. D., & Gaylord, B. (2013). The influence of food supply on the response of Olympia oyster larvae to ocean acidification. *Biogeosciences*, 10, 6629–6638. <https://doi.org/10.5194/bg-10-6629-2013>.
- IPCC (2007). Summary for policymakers. In S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, & K. B. Averyt, (Eds.). *Climate change 2007: The physical science basis. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- IPCC (2013). Summary for policymakers. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, & J. Boschung, (Eds.). *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- Kleypas, J. A., Buddemeier, R. W., Eakin, C. M., Gattuso, J. P., Guinotte, J., Hoegh-Guldberg, O., et al. (2005). Comment on “Reef calcification and climate change: The effect of ocean warming”. *Geophysical Research Letters*, 32, 1–3. <https://doi.org/10.1029/2004GL022329>.
- Marlowe, I. T., Green, J. C., Neal, A. C., Brassell, S. C., Eglinton, G., & Course, P. A. (1984). Long chain (n-C₃₇–C₃₉) alkenones in the Prymnesiophyceae. Distribution of alkenones and other lipids and their taxonomic significance. *British Phycological Journal*, 19, 203–216. <https://doi.org/10.1080/00071618400650221>.
- Martinez-Roldan, A. J., Perales-Vela, H. V., Canizares-Villanueva, R. O., & Torzillo, G. (2014). Physiological response of *Nannochloropsis* sp. to saline stress in laboratory batch cultures. *Journal of Applied Phycology*, 26, 115–121. <https://doi.org/10.1007/s10811-013-0060-1>.
- Melzner, F., Stange, P., Trubenbach, K., Thomsen, J., Casties, I., Panknin, U., et al. (2011). Food supply and seawater pCO₂ impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*. *PLoS One*, 6, e24223. <https://doi.org/10.1371/journal.pone.0024223>.
- Nakanishi, A., Aikawa, S., Ho, S.-H., Chen, C.-Y., Chang, J.-S., Hasunuma, T., et al. (2014). Development of lipid productivities under different CO₂ conditions of marine microalgae *Chlamydomonas* sp. JSC4. *Bioresource Technology*, 152, 247–252. <https://doi.org/10.1016/j.biortech.2013.11.009>.
- Pal, D., Khozin-Goldberg, I., Cohen, Z., & Boussiba, S. (2011). The effect of light, salinity, and nitrogen availability on lipid production by *Nannochloropsis* sp. *Applied Microbiology and Biotechnology*, 90, 1429–1441. <https://doi.org/10.1007/s00253-011-3170-1>.
- Parker, L., Ross, P., O'Connor, W., Borysko, L., Raftos, D., & Pörtner, H. (2012). Adult exposure influences offspring response to ocean acidification in oysters. *Global Change Biology*, 18, 82–92. <https://doi.org/10.1111/j.1365-2486.2011.02520.x>.
- Parrish, C. C., Wells, J. S., Yang, Z., & Dabinett, P. (1999). Growth and lipid composition of scallop juveniles, *Placochlamys magellanicus*, fed the flagellate *Isochrysis galbana* with varying lipid composition and the diatom *Chaetoceros muelleri*. *Marine Biology*, 133, 461–471. <https://doi.org/10.1007/s002270050486>.
- Priyadarshani, I., Sahu, D., & Rath, B. (2012). Algae in aquaculture. *International Journal of Medical and Health Sciences Research*, 2, 108–114.
- Riebesell, U., Fabry, V. J., Hansson, L., & Gattuso, J.-P. (2010). *Guide to best practices for ocean acidification research and data reporting*. Luxembourg: Publications Office of European Union.
- Riebesell, U., Revill, A. T., Holdsworth, D. G., & Volkman, J. K. (2000). The effects of varying CO₂ concentration on lipid composition and carbon isotope fractionation in *Emiliania huxleyi*. *Geochimica et Cosmochimica Acta*, 64, 4179–4192.
- Rossell, D., Bermudez, R., Hauss, H., Schulz, K. G., Riebesell, U., Sommer, U., et al. (2012). Ocean acidification-induced food quality deterioration constrains trophic transfer. *PLoS One*, 7, e34737. <https://doi.org/10.1371/journal.pone.0034737>.
- Servel, M.-O., Claire, C., Derrien, A., Coiffard, L., & De Roeck-Holtzauer, Y. (1994). Fatty acid composition of some marine microalgae. *Phytochemistry*, 36, 691–693. [https://doi.org/10.1016/S0031-9422\(00\)89798-8](https://doi.org/10.1016/S0031-9422(00)89798-8).
- Shumway, S. E., Cucci, T. L., Newell, R. C., & Yentsch, C. M. (1985). Particle selection, ingestion, and absorption in filter-feeding bivalves. *Journal of Experimental Marine Biology and Ecology*, 91, 77–92. [https://doi.org/10.1016/0022-0981\(85\)90222-9](https://doi.org/10.1016/0022-0981(85)90222-9).
- Singh, S. P., & Singh, P. (2014). Effect of CO₂ concentration on algal growth: A review. *Renewable and Sustainable Energy Reviews*, 38, 172–179. <https://doi.org/10.1016/j.rser.2014.05.043>.
- Soudant, P., Le Coz, J.-R., Marty, Y., Moal, J., Robert, R., & Samain, J.-F. (1998). Incorporation of microalgae sterols by scallop *Pecten maximus* (L.) larvae. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 119, 451–457. [https://doi.org/10.1016/S1095-6433\(97\)00451-0](https://doi.org/10.1016/S1095-6433(97)00451-0).
- Tanadul, O.-u.-m., Vanderghenst, J. S., Beckles, D. M., Powell, A. L. T., & Labavitch, J.

- M. (2014). The impact of elevated CO₂ concentration on the quality of algal starch as a potential biofuel feedstock. *Biotechnology and Bioengineering*, 111, 1323–1331. <https://doi.org/10.1002/bit.25203>.
- Thomsen, J., Casties, I., Pansch, C., Körtzinger, A., & Melzner, F. (2013). Food availability outweighs ocean acidification effects in juvenile *Mytilus edulis*: Laboratory and field experiments. *Global Change Biology*, 19, 1017–1027. <https://doi.org/10.1111/gcb.12109>.
- Tsuzuki, M., Ohnuma, E., Sato, N., Takaku, T., & Kawaguchi, A. (1990). Effects of CO₂ concentration during growth on fatty acid composition in microalgae. *Plant Physiology*, 93, 851–856.
- Tunnicliffe, V., Davies, K. T. A., Butterfield, D. A., Embley, R. W., Rose, J. M., & Chadwick, W. W., Jr. (2009). Survival of mussels in extremely acidic waters on a submarine volcano. *Nature Geoscience*, 2, 344–348. <https://doi.org/10.1038/ngeo500>.
- Wang, W. X., & Fisher, N. S. (1996). Assimilation of trace elements and carbon by the mussel *Mytilus edulis*: Effects of food composition. *Limnology & Oceanography*, 41, 197–207. <https://doi.org/10.4319/lo.1996.41.2.0197>.
- Yao, W. S., & Byrne, R. H. (1998). Simplified seawater alkalinity analysis: Use of linear array spectrometers. *Deep-Sea Research Part I Oceanographic Research Papers*, 45, 1383–1392. [https://doi.org/10.1016/S0967-0637\(98\)00018-1](https://doi.org/10.1016/S0967-0637(98)00018-1).
- Zhu, C. J., Lee, Y. K., & Chao, T. M. (1997). Effects of temperature and growth phase on lipid and biochemical composition of *Isochrysis galbana* TK1. *Journal of Applied Phycology*, 9, 451–457. <https://doi.org/10.1023/A:1007973319348>.
- Zittelli, G. C., Rodolfi, L., Biondi, N., & Tredici, M. R. (2006). Productivity and photosynthetic efficiency of outdoor cultures of *Tetraselmis suecica* in annular columns. *Aquaculture*, 261, 932–943. <https://doi.org/10.1016/j.aquaculture.2006.08.011>.