

Higher dietary micronutrients are required to maintain optimal performance of Atlantic salmon (*Salmo salar*) fed a high plant material diet during the full production cycle

Luisa M. Vera¹, Kristin Hamre², Marit Espe², Gro-Ingunn Hemre², Kaja Skjærven², Erik-Jan Lock², Antony Jesu Prabhu², Daniel Leeming³, Herve Migaud¹, Douglas R. Tocher¹, John F. Taylor^{1*}

¹ Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling, UK

² Institute of Marine Research, Nordnes, 5817 Bergen, Norway

³ BioMar Ltd., North Shore Road, Grangemouth FK9 8UL, Scotland, UK

*Correspondence: E-mail: j.f.taylor@stir.ac.uk (J. Taylor)

Tel: 00-44-01786 467929

Accepted refereed manuscript of: Vera LM, Hamre K, Espe M, Hemre G, Skjærven K, Lock E, Prabhu AJ, Leeming D, Migaud H, Tocher DR & Taylor JF (2020) Higher dietary micronutrients are required to maintain optimal performance of Atlantic salmon (*Salmo salar*) fed a high plant material diet during the full production cycle. *Aquaculture*, 528, Art. No.: 735551. <https://doi.org/10.1016/j.aquaculture.2020.735551>

© 2020, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

16 **Abstract**

17 A full life cycle (parr to harvest) study of growth and performance was conducted in Atlantic salmon
18 fed diets high in plant ingredients supplemented with graded levels of a nutrient premix (NP),
19 containing selected amino acids, taurine, cholesterol, vitamins and minerals to re-evaluate current
20 nutrient recommendations. Triplicate groups were fed one of three NP levels included at 1x, 2x and
21 4x, where 1x corresponds to recommendations of NRC (2011). Whole body and specific tissue
22 concentrations of nutrients were monitored throughout the experiment as requirement markers.
23 Growth in parr was significantly enhanced in 2xNP, but restricted in 4xNP, while in post-smolts
24 growth was positively correlated with NP level. Spinal deformity decreased linearly with increased
25 NP level in both smolt and post-smolts. When fishmeal and fish oil are present at very low levels, as
26 in the present study, we found beneficial effects with moderate increased levels of the B-vitamins
27 niacin, riboflavin and cobalamin. Further, vitamin C should be increased, based on metabolic
28 responses, although it did not influence growth. Increased Zn and Se affected fish metabolism in a
29 positive manner. Alterations in hepatic transcriptome profiles and expression of specific genes of
30 metabolic pathways were evident in response to micronutrient supplementation level. Collectively,
31 increasing the levels of the micro-nutrient package to a too high level, showed a negative effect and
32 cannot be recommended. NRC (2011) recommendations should therefore be revised for diets in which
33 plant ingredients form the major part of the formulation.

34

35 **Keywords:** Fishmeal, fish oil, plant proteins, vegetable oil, vitamins, minerals, skeletal deformity,
36 sustainable feeds.

37 **1. Introduction**

38 Over the last decade and a half, feed formulations for farmed Atlantic salmon (*Salmo salar*) have
39 changed with marine raw materials, fish oil (FO) and fishmeal (FM), being increasingly replaced by
40 high levels of plant ingredients (Gatlin III et al., 2007; Hardy, 2010; Turchini et al., 2011). Levels of
41 marine ingredients, FM and FO, in salmon feeds are now approximately 20 % of total feed inclusion
42 compared with 90 % around 20 years ago (Ytrestøyl et al., 2015). Over this period, many studies
43 investigated the effects of the inclusion of plant-derived ingredients in feeds. The majority of studies
44 focussed on the impacts on growth performance and feed utilisation (Torstensen et al., 2000; Opstvedt
45 et al., 2003; Mundheim et al., 2004; Espe et al., 2006; Torstensen et al., 2005; Pratoomyot et al., 2008;
46 Waagbø et al., 2013), and also on product quality for human consumers (Waagbø et al., 1993;
47 Mundheim et al., 2004; Torstensen et al., 2005; Menoyo et al., 2007). However, overall, the
48 development and application of modern, plant-based feed formulations has been largely successful
49 enabling salmon production to be maintained and/or increased despite the finite and limited supplies
50 of FM and FO (Shepherd & Jackson, 2013).

51 Despite the above, formulating salmon feeds with high levels of plant meals and vegetable
52 oils significantly affects a range of nutrients (Sissener et al., 2013), including changing micronutrient
53 concentrations and their chemical forms, as well as introducing compounds that interact with
54 micronutrient uptake and metabolism (Olsvik et al., 2013). Importantly, in addition to affecting fish
55 growth, feed efficiency and product quality, these changes in nutrient composition could also affect
56 fish health and welfare (Oxley et al., 2005; Seierstad et al., 2005; Waagbø, 2006; Hemre & Sandnes,
57 2008; Waagbø, 2008; Seierstad et al., 2009). As a result, there has been increasing interest in
58 improving knowledge of practical nutrient requirements of Atlantic salmon when fed modern, plant
59 based feeds (Hansen et al., 2015). In particular, there has been a focus on micronutrients (minerals,
60 vitamins etc.) as it is believed that these were likely supplied in surplus in feeds containing high levels
61 of FM and FO, and that levels of some micronutrients in terrestrial raw materials can be lower
62 (Sissener et al., 2013). When feeding plant-based diets enriched with the micronutrients folate,

63 vitamin B12, vitamin B6, methionine and choline to zebrafish (*Danio rerio*) over the entire life cycle,
64 fish responded with increased growth, higher fecundity and altered metabolic profile of several N-
65 metabolites, indicating an overall positive effect of a micronutrient-enriched plant-based diet
66 (Skjaerven et al., 2016). As a consequence, micronutrient and mineral requirements need careful re-
67 evaluation when substituting raw materials, as the current established micronutrient supplements /
68 premixes (National Research Council, NRC 2011) may not be sufficient to satisfy requirements for
69 Atlantic salmon, especially throughout the whole life cycle.

70 To investigate this, micronutrient requirements of Atlantic salmon fed diets formulated with
71 high levels of plant meals and vegetable oil were studied in two regression experiments in parr in
72 freshwater and post-smolts in seawater (Hamre et al., 2016; Hemre et al., 2016; Prabhu et al. 2019a).
73 A nutrient package, consisting of vitamins B, C, D₃, E, minerals, cholesterol, taurine and histidine
74 was fed at various levels from below NRC (2011) recommended levels to well above NRC levels.
75 Growth, health and welfare parameters responded to nutrient packages in parr, but not to the same
76 extent in post-smolts (Hemre et al., 2016). Thus, parr fed diets containing the nutrient package
77 supplying micronutrients above NRC (2011) recommended levels showed improved protein retention,
78 and reduced liver and viscera indices, whereas the response was less pronounced in post-smolt
79 (Hemre et al., 2016). Significant regressions were obtained in body compartments for vitamin C and
80 several of the B-vitamins and, based on these results, the authors recommended that B-vitamin
81 supplements should be adjusted in plant-based diets for Atlantic salmon (Hamre et al., 2016; Hemre
82 et al., 2016). There were no indications of vitamin E deficiency when vitamin E was not supplemented,
83 indicating that the ingredients delivered sufficient amounts of vitamin E. Nevertheless, it was
84 recommended to supplement 150 mg kg⁻¹ vitamin E to compensate in periods of oxidative stress
85 (Hamre et al., 2016). Regarding the micro-minerals, requirements of Atlantic salmon fed the plant-
86 based diet were met at NP inclusion of 100-150 % and 150-200 % in parr and post-smolt, respectively
87 (Prabhu et al., 2019a).

88 More recently, the impact of micronutrients was investigated in both diploid and triploid
89 Atlantic salmon parr fed graded levels of a nutrient package (NP) from around 30 g to seawater
90 transfer (Taylor et al., 2019). The diets were formulated with very low levels of marine ingredients
91 and supplemented with three levels of NP (L1, 100 %; L2, 200 % and L3, 400 % NP), which contained
92 24 micronutrients including selected amino acids, vitamins, minerals, taurine and cholesterol. The
93 NP was based on the minimum nutrient recommendations for Atlantic salmon (NRC, 2011). Diploid
94 parr fed L2 showed significantly improved growth and reduced liver (HSI) and viscera (VSI) indices,
95 and hepatic steatosis, compared to fish fed L1, while diploids fed L3 also showed improved growth
96 in the first 14 weeks, although growth rate was subsequently reduced (Taylor et al., 2019). In contrast,
97 dietary NP level had less effect on triploid growth rate, VSI or HSI, and smoltification was not
98 affected by NP in either ploidy. Overall, results suggested that, while micromineral requirements
99 were met, other micronutrients may require to be supplemented above current NRC (2011)
100 recommendations for optimal growth and liver function of diploid Atlantic salmon fed plant-based
101 diets in freshwater (Taylor et al., 2019). In addition, the above study also investigated the impact of
102 NP level on skeletal deformity and bone health including the prevalence, localisation and pathology
103 of spinal malformation, and vertebral expression of bone biomarker genes (Vera et al., 2019). This
104 study showed that prevalence of radiologically detectable spinal deformities decreased with
105 increasing micronutrient supplementation when fed from parr to smolt. Concomitantly, expression of
106 many osteogenic genes (bone morphogenic protein, osteocalcin, alkaline phosphatase, matrix
107 metalloproteinase, insulin-like growth factor) increased with increasing NP inclusion. The
108 observations suggested that components thereof, or interactions between micronutrients within the
109 NP may be affecting downstream processes involved in bone formation and remodelling, which also
110 occur through active epigenetic regulatory mechanisms of gene expression (Saito et al., *unpublished*
111 *data*).

112 The previous trials investigating dietary micronutrients in salmon were all carried out during
113 the freshwater phase other than one short-term (5-month) trial in post-smolt in seawater (Hamre et

114 al., 2016; Hemre *et al.*, 2016; Taylor *et al.*, 2019; Vera *et al.*, 2019; Prabhu, *et al.*, 2019a). Therefore,
115 the aim of the present study was to determine the influence of micronutrients in Atlantic salmon fed
116 diets formulated with very low levels of marine ingredients over the entire life cycle. To this end,
117 diploid Atlantic salmon were fed from around 30 g to 3 Kg with plant-based feeds supplemented with
118 the same NP as described above to produce three experimental diets (L1, 100 %; L2, 200 % and L3,
119 400 % NP). The impacts of NP levels on growth, feed efficiency, nutrient retention, skeletal
120 development and liver gene expression were determined.

121

122 **2. Methods & Materials**

123 **2.1 Fish and feeds**

124 The trial was carried out with Atlantic salmon obtained from SalmoBreed AS (Norway). All
125 experimental procedures and husbandry practices were conducted in compliance with the Animals
126 Scientific Procedures Act 1986 (Home Office Code of Practice) in accordance with EU regulation
127 (EC Directive 86/609/EEC) and approved by the Animal Welfare and Ethical Review Board
128 (AWERB) of the University of Stirling. All fish were monitored daily by the Named Animal Care
129 and Welfare Officer (NACWO).

130 Feeds were formulated to reflect standard practice in commercial salmon feeds in terms of
131 protein, oil and energy contents. Thus, feeds were initially formulated to contain 48 % protein and
132 20 % lipid (~22 MJ), with protein content decreasing and lipid content increasing with increasing
133 pellet size to reach 36 % protein and 34 % lipid (~24 MJ) in the largest pellet size in seawater. The
134 experimental feeds were a low FM/FO formulation (initially 15 % and 8 % in freshwater, decreasing
135 to 5% and 3 % respectively in seawater, Tables 1 & 2). Feeds were supplemented with a nutrient
136 package (NP, Tables 3 and 4) at one of three inclusion levels to produce 3 dietary treatments: L1,
137 100 % NP; L2, 200 % NP; L3, 400 % NP (Table 4), the assumption being that the 100 % NP package
138 should contain 100 % of assumed requirement based on the levels reported for Atlantic salmon at the
139 time (NRC, 2011). Specifically, the NP contained 24 nutrients in total these being; vitamins (A, D3,

140 E, K3, C, thiamine, riboflavin, B6, B12, niacin, pantothenic acid, folic acid and biotin), minerals (Ca,
141 Co, I, Se, Fe, Mn, Cu and Zn), crystalline amino acids (L-histidine and taurine) and cholesterol. Total
142 and available phosphorus were fixed in all diets at 12.0 and 9.0 g kg⁻¹ respectively, and magnesium
143 at 1.5 g kg⁻¹, and were not part of the NP. Pellet size (2 mm, 3.5 mm, 5 mm, 7 mm, 9 mm) was
144 adjusted according to fish weight. All non-oil ingredients were mixed and pellets produced by
145 extrusion to produce three base pellets that had oil added by vacuum coating. All feeds were produced
146 at the BioMar Tech-Centre (Brande, Denmark).

147

148 **2.2 Feeding trial**

149 The freshwater phase was carried out at the Niall Bromage Freshwater Research Facility (Stirlingshire,
150 UK). Initially, 500 diploid salmon pre-smolt (initial mean weight, 38.2 ± 0.6 g) were stocked (2 Sept-
151 14) into nine x 1.6 m³ circular fibreglass tanks (3 tanks / diet, initial stocking density 12.0 ± 0.1 kg
152 m⁻³). Fish were acclimated to the experimental tanks for 2 weeks (Fed BioMar Inico Plus) before
153 being fed the experimental diets. Fish were fed continuously during the light period of the light:dark
154 cycle by automatic feeders (Arvotec T2000, Arvotec, Finland) controlled by a PC system. Specific
155 feeding rates (SFR; % tank biomass per day) were adjusted automatically according to predicted
156 growth and daily temperature. An out-of-season photoperiod (LL – 400 °days LD14:10 – 400 °days
157 LL) and ambient water temperature (12 - 16 °C) was applied to produce S0+ smolts (final stocking
158 density 20.2 – 22.8 kg m⁻³), with lighting provided by two 28 W fluorescent daylight bulbs (4000 °K;
159 RS Components, UK) mounted centrally within the tank lid. Water was supplied by an upstream
160 reservoir under flow-through conditions (10 L min⁻¹). Oxygen levels were always higher than 8 mg
161 L⁻¹. Uneaten feed recovery was not feasible during the freshwater phase.

162 Smolts were transferred (4 Nov-14) to the Marine Harvest (Mowi) Feed Trial Unit (Ardnish,
163 Scotland) and on-grown for 11 months in nine 5x5x5 m sea pens under natural photoperiod and
164 ambient water temperatures ranging from 6 to 16 °C. Triplicate groups of 250 post-smolts from
165 respective tanks were stocked per pen (Initial stocking density 0.17 – 0.19 kg m⁻³) and on-grown to

166 a final size of ~2.5 Kg (15 Sept-15, Final stocking density 4.52 – 5.18 kg m⁻³). Fish were fed
167 continuously during daylight by automatic feeders (Arvotec T2000) controlled by a PC system.
168 Specific feeding rates (SFR; % pen biomass per day) were adjusted weekly according to predicted
169 growth and water temperature. Waste feed was collected per pen by means of air uplifts following
170 meal delivery ensuring satiation and allowing calculation of total daily feed intake.

171

172 **2.3 Growth and feed efficiency**

173 Fish were sampled at start and end of the freshwater phase, and then at approximately 250 g, 500 g,
174 1 kg, and ~2.5 kg in seawater prior to dietary pellet size/formulation changes. In freshwater at each
175 sampling point, 50 fish / tank were anaesthetised (50ppm Tricaine, PHARMAQ, UK), individual
176 weights (± 0.1 g) and fork lengths (± 1.0 mm) measured, while in seawater, all fish per pen were
177 counted and individually measured. Following measurement, all fish were allowed to recover in
178 aerated water before returning to their original experimental tanks/pens. Fulton's condition factor (K)
179 was calculated using: $K = (WL^{-3})100$; where W is body weight (g) and L is fork length (mm). Weight
180 data were used to calculate specific growth rate (SGRwt), thermal growth coefficient (TGC) and feed
181 conversion rate (FCR) for each sampling period where SGRwt was calculated as: $(e^{g-1}) \times 100$, where
182 $g = (\ln(W_f) - \ln(W_i)) \times (t_2 - t_1)^{-1}$ and TGC was calculated as: $(W_{f1/3} - W_{i1/3}) \times (\sum D_o)^{-1}$, where W_f is the
183 final body weight, W_i is the initial body weight and D_o is the cumulative sum of water temperature in
184 degrees per day. FCR was calculated as: $F_i / (B_f - B_i + B_m)$ where F_i is the fed intake (kg), B_f is the final
185 biomass (kg), B_i is the initial biomass (kg) and B_m is the mortality biomass for the period (kg).
186 Maturation at harvest was determined by assessment of external appearance of secondary sexual
187 characteristics and gonad development ($n = 30$ / pen), where gonadosomatic-index (GSI) was
188 calculated as $GSI (\%) = (G_w \times 100) / B_w$. Fish were classified as sexually recruited based on a threshold
189 value of $GSI > 0.20\%$ or $> 1.0\%$ for males and females respectively (Kadri et al., 1997).

190

191 **2.4 Biochemical composition and nutrient retention**

192 At the termination of the trial, 10 fish per pen (30 / diet) were euthanised using lethal anaesthesia
193 ($>200 \text{ mg L}^{-1}$ Tricaine, PHARMAQ UK) to provide samples for biochemical analyses. Five whole
194 fish per pen (15 / diet, in 3 pen pools, $n = 3$) were taken for the determination of proximate
195 composition (MOPA), minerals, and fatty acid composition. In addition, muscle/flesh (Norwegian
196 Quality Cut, NQC) were collected from 5 additional fish per pen (15 fish/diet in 3 pen pools, $n = 3$)
197 for MOPA, minerals, carotenoid (flesh), fatty acid composition, and total amino acid.

198

199 ***2.4.1. Proximate composition***

200 Whole fish and diets were ground before determination of proximate composition according to
201 standard procedures (AOAC, 2000). Five fish were pooled per pen and three technical replicates for
202 the single batch feeds were analysed. Moisture contents were obtained after drying in an oven at
203 110°C for 24 h and ash content determined after incineration at 600°C for 16 h. Crude protein content
204 was measured by determining nitrogen content ($\text{N} \times 6.25$) using automated Kjeldahl analysis (Tecator
205 Kjeltex Auto 1030 analyzer, Foss, Warrington, UK) and crude lipid content determined
206 gravimetrically after Soxhlet lipid extraction (Tecator Soxtec system 2050 Auto Extraction apparatus).

207

208 ***2.4.2. Amino acid, mineral and vitamin analyses***

209 Free amino acids and nitrogen metabolites were analysed using the Biochrome with post-column
210 derivatisation with ninhydrin as described by Espe et al. (2014). The total amino acids in whole body
211 and diets were analysed after being hydrolysed in 6N HCl for 22 h as previously described (Espe et
212 al., 2014) and used for calculation of amino acid retention (% deposited of consumed amino acid).
213 The B-vitamins biotin, niacin, folate, pantothenic acid and cobalamin were all determined by
214 microbiological methods as described in detail previously (Mæland et al., 2000; Feldsine et al., 2002).
215 Some of the B-vitamins were determined by HPLC by standard methods; thiamine (CEN, 2003),
216 vitamin B6 (CEN, 2006) and riboflavin (Brønstad et al., 2002). Multi-element determination was
217 done by ICP-MS (inductively coupled plasma mass spectrometry) (Julshamn et al., 1999). HPLC was

218 used for determination of ascorbic acid (Mæland and Waagbø, 1999) and tocopherols were analysed
219 according to standard methodology (CEN, 1999). Vitamins A and D were analysed according to
220 Moren et al. (2004) and Horvli and Lie (1994), respectively.

221

222 ***2.4.4. Lipid content and fatty acid composition***

223 Samples of whole bodies and NQC from five fish per tank were prepared as pooled pen homogenates
224 (n = 3 per diet). Total lipid was extracted from approximately 1 g of sample by homogenising in
225 chloroform/methanol (2:1, v/v) using an Ultra-Turrax tissue disrupter (Fisher Scientific,
226 Loughborough, UK), and content determined gravimetrically (Folch et al., 1957). Fatty acid methyl
227 esters (FAME) were prepared from total lipid by acid-catalysed transesterification at 50 °C for 16 h
228 (Christie, 2003), and FAME extracted and purified as described previously (Tocher and Harvie, 1988).
229 FAME were separated and quantified by gas-liquid chromatography using a Fisons GC-8160
230 (Thermo Scientific, Milan, Italy) equipped with a 30 m × 0.32 mm i.d. × 0.25 µm ZB-wax column
231 (Phenomenex, Cheshire, UK), on-column injector and a flame ionisation detector. Data were
232 collected and processed using Chromcard for Windows (version 2.01; Thermoquest Italia S.p. A.,
233 Milan, Italy). Individual FAME were identified by comparison to known standards and published
234 data (Tocher and Harvie, 1988).

235

236 ***2.4.5. Pigment analysis***

237 Flesh carotenoid levels were determined using a modified version of the JX Nippon Oil Corporation's
238 Analytical methods for Panaferd-AX. Briefly, three grams of homogenised flesh samples were
239 transferred to a 50 ml Quickfit tube. Measurements of the samples were done to two decimal places.
240 One mL of deionised water was added to the tubes. Subsequently, 5 ml of tetrahydrofuran
241 (THF)/methanol (20:1, v/v) were added and the mixture shaken vigorously for 1 min and then mixed
242 using a shaker for 5 min. Ten millilitres of iso-hexane were then added and the contents mixed
243 thoroughly before being centrifuged at 340 g for 10 min. The supernatant was transferred to a 50 ml

244 volumetric flask. Five mL of tetrahydrofuran (THF)/methanol (20:1, v/v) were added to the initial
245 tube and the process repeated three times, making a pool of four extractions in the volumetric flask.
246 Then, iso-hexane/THF/methanol (40:20:1, v/v/v) was added to the flask making up the volume of the
247 pooled extractions to 50 ml. Subsequently, 10 ml of the samples were removed and evaporated to
248 dryness under a nitrogen stream at room temperature, before being re-suspended in 2 ml isohexane
249 prior to HPLC analysis. Samples were injected on a Thermo Scientific Ultimate 300 HPLC system
250 equipped with a 150 x 4.6 mm 3 μ Phenomenex silica column with detection at a wavelength of 474
251 nm and a column temperature of 25 °C. The carotenoids were quantified using an external standard
252 of astaxanthin obtained from DSM (Heerlen, Netherlands).

253

254 **2.5 Skeletal development and spinal deformity**

255 Skeletal development and spinal malformation in smolt and harvest fish was assessed by x-ray
256 radiography. Twenty-five fish per tank/pen (n = 3) were euthanised using lethal anaesthesia (>200
257 mg L⁻¹ Tricaine, PHARMAQ UK), placed on flat trays and frozen at -20 °C prior to x-ray radiography.
258 Frozen fish were x-radiographed using a portable x-ray unit (Celtic SMR PX40 HF) with an
259 extremities plate measuring 24 x 30 cm. Each radiograph was exposed for 32 mAs and 40kV and the
260 image digitised (AGFA CR-35X). Radiographs were analysed using Adobe Photoshop CS 6 (version
261 13.0.1, Adobe system Incorporated, California, USA) with the spine divided into four regions, R1, 2,
262 3, and 4 as per Kacem et al. (1998) and deformity type categorised according to Witten et al. (2009).
263 Example pathologies were provided previously (Vera et al., 2019). Observations were made in the
264 anterior-caudal direction and the total number of vertebra was recorded for each fish.

265

266 **2.6 Gene expression**

267 At the termination of the trial, a further 5 fish per pen were euthanised using lethal anaesthesia (>200
268 mg L⁻¹ Tricaine, PHARMAQ UK) to provide samples for molecular analyses. Liver and muscle
269 samples were aseptically dissected and placed in microcentrifuge tubes containing RNA-later for the

270 assessment of gene expression using microarray (liver transcriptome) and targeted gene expression
271 through the application of real-time qPCR analysis of liver and muscle. The tissue samples were
272 homogenised in 1 mL of TRIzol® (Invitrogen, UK) and total RNA extracted in accordance with the
273 manufacturer's instructions. RNA pellets were rehydrated in MilliQ water, and total RNA
274 concentration determined using an ND-1000 Nanodrop spectrophotometer (Labtech Int., East Sussex,
275 UK), and RNA integrity assessed by electrophoresis.

276

277 ***2.6.1. Transcriptomic (microarray) analysis***

278 Transcriptome analysis of liver (male only) was performed using an Atlantic salmon custom-made
279 oligoarray with 44 k features per array in a four-array-per-slide format (Agilent Technologies UK
280 Ltd., Wokingham, UK). A dual-label experimental design was used for the microarray hybridisations
281 with Cy3-labelled test samples competitively hybridised to a common Cy5-labelled pooled-reference
282 per array. A total of 15 arrays were employed, one array per individual fish. The common reference
283 was a pool of equal amounts of amplified RNA from all test samples. Indirect labelling was employed
284 in preparing the microarray targets. Amplified antisense RNA (aRNA) was produced from each RNA
285 sample using TargetAmp™ 1-Round Aminoallyl-aRNA Amplification Kit 101 (Epicentre, Madison,
286 Wisconsin, USA), as per manufacturer's methodology, followed by Cy3 or Cy5 fluor (PA23001 or
287 PA25001, GE HealthCare) incorporation through a dye-coupling reaction, as described by Betancor
288 et al. (2016). The hybridisations were performed using SureHyb hybridisation chambers (Agilent) in
289 a DNA Microarray Hybridisation Oven (Agilent). Sample order was semi-randomised, with one
290 replicate per experimental group being loaded into each slide. For each hybridisation, 825 ng of Cy3-
291 labelled experimental biological replicate and Cy5-labelled reference pool were combined, following
292 the protocol described by Morais et al. (2012).

293

294 ***2.6.2. Quantitative Real Time PCR***

295 cDNA was reverse transcribed from 1 µg of total liver or muscle RNA using QuantiTect Reverse
296 Transcription kit (Qiagen Ltd., Manchester, UK). The resulting cDNA was diluted 20-fold with
297 milliQ water. Real-time PCR was performed using Luminaris color Hlgreen qPCR Master mix
298 (Thermo Fisher Scientific, MA, USA) and Mastercycler RealPlex 2 thermocycler (Eppendorf, UK)
299 which was programmed to perform the following protocol: 50 °C for 2 min, 95 °C for 1 min, followed
300 by 40 cycles at 95 °C for 15 s, annealing temperature (see Additional File 1) for 15 s and 72 °C for
301 30s. This was followed by a temperature ramp from 70 to 90 °C for melt-curve analysis to verify that
302 no primer-dimer artefacts were present and only one product was generated from each qPCR assay.
303 qPCR was performed in 96-well plates in duplicates per sample. The final volume of the PCR reaction
304 was 10 µL: 2.5 µL of cDNA, 5 µL of the qPCR Master Mix and 2.5 µL of forward and reverse primers.
305 The efficiency of the primers was verified and validated by doing standard curves for all genes
306 investigated. The new primers used in this study were designed using the software PRIMER3
307 (Untergasser et al., 2012). Target specificity was checked in silico using Blast (NCBI). Only primer
308 pairs with no unintended targets were selected (Additional File 1). The relative expression of target
309 genes in liver and muscle of salmon post-smolts was calculated by the $\Delta\Delta C_t$ method (Pfaffl, 2001),
310 using *β-actin*, *ef1a*, *polr2f* and *rpl1* as the reference genes, which were chosen as the most stable
311 according to RefFinder (Xie et al., 2012).

312 In liver, the genes investigated were involved in sterol metabolism (*apolipoprotein B*, *acetyl-*
313 *CoA carboxylase* and *cholesterol 7 alpha-hydroxylase*), long-chain polyunsaturated fatty acid (LC-
314 PUFA) metabolism (*fatty acyl elongase 2*, *fatty acyl elongase 5 isoform b*, *fatty acyl elongase 6* and
315 *delta-6 fatty acyl desaturase isoform a*), immune system (*interleukin 8*, *interleukin 1 beta*, *major*
316 *histocompatibility complex I* and *fas cell surface death receptor*), 1-carbon metabolism
317 (*cystathionine-β-synthase*, *betaine-homocysteine S-methyltransferase*, *methionine*
318 *adenosyltransferase1*) and *growth hormone receptor*. In muscle, the expression levels of genes
319 involved in protein and energy metabolism (*glucose transporter type 4*, *muscle-specific RING finger*
320 *protein 1a*, *muscle-specific RING finger protein 1b*, *autophagy-related protein 4 homolog B*,

321 *autophagy related 12, microtubule-associated proteins 1A/1B light chain 3B precursor* and *GABA(A)*
322 *receptor-associated protein*), immune function (*interleukin 1 beta*) and growth (*growth hormone*
323 *receptor* and *insulin-like growth factor*) were studied.

324

325 **2.7 Statistical analysis**

326 Growth and biochemical data were presented as means \pm SD and analysed using one-way analysis of
327 variance (ANOVA), general linear model, and pairwise comparison (Tukey) of means. Data were
328 checked for homogeneity of the variances by the Bartlett test and, where necessary, the data were
329 transformed (arcsine or ln) for normalisation before further statistical analysis. Statistical tests were
330 performed using Minitab statistical software (Version 17©, University of Stirling, 2016). Data on
331 vitamins, minerals, free and hydrolysed amino acids were analysed using Statistica (ver13.4, Sibco
332 software inc.) by one way ANOVA and Tukey's post-hoc test when variances were homogenous,
333 otherwise with the Kruskal-Wallis ANOVA by Ranks. Differences were considered statistically
334 significant at p values < 0.05 . Statistical analysis of microarray data was performed in GeneSpring
335 GX (Agilent Technologies, Wokingham, Berkshire, UK) using a Moderated t-test, at 0.05
336 significance. Data were submitted to the Kyoto Encyclopedia of Genes and Genomes (KEGG;
337 Kanehisa and Goto, 2000) for biological function analysis. The significance of differences in RT-
338 qPCR data between dietary groups was determined using a one-way ANOVA followed by Tukey's
339 test, using SPSS v.19 software (IBM, Armonk, NY), with significance level fixed at $p < 0.05$.

340

341 **3. Results**

342 **3.1 Mortality, growth and maturation**

343 There were no significant differences in cumulative mortality between dietary treatments in
344 freshwater or seawater (Table 5). Diet had a significant effect on weight, growth rate (TGC_{wt}) and
345 bFCR in both freshwater and saltwater (Fig. 1, Table 5). Fish fed diet L2 achieved the highest final
346 smolt weight, with fish fed diet L1 significantly lower, and diet L3 intermediary to both. This pattern

347 was mirrored by similar significant dietary effects on TGC_{wt} and bFCR. Final smolt K factor
348 decreased with increasing NP inclusion level. Diet did not significantly affect VSI, while HSI was
349 significantly lower in salmon fed diets L2 and L3 than fish fed diet L1 at smolt.

350 In seawater, fish fed diets L2 and L3 achieved a significantly higher final weight than those
351 fed diet L1, but did not differ significantly between each other. Moreover, a trend was clearly evident
352 in TGC_{wt} , which increased with increasing NP inclusion level (Table 5). In particular, salmon fed diet
353 L3 were transferred to sea at a significantly lower smolt weight than those fed diet L2, but achieved
354 a comparable body weight to fish fed diet L2 within 4 months of seawater transfer (Fig. 1). Final K
355 factor also showed a significant increase with increasing dietary NP inclusion level, and bFCR was
356 significantly lower in fish fed diets L2 and L3 than diet L1. Both VSI and HSI were not significantly
357 different at harvest between diets, and no statistical differences in maturation rates (predominantly
358 males), were observed between fish fed the different diets.

359

360 **3.2 Radiological deformity**

361 Externally visible malformation was <1 % at the end of both freshwater and seawater phases, but
362 radiologically detected spinal deformity was significantly affected by diet (Fig. 2). There was a
363 progressive reduction in detectable malformation in fish with increasing NP inclusion level. At smolt,
364 fish fed diet L3 had significantly lower detectable deformity than fish fed diet L1, and diet L2 was
365 intermediary to both. The predominant region affected with spinal malformation was in the caudal
366 trunk region (R2), with v24-28 the most commonly affected vertebral bodies. At the end of the
367 seawater phase, there was an increase, albeit non-significant, in detectable deformity in fish fed diets
368 L1 and L2 to that observed at smolt, and the pattern was maintained whereby salmon fed diet L3 had
369 significantly lower deformity than those fed diet L1, and diet L2 was intermediary to both. The
370 predominant region affected with spinal malformation was in the tail region (R3), posterior to the
371 dorsal fin, with v38 - 44 the most commonly affected vertebral bodies. Most deformities presented
372 themselves as compressive type pathologies (~53 % of total pathology) followed by reduction in

373 intervertebral spacing (~30 %) and, finally, more severe, fusion-type lesions (~15 %), and was
374 generally similar across all diets.

375

376 **3.3 Whole body composition and nutrient retention**

377 Post-smolts had similar whole body composition independent of dietary NP inclusion level, with the
378 exception of whole body ash that was significantly higher in salmon fed diet L3 than those fed diets
379 L1 or L2. Whole body Zn and Se concentration increased significantly with increasing NP inclusion
380 (Table 6). Retention of Cu decreased with increased NP inclusion, while Se retention was
381 significantly higher in L3 than both L1 and L2. Remaining nutrient retentions did not differ between
382 fish fed the different diets. NQC composition was generally comparable between diets, however there
383 was a significantly higher total lipid in fish fed diet L3 than those fed diets L1 or L2. As a percentage
384 of total fatty acids, n-6 long-chain polyunsaturated fatty acids (LC-PUFA) and docosahexaenoic acid
385 (DHA; 22:6n-3) decreased with increasing NP inclusion level

386

387 **3.4 Total amino acid concentration in whole body and free amino acids and N-metabolites in** 388 **white muscle tissue**

389 Whole body histidine and lysine retention decreased with increasing NP inclusion level, while all
390 other amino acid retentions were similar between fish fed the different diets (Table 7). Muscle content
391 of free OH-Pro, serine and glycine significantly decreased with increasing NP inclusion, while free
392 histidine, increased significantly with NP inclusion level (Table 8).

393

394 **3.5 Tissue water soluble vitamin status and retention**

395 Vitamin C showed higher concentrations in whole body of fish fed L2 and L3 than in those fed L1.
396 Vitamin C concentration in liver was not significantly different between fish fed the three diets, but
397 the trend was that vitamin C increased from L1 to L2 and then decreased again in fish fed L3. Whole
398 body pantothenic acid and niacin increased significantly with increasing NP inclusion, while

399 concentrations of the other B-vitamins in whole body were comparably independent of dietary NP
400 inclusion level (Table 9). Riboflavin retention and the retention of folate decreased significantly with
401 increasing NP inclusion, while thiamine, niacin and biotin showed no significant effect of NP
402 inclusion. Tissue specific concentrations (liver, gill and muscle) of the measured B-vitamins were
403 comparable and independent of dietary NP inclusion level, although there was a trend for increasing
404 gill pantothenic acid and muscle vitamin B6 concentration with increasing NP inclusion.

405

406 **3.6 Tissue lipid soluble vitamin status and retention**

407 With the exception of vitamin K₁ and γ -TOH (tocopherol), all other lipid soluble vitamins in whole
408 body and tissue (liver and muscle) showed a significant increase in concentration with increasing NP
409 inclusion level (Table 10). Whole body α -TOH retention increased with NP inclusion level.

410

411 **3.7 Gene expression**

412 ***3.7.1 Liver transcriptome***

413 Transcriptome analysis was performed on liver tissue in individual male fish (n = 5). Statistical
414 analysis of the microarray data returned a list of 306 and 360 annotated differentially expressed genes
415 (DEG) in liver of salmon fed the L2 and L3 diets, respectively, when compared to fish fed diet L1
416 (moderated t-test, $p < 0.05$ and fold-change > 1.3). In addition, a total of 396 genes were differentially
417 expressed (fold-change > 1.3) in the liver of salmon fed diet L3 when compared to fish fed diet L2.
418 Functional analysis for all contrasts (Kyoto Encyclopedia of Genes and Genomes, [KEGG]
419 Orthology) showed a similar distribution by categories of DEG, with the major categories being
420 metabolism, signal transduction and immune system. Within metabolism, the most represented
421 category was lipid metabolism in all cases (Fig. 3). Pathway analysis showed that the top significant
422 differentially expressed pathways in the L2 vs. L1 comparison were: PI3K-Akt signalling pathway,
423 protein processing in endoplasmic reticulum, steroid biosynthesis, sphingolipid signalling pathway,
424 phagosome and focal adhesion (Additional File 2). Moreover, the most represented metabolic

pathways suggested a downregulation of lipid metabolism (steroid biosynthesis, sphingolipid metabolism, glycerophospholipid metabolism, glycerolipid metabolism and arachidonic acid metabolism) in fish fed diet L2 compared to the L1 diet (Fig. 4A, Additional File 3). Regarding the L3/L1 contrast, the top significant differentially expressed pathways were PI3K-Akt signalling pathway, regulation of actin cytoskeleton, MAPK signalling pathway, Rap1 signalling pathway, NF-kappa B signalling pathway, focal adhesion and chemokine signalling pathway (Additional File 4). In this case, most genes involved in lipid and amino acid metabolism were downregulated whereas oxidative phosphorylation and glycolysis were upregulated (Fig. 4B, Additional File 3).

In order to identify common functional categories affected by micronutrient supplementation results from each pairwise contrast were intersected, obtaining a list of 138 common features corresponding to a total of 58 annotated genes. The most affected biological categories for L3/L1 were metabolism (27 %), signal transduction (15 %), immune system (9 %) and endocrine system (8 %). Within metabolism the most represented category was lipid metabolism at 11 % (Additional File 5).

3.7.2 Gene expression by qPCR

In liver, the expression of genes involved in sterol metabolism (*apoB*, *cyp7a1*) did not differ significantly between diets, although there was a trend for increased *apoB* expression with NP inclusion level (Table 11). Regarding genes involved in fatty acid and LC-PUFA metabolism, *fads2d6a* showed a significant decrease in fish fed diet L2 relative to fish fed diet L1, and diet L3 intermediary to both. Expression of *elovl2* was significantly higher in fish fed diet L2 than those fed diet L3, with diet L1 intermediary to both. Other genes (*acc*, *elovl5*, *elovl6*, *fas*) showed no significant effect of diet, although there was a trend for increased *acc* expression with increasing NP inclusion. Genes involved in 1-carbon metabolism (*cbs*, *bhmt*, *mat1*) and immune functions (*il8*, *il1b*) did not differ between fish fed the different diets. Finally, *ghr* expression was significantly higher in salmon fed diet L1 than those diet L2, with diet L3 intermediary to both.

451 In muscle, the expression of some genes involved in autophagy and protein turnover (*atg12*,
452 *gabapap*, *murfla*, *mlp3b*) showed significantly increased expression in salmon fed diet L3 relative to
453 those fed diet L1, and diet L2 intermediary to both (Table 11). In addition, *atg4b*, and *murflb* showed
454 the same trend although differences between groups were not statistically significant. Differences
455 between dietary groups for genes involved in growth and glucose transport (*ghr*, *igf*, *glut4*) were not
456 significant. However, the pattern of *ghr* expression in muscle mirrored that of *ghr* in the liver, with
457 fish fed diet L2 having lowest expression and those fed diet L1 highest, while *igf* expression showed
458 a trend of increased expression with increasing NP inclusion. Finally, genes associated with immune
459 function (*illbm*) were not statistically different in muscle of post-smolt salmon.

460

461 **4. Discussion**

462 **4.1 Growth**

463 In the current long-term study utilising S0+ Atlantic salmon parr, growth in freshwater and the pattern
464 of final smolt body weight between diets mirrored that previously reported in S1+ Atlantic salmon
465 parr fed the same diet formulation (Taylor et al., 2019). In both studies, fish fed diet L2 (2xNP)
466 achieved a significantly higher final smolt weight, than fish fed diets L1 and L3, which was reflected
467 in an improved growth rate (TGC_{bw}) and more efficient FCR. In both studies, irrespective of smolt
468 regime, diet L3 (4xNP) initially showed an increased growth rate relative to diet L1 (1xNP), but
469 subsequently decreased, resulting in a final smolt weight intermediary to diets L1 and L2. This growth
470 reduction occurred around 14 weeks post-feeding in the S1 salmon (Taylor et al., 2019) and within 4
471 weeks of feeding under the S0+ regime in the current study. These results are in contrast to Hemre et
472 al. (2016) and Prabhu et al. (2019), where parr were fed similar dietary formulations for 12 weeks but
473 a decrease in growth rate beyond 2xNP was not observed. However, parr were not taken through parr-
474 smolt transformation (PST) in that short-term study. Of note, in both S1+ (Taylor et al., 2019) and
475 the current S0+ parr study, the occurrence of growth reduction commenced at the start of the rising
476 day length in spring (ambient in S1+; and following switch from LD 14:10 to LL in S0+). Given that

there are significant morphological, physiological and metabolic transformations during PST (Hoar, 1988), then there may be differing dietary requirements and nutrient re-profiling during this transition. As such, the current study suggested certain components in NP could be in “excess” or detrimental to growth when supplemented at >2xNP levels, particularly in Atlantic salmon undergoing PST. However, both the current study and that of Taylor et al. (2019) have not been able to conclusively ascertain which components and/or interactions within the NP were detrimental, and further studies are clearly warranted to elucidate this apparent phenomenon.

In seawater, growth rate (TGC_{bw}) improved linearly with increasing NP, and bFCR was more efficient in fish fed diets L2 and L3 than L1, such that fish fed diet L3 while initially stocked at a significantly smaller body weight at smolt, attained a body weight comparable to that of fish fed diet L2 at harvest. By contrast, Hemre et al., (2016) did not find a strong effect of graded NP on growth rate in post-smolts, but this may in part be related to differences between the former and current study: shorter duration (5 months vs. 12 months), cooler water temperatures experienced (4.1-10 °C vs. 6-16 °C) and subsequently lower TGC_{bw} (2.37-2.55 vs. 2.9-3.1). Nonetheless, with the progressive reduction of FM and FO down to 5 % and 3 % during the current trial, results do confirm and support the earlier shorter-term studies (Hamre et al., 2016; Hemre et al., 2016; Taylor et al., 2019; Prabhu et al., 2019) that specific components of the NP are required to be supplemented at between 2x and 4x current recommendation when utilising high plant-based diets for the duration of the life cycle. The specific components of the NP that are associated with improvements in growth and performance will be discussed in subsequent nutrient status sections.

4.2 Skeletal Health

In contrast to growth, we observed a clear linear decrease in radiological deformity with increasing NP inclusion in freshwater which concurs with results from our previous short-term feeding study (Vera et al 2019). Further, in the current study, there was also a progression of spinal pathology in post-smolts when reassessed at harvest compared to initial smolt in fish fed diets L1 and L2, while

diet L3 showed no further increase in pathology. Concomitantly, spinal deformity was also inversely correlated with final whole body ash concentration, a widely accepted indicator of body/bone mineralisation (Baeverfjord et al., 2019), indicating increasing NP level improved post-smolt mineral status. Previously, upregulation of bone gene expression associated with extracellular matrix (ECM) mineralisation (*ALP*, *Mgp*, *Ostcn*), and corresponding decrease in skeletal pathology was observed following increased NP supplementation in parr (Vera et al., 2019). Effects were hypothesised to be mediated through altered P-Ca homeostasis pathways, as reflected by increased whole-body Ca:P ratios in smolts. Candidate components of the NP were suggested as vitamins D₃ and K as cofactors of calcium sequestering. However, in the current study, improved mineralisation was not explained by differences in whole body macromineral concentration or retention in post-smolts which were similar across all diets and in the normal range for salmon (Shearer et al., 1994). Similarly, Fjelldal et al. (2009) reported a reduction in vertebral deformity at 4.1 kg in S0+ post-smolts fed a high mineral diet for the first 8 weeks post sea entry, and while analysis of plasma Ca, P and D-vitamin metabolites reflected changes in P homeostasis, these factors could not explain the preventive effect of a high mineral diet on development of bone deformities. To this end, given that analysed dietary vitamin D₃ and K in the current formulation were within established requirement ranges of 0.15-0.19 mg kg⁻¹ (NRC, 2011) and >0.1 mg kg⁻¹ (Krossøy et al., 2009) respectively, it is unlikely these components of the NP are directly attributable to the graded decrease in spinal pathology observed in post-smolts, and other components of the NP are worth consideration.

Whole body Zn and Se increased across diets with NP inclusion rate, with both being identified as the principal limiting micro-minerals in plant-based diets (Prabhu et al., 2018; Prabhu et al., 2019). Selenium is involved in antioxidant mechanisms and redox reactions in liver and blood cells, yet the role of Se in bone metabolism remains unexplored in salmonids to date (Baeverfjord et al., 2019). Recent studies have shown that Se, in the form of selenoproteins, plays a vital role in bone metabolism in humans (Pietschmann et al., 2014), with plasma selenoprotein P concentrations found to positively correlate with bone mineral density in elderly women (Zhang et al., 2014). As such,

529 although whole body Se levels across all diets were within the normal reported range for Atlantic
530 salmon (Prabhu et al., 2016), the role of Se in teleost bone development should be investigated.

531 It is well established that Zn is the most abundant micro-mineral in the vertebrae of many
532 teleost fish species and has functional importance in the activity of ALP, and associated processes
533 related to mineralisation of bone (Prabhu et al., 2016). Thus, improved Zn status and downstream
534 processes may be directly attributable to the observed reduction in spinal pathology and progression
535 as post-smolts, particularly in fish fed diet L3 (4xNP). While studies on spinal development have
536 often focused on dietary macro- and micro-minerals, the role of specific vitamins within the NP
537 should not be overlooked as having a beneficial effect on skeletal health. Whole body vitamin A
538 levels suggested that diet L1 covered minimum requirement ($>0.75\text{mg kg}^{-1}$, Moren et al., 2004), and
539 diet L3 had significantly lower skeletal deformity. Diet L1-L3 in this study contained 3.7-12.2mg
540 kg^{-1} VA, well below the upper limits shown to induce skeletal pathology ($>37\text{ mg kg}^{-1}$) (Ørnsrud
541 et al., 2002; Ørnsrud et al., 2013). Therefore, it is unlikely that vitamin A deficiency or
542 hypervitaminosis is attributable to the differences in spinal deformity rates observed. The specific
543 role of B-vitamins on bone development in humans has been documented to a degree (Dai and Koh,
544 2015) but, as yet, has not been investigated in fish (Lall & McCrea, 2007). Evidence from humans as
545 reviewed by Dai and Koh (2015) has shown a variety of effects of vitamin B deficiencies including:
546 reduced dietary intake of B6 is associated with lower bone mineral density; synthesis of methionine
547 depends on both folate and B12 for remethylation, and deficiency of either B vitamin results in
548 megaloblastic changes in bone marrow; B vitamin deficiency affects the anatomical/biochemical
549 properties of bone if it is profound enough to cause hyperhomocysteinemia that is sufficient to induce
550 an accumulation of homocysteine in the bone tissue. In light of the growing research in B vitamin
551 involvement in bone metabolism in humans, greater consideration of B vitamin requirements in fish
552 would also warrant consideration in relation to skeletal health.

553

554 **4.3 Protein and amino acid status and retention**

555 The present study increased dietary taurine and histidine keeping the methionine constant in high
556 plant protein diets for Atlantic salmon growing from parr to approximately 2.5 kg BW. NRC
557 requirement for methionine in Atlantic salmon is reported to be 7 g kg⁻¹ (NRC 2011). However, it
558 was previously reported that dietary methionine of 7.9 vs 11.4 g kg⁻¹ methionine improved growth
559 and hepatosomatic index in salmon fed the latter diet (Espe et al., 2014). In another study, Atlantic
560 salmon fed high plant protein diets containing either 8 or 11 g methionine per kg of diet, the weight
561 gain in the latter diet did not differ from the control group fed a high fishmeal diet (760 g kg⁻¹ diet).
562 Thus, in the current trial it was decided to maintain the dietary methionine constant at around 9 g kg⁻¹
563 diet, while histidine and taurine were increased in the nutrient packages.

564 The retention of amino acids is known to be high when an amino acid is slightly deficient and
565 then decrease as the requirement for amino acid deposition is met or increased above requirement for
566 protein deposition (Espe et al., 2017). In the current study this is also true for histidine, where
567 retention decreased as dietary histidine increased. In addition to histidine, lysine retention was also
568 reduced in salmon fed the L3 diet as compared to fish fed the L1 and L2 diets. Lysine is mainly used
569 for protein deposition (Espe et al., 2007) and the reduced retention of lysine in fish fed the highest
570 nutrient package thus might indicate a lower protein gain in muscle of these fish. However, as protein
571 retention did not differ in fish fed any of the diets, reduced lysine retention does not seem to be
572 associated with a reduced muscle protein deposition in the current trial. Thus, to address the reduced
573 retention of lysine, interaction with other nutrients in the NP needs to be further explored. It is well
574 known that both amino acid supply and the balance between dietary amino acids affect growth
575 hormone, IGF and GH (Hevrøy et al., 2007; Bower et al., 2008; Valente et al., 2012). In the current
576 trial these were not affected, which supports the fact that protein retention did not differ between fish
577 fed any of the diets in the current trial.

578 Any imbalanced supply of amino acids may increase the ubiquitination and thereby
579 degradation of proteins (Bower et al., 2008; Valente et al., 2012). The gene expression level of *murf*,
580 *gabarab* and *atg12* were higher in the higher nutrient package in the current trial, but this is more

likely due to a higher degradation of soluble protein than the degradation of muscle protein. However, this also needs to be addressed in new trials using fewer variables.

As expected, free histidine in muscle increased as dietary histidine increased simply reflecting the higher dietary histidine content in line with previous results (Remø et al., 2014). Taurine in muscle on the other hand did not increase following increased dietary taurine. This is most likely due to increased excretion of excess taurine or that the taurine fed is directed to other tissues such as the liver (Espe et al., 2012a, 2012b). Both dietary histidine and taurine are more associated with oxidation and protection against ROS than with growth and metabolism (Remø et al., 2014) as well as survival of the cells (Espe and Holen, 2013), but these aspects were not addressed in the current trial. In addition, dietary taurine has been reported to reduce the visceral fat in salmon (Espe et al., 2012a, 2012b), but as neither HSI nor VSI were affected by increased inclusion of NP, the L1 diet appears to contain sufficient taurine to not significantly affect adiposity in salmon in the current trial.

Finally, free OH-proline decreased with increased nutrient package which might be due to an interaction with some of the other micronutrients fed, e.g. vitamin C. It is, however, unlikely that the lower OH-pro in fish muscle of fish fed the L3 diet is associated with any increased muscle degradation as deposition and growth actually was greatest in this group, although protein retention did not differ between the groups.

4.4 Water-soluble vitamins

Since growth was improved when increasing the nutrient package, without increasing methionine, the other components must have affected growth positively without the present level of methionine being critical, e.g. some of the B-vitamins will be critical for salmon to obtain maximum growth (Hemre et al., 2016), as well as some of the minerals (Prabhu et al., 2019a, b). In an experiment with salmon parr, it was estimated that the 2xNP would cover requirement for several micronutrients, especially when given with high plant raw material diets (Hemre et al., 2016, Hamre et al., 2016, Prabhu et al., 2019a). The present study, therefore, confirmed earlier conclusions that plant-based

607 diets need new requirement estimates for several micronutrients. The suggested body-concentrations,
608 health outcomes and growth response used as requirement markers were not sufficient in the former
609 study (Hemre et al., 2016), and the present study confirmed this.

610 Pantothenic acid in whole body was higher in fish fed diets L2 and L3 compared to diet L1,
611 while niacin increased gradually from L1 to L3, significantly at each step. Pantothenic acid in gill
612 tissue also increased but differences were not significant. There was also a tendency for higher B6
613 concentration in whole body in fish fed diets L2 and L3 compared to L1. This is in agreement with
614 earlier findings for salmon post-smolt (Hemre et al., 2016). These results suggest that these three B-
615 vitamin optimal levels should be raised when compared to NRC (2011), to the 2xNP level for
616 pantothenic acid and B6 and to the 4xNP level for niacin.

617 Vitamin B6 interacts highly with protein metabolism (Hansen et al., 2015). Pyridoxine
618 functions as phosphorylated coenzymes pyridoxal phosphate (PLP) in many reactions, such as
619 transaminases in amino acid metabolism. Pyridoxine is a cofactor for the enzyme aromatic amino
620 acid decarboxylase, which is required for the synthesis of neurotransmitters (Rorsman et al., 1995).
621 Therefore, symptoms and consequences of vitamin B6 deficiency are many, diverse and severe, and
622 include nervous disorders and abnormal behavior, none of which was registered in the present study.
623 Tissues with mitochondria-rich cells and cells that undergo rapid cell divisions like gill tissue are
624 especially sensitive to pantothenic acid deficiency. The 1xNP levels seemed to be sufficient for the
625 other B vitamins (thiamine, riboflavin, folic acid and biotin), as found by Hemre et al. (2016). There
626 was, however, a tendency for a decrease in retention of B-vitamins, significant only for riboflavin
627 and folate. This may point to either an interaction and/or control factor at the gut-barrier level, or
628 excretion of excess vitamin, since whole body levels were maintained or increased.

629 Vitamin C in whole body increased from diet L1 to L2 and was then unchanged in fish fed
630 diet L3. This is in accordance with previous results (Hamre et al., 2016). In the liver, changes in
631 vitamin C concentrations were not significant ($p = 0.06$), but there was a tendency for an increase in
632 fish fed diet L2 compared to L1, and then a decrease in fish fed L3. The variation was very high in

633 both the L2 and L3 groups. It is well known that vitamin C may act as a pro-oxidant at high
634 concentrations (Hamre et al., 1997), and this may indicate that diet L3, and perhaps L2, contained too
635 much vitamin C, and an optimal vitamin C level at the 2xNP level or slightly less is needed.

636

637 **4.5 Lipid soluble vitamins**

638 Lipid soluble vitamins have a tendency to accumulate in the body in response to increased
639 supplementation. Therefore, it is not possible to use body concentrations as an indication of
640 requirement. This is true for α -tocopherol (TOH) which normally shows a linear relationship between
641 supplementation and whole body concentration (Hamre et al., 1997; Hamre et al., 2016) as confirmed
642 in the present study. Hamre et al. (2016) recommended supplementation with at least 150 mg kg⁻¹ α -
643 TOH to compensate for variations in composition of feed ingredients and variation in oxidative stress,
644 in response to handling, seasons and disease. TOH was not supplemented, but came from the feed
645 ingredients, and the feed and whole body concentrations did not vary among the dietary treatments.

646 For vitamin A, the situation is more complicated due to the presence of many different forms.
647 Most of the vitamin A is present in the storage form retinol-esters (Moren et al., 2004), and the method
648 used here determined esters and free forms combined. However, both free retinol and retinol esters
649 increased exponentially with increasing concentration of vitamin A in the diet, and there was a large
650 increase in body stores with an increase of dietary retinol equivalents from 0.75 to 2.5 mg kg⁻¹ diet.
651 Nevertheless, a requirement was found in Atlantic halibut of 2.5 mg kg⁻¹ diet based on cell
652 proliferation in the intestine as measured by DNA content, PCNA and maturity of the brush border
653 membrane by enzyme activities (Moren et al., 2004). With reference to Moren et al. (2004) and the
654 whole body levels found here, diet L1 had enough vitamin A (3.7mg kg⁻¹) to cover the requirement
655 (≥ 0.75 mg kg⁻¹ NRC, 2011).

656 The diets were supplemented with the synthetic pro-vitamin K, menadione (vitamin K3).
657 Menadione is very unstable during feed processing (Krossøy et al., 2009), and the analysed dietary
658 levels of K3 were only 1/10 of the supplemented levels. Menadione is metabolised to menakinones

(MK4-13, vitamin K2) in the body (Krossøy et al., 2011) and therefore, MK4 increased in response to increase in dietary vitamin K3, while whole body vitamin K1 was similar in fish fed the different treatments. Krossøy et al. (2009) fed Atlantic salmon eight diets supplemented with 0 to 50 mg kg⁻¹ vitamin K3 and found no effects on fish performance, or blood coagulation time and vertebrae stiffness. They concluded that the feed ingredients probably contained enough vitamin K (0.1 mg kg⁻¹) to cover the requirement, which is consistent with the findings in the present study. Estimated requirements in other species range from 0.2 to 2 mg kg⁻¹ (NRC 2011).

Vitamin D was supplemented at 0.05, 0.10 and 0.20 mg kg⁻¹, but the analysed levels in muscle were 0.15-0.19 mg kg⁻¹, reflecting that the ingredients contained significant amounts of vitamin D₃. Marine feed ingredients usually contain ample amounts of vitamin D and this may be the reason that vitamin D requirements in Atlantic salmon have been difficult to define. Furthermore, high doses of vitamin D up to 57 mg kg⁻¹ seem to be well tolerated by Atlantic salmon (Horvli et al., 1998; Graff et al., 2002). In other fish species estimated requirements vary from 0.006 to 0.060 mg kg⁻¹ (NRC 2011), well below the supplementation levels in the present study.

673

674 **4.6 Minerals**

Optimal supply of dietary micro-minerals is essential to maintain body mineral homeostasis (Watanabe et al., 1997). Deficient or sub-optimal supply of dietary micro-minerals will deplete body mineral status, especially over the long term (Maage & Julshman, 1993). The NPs had graded levels of micro-minerals of which Zn and Se increased with increasing NP inclusions from diets L1 to L3. In plant-based salmonid feeds, Zn and Se have been identified as the most limiting micro-minerals (Prabhu et al., 2018, 2019a). In Atlantic salmon parr and post-smolt, dietary Zn levels of 101-132 mg kg⁻¹ and 140-177 mg kg⁻¹ respectively supplied by NP inclusion of 100-150 and 150-200 were able to meet the Zn requirement over the short-term (Prabhu et al., 2019a). Herein, on a long-term basis with an almost 30-fold increase in weight during the trial period, the L1 diet with 94 mg Zn kg⁻¹ decreased the Zn status to half and one-third in L2 and L3 fed-fish, respectively. Similar reductions

in whole body Zn ranging from $\frac{1}{3}$ to $\frac{1}{2}$ were observed in Atlantic salmon parr fed sub-optimal Zn levels (Maage & Julshman, 1993). The L2 diet with 156 mg Zn kg⁻¹ was able to sustain the body Zn status on parr to the initial Zn status (18 mg kg⁻¹ wet weight). However, the initial Zn status of the Atlantic salmon in this study was lower than values reported in the literature for similar size fish (Maage & Julshman, 1993; Shearer et al., 1994). The Zn status of Atlantic salmon fed the L2 diet (18 mg kg⁻¹ wet weight, averaging 2.3 kg) was considerably lower than the Zn status of 27-29 mg kg⁻¹ of 500 g post-smolt fed similar Zn levels in a previous trial (Prabhu et al. 2019a). Se levels in all three dietary groups were in the normal range observed in salmonids or Atlantic salmon specifically (Prabhu et al., 2016), mostly because the high background Se levels in basal diet already met the requirement. Although the NP composition did not change between the present and previous studies (Hamre et al., 2016; Prabhu et al., 2019a), the total dietary Se levels were considerably different for corresponding NP inclusions, thereby Se recommendations differ between studies. Variation in Se content of feed ingredients have to be considered in feed formulations and regulative legislation. The supplementation of manganese of 12 mg/kg diet in diet L1 corresponding to a total dietary Mn of 42 mg/kg diet was sufficient to maintain normal body status. Recently, Prabhu et al. (2019b) also showed that Mn supplementation of 14 mg/kg as MnSO₄ satisfied the minimal dietary requirement of Atlantic salmon post-smolt fed plant-based diets. The macro-mineral composition was within normal range and unaltered in Atlantic salmon fed the different diets implying the diets met the macro-mineral needs of Atlantic salmon over the entire study period. Overall, Atlantic salmon post-smolt in seawater were able to maintain their body status in the long run with minimal dietary supply of most micro-minerals from the L1 diet, except for Zn which required L2 levels.

706

707 **4.7 Gene Expression**

708 In general, the number of significantly differentially regulated functional pathways were greater for
709 the L3 vs. L1 dietary comparison, than for the L2 vs. L1 comparison. In fact, not only the number of
710 up- or down-regulated pathways was higher in L3/L1, but also the number of DEGs involved in each

711 pathway was greater, suggesting a correlation between increased dietary micronutrient
712 supplementation in low marine diets and changes in liver transcriptome. Regarding the effects on
713 specific pathways, in both diet comparisons (L3/L1 and L2/L1), the terpenoid backbone and steroid
714 biosynthesis pathways were down-regulated. A micronutrient dependent epigenetic gene regulation
715 of the upstream regulator (acetyl-CoA carboxylase alpha, *acaca*) that provides malonyl-CoA
716 substrate for the biosynthesis of fatty acids has been demonstrated (Saito et al, *publication pending*).
717 The terpenoid backbone pathway underpins cholesterol (sterol), heme and vitamin (retinoids, retinol,
718 retinoic acid, menadione) synthesis while both biosynthetic pathways are involved in steroid (e.g.
719 testosterone) hormone production. The steroid and sterol biosynthetic pathways were also the most
720 significantly affected in zebrafish offspring after feeding the parental generation a diet deficient in
721 micronutrients (vitamin B12, vitamin B6, folic acid, methionine and choline) (Skjaerven et al., 2018).
722 Further, the offspring from deficient parents had accumulated higher levels of lipid inclusion in the
723 hepatocytes and showed a downregulation in the steroid biosynthetic process and widespread
724 epigenetic changes of gene regulation in the liver (Skjaerven et al., 2018). This suggests that
725 micronutrient supplementation may not only be affecting nutrient retention directly, but also via
726 epigenetic mechanisms altering gene regulation and downstream biochemical process.

727 These effects on gene expression may be in response to the increased supplementation of the
728 L2 and L3 diets with cholesterol, menadione (vitamin K) and retinol (vitamin A) as part of the
729 micronutrient premix and therefore reflect an increased requirement of these compounds in fish fed
730 diet L1 (Kortner et al., 2014). The lower level of dietary pantothenic acid could also have affected
731 the up-regulation of several metabolic pathways in fish fed diet L1. Pantothenic acid is required for
732 synthesis of coenzyme A, which participates in a range of acyl transfer reactions in energy production,
733 fatty acid oxidation, cholesterol and steroid synthesis, heme synthesis, amino acid catabolism,
734 acetylcholine synthesis and many other acetylation and acylation reactions (Olsvik et al., 2013).

735 Increased expression of genes related to glyoxylate/dicarboxylate metabolism was observed
736 in salmon fed diet L1 in comparison to fish fed L3, suggesting lower levels of glucose in liver of L1

737 salmon and therefore the need to synthesise it from fatty acids. Meanwhile, an activation of
738 intermediary metabolism in fish fed diet L3 may indicate higher levels of glucose available for energy
739 production, and an enhancement of cellular energy levels as suggested by the increased number of
740 up-regulated genes in glycolysis/gluconeogenesis, and oxidative phosphorylation pathways in fish
741 fed diet L3. The differences in hepatic metabolic activities, energy generation and transport and
742 growth performance between salmon fed diets L3 and L1 could be linked to the higher levels in L3
743 fish of B vitamins (riboflavin, niacin), which are involved in a range of cellular processes, most
744 importantly energy generation and transport that enables fish fed diet L3 to sustain vital metabolic
745 processes without compromising the amount of energy available for growth (Hansen et al., 2015).

746 The q-PCR results of genes involved in LC-PUFA biosynthesis showed down-regulation of
747 biosynthesis with increasing micronutrient supplementation. A down-regulation of *fads2d6a* was
748 found, indicating greater LC-PUFA biosynthesis capacity in the liver of L1 salmon. However, these
749 differences were not fully reflected in hepatic n-3 LC-PUFA profiles in the present study, which
750 showed no major differences between the dietary groups. In muscle, *mlp3b*, *atg12* and *gabarap* genes
751 were found to be upregulated in L3-fed salmon. These genes are involved in autophagosome mediated
752 autophagy, playing an important role in autophagosome formation and sequestration of cytosolic
753 cargo into double membrane vesicles, leading to subsequent degradation after fusion with lysosomes
754 (Mannackand Lane, 2014).

755

756 5. Conclusions

757 The present study as part EU ARRANA project, confirmed earlier studies (Hamre et al., 2016;
758 Hemre et al., 2016; Prahbu et al., 2019; Taylor et al., 2019; Vera et al., 2019) that plant-based diets
759 for Atlantic salmon need new requirement estimates for several micronutrients. In particular, these
760 optimal levels may differ between freshwater and seawater phases of the production cycle, and that
761 requirements for growth and skeletal health may also differ. Based on the present results and
762 conclusions from earlier studies (Hamre et al., 2016; Hemre et al., 2016; Prahbu et al., 2019a; Taylor

et al., 2019; Vera et al., 2019), NRC (2011) should be revised for diets in which plants form the major part of the ingredient mix. For these plant-based diets most B-vitamin optimal levels were met at the NRC (2011) recommendations, but beneficial effects were found with moderately increased levels (L2) of niacin, riboflavin and cobalamin, vitamin C, and the minerals Zn and Se. However, as indicated in all previous studies, too high levels of the micro-nutrient package (L3) may reduce these beneficial effects.

769

770 5. Acknowledgements

771 This study was funded by the European Commission FP7 Integrated Project No. 288925, Advanced
772 Research Initiatives for Nutrition & Aquaculture (ARRAINA). A special thanks goes to all technical
773 staff at the Niall Bromage Freshwater Research Facility, Stirling and the Ardnish Feed Trial Unit,
774 Lochailort, MOWI Scotland for care and maintenance of the animals.

775

776 6. References

- 777 Baeverfjord, G., Prabhu, A.J.P., Fjelldal, P.G., Albrektsen, S., Hatlen, B., Denstadli, Y., Ytteborg, E.,
778 Takle, H., Lock, E.-J., Berntssen, M.H.G, Lundebye, A.K., Åsgård, T., Waagbø, R. 2019.
779 Mineral nutrition and bone health in salmonids. Rev. Aquacult. 11, 740-765.
- 780 Betancor M.B., Olsen, R.E., Solstorm, D., Skulstad, O.F., Tocher, D.R. 2016. Assessment of a land-
781 locked Atlantic salmon (*Salmo salar* L.) population as a potential genetic resource with a focus
782 on long-chain polyunsaturated fatty acid biosynthesis. Biochim Biophys Acta 1861, 227-238.
- 783 Bower, N.I., Li, X., Taylor, R., Johnston I.A. 2008. Switching to fast growth: the insulin-like growth
784 factor (IGF) system in skeletal muscle of Atlantic salmon. J. Exp Biol. 211, 3859-3870
- 785 CEN, 1999. Foodstuffs – Determination of vitamin E by high performance liquid chromatography -
786 Measurement of alpha-, beta-, gamma- and delta-tocopherols. prEN 12822, The European
787 Committee for Standardization.
- 788 CEN. 2003. Foodstuffs -Determination of vitamin B1 by high performance liquid chromatography.
789 prEN 14122, The European Committee for Standardization.
- 790 CEN. 2006. Foodstuffs -Determination of vitamin B6 by high performance liquid chromatography.
791 prEN 14164, The European Committee for Standardization.

792 Dai, Z., Koh, W.P. 2015. B-Vitamins and Bone Health—A Review of the Current Evidence. *Nutrients*,
793 7, 3322-3346.

794 Espe, M., Lemme, A., Petri, A., El-Mowafi, A., 2006. Can Atlantic salmon (*Salmo salar*) grow on diets
795 devoid of fish meal? *Aquaculture* 255, 255-262.

796 Espe, M., Lemme A., Petri, A., El-Mowafi, A., Espe, M., Ruohonenm K., El-Mowafi, A. 2012a. Effect
797 of taurine supplementation on metabolism and body lipid to protein ratio in juvenile Atlantic
798 salmon *Salmo salar*. *Aquacult. Res.* 43, 349-360

799 Espe, M., Ruohonen, K., El-Mowafi, A. 2012b. Dietary FPC reduces the viscera somatic index in on-
800 growing Atlantic salmon fed diets containing 20% fishmeal inclusion.. *Aquacult. Nutr.* 18, 599-
801 609.

802 Espe, M., Andersen, S.M., Holen, E., Rønnestad, I., Veiseth-Kent, E., Zerrahn, J-E., Aksnes, A. 2014.
803 Methionine deficiency does not increase polyamine turnover through depletion of liver S-
804 adenosylmethionine (SAM) in juvenile Atlantic salmon. *Br. J. Nutr.* 112, 1274-1283.

805 Espe, M., Andersen, S.M., Veiset-Kent, E., Rønnestad, I., Holen, E., Zerrahn, J.-E., Aksnes, A. 2017.
806 Choline supplementation slightly increased lipid gain, but had no impact on growth and protein
807 gain, while surplus methionine improved growth due to increased protein accretion without
808 affecting lipid deposition. *Aquacult. Nutr.* 23. 1086-1094.

809 Feldsine, P., Abeyta, C., Andrews, W.H., 2002. AOAC International methods committee guidelines
810 for validation of qualitative and quantitative food microbiological official methods of analysis.
811 *J. AOAC Int.* 85, 1187–1200.

812 Fjelldal, P.G., Hansen, T., Breck, O., Sandvik, R., Waagbø, R., Berg, A. & Ørnsrud, R. 2009.
813 Supplementation of dietary minerals during the early seawater phase increase vertebral strength
814 and reduce the prevalence of vertebral deformities in fast growing under-yearling Atlantic
815 salmon (*Salmo salar* L.) smolt. *Aquacult.Nutr.* 15, 366–378.

816 Gatlin III, D.M., Barrows, F.T., Brown, P., Dabrowski, K., Gaylord, T.G., Hardy, R.W., Herman, E.,
817 Hu, G., Krogdahl, A., Nelson, R., Overturf, K., Rust, M., Sealey, W., Skonberg, D., Souza, E.J.,
818 Stone, D., Wilson, R., Wurtele, E., 2007. Expanding the utilization of sustainable plant products
819 in aquafeeds: A review. *Aquacult. Res.* 38, 551-579.

820 Hamre, K., Waagbø, R., Berge, R.K., Lie, O., 1997. Vitamins C and E interact in juvenile Atlantic
821 salmon (*Salmo salar*, L.). *Free Radical Biol. Med.* 22, 1-2.

822 Hamre, K., Sissener, N.H., Lock, E.J., Olsvik, P.A., Espe, M., Torstensen, B.E., Silva, J., Johansen,
823 J., Waagbø, R., Hemre, G.I., 2016. Antioxidant nutrition in Atlantic salmon (*Salmo salar*) parr
824 and post-smolt, fed diets with high inclusion of plant ingredients and graded levels of
825 micronutrients and selected amino acids. *PeerJ* 4:e2688.

826 Hansen, A.-C., Waagbø, R., Hemre, G.-I., 2015. New B vitamin recommendations in fish when fed
827 plant-based diets. *Aquacult. Nutr.* 21, 507-527.

828 Hardy, R.W., 2010. Utilization of plant proteins in fish diets: Effects of global demand and supplies
829 of fishmeal. *Aquacult. Res.* 41, 770-776.

830 Hemre, G.-I., Sandnes, K., 2008. Seasonal adjusted diets to Atlantic salmon (*Salmo salar*):
831 Evaluations of a novel feed based on heat-coagulated fish mince, fed throughout 1 year in sea:
832 Feed utilisation, retention of nutrients and health parameters. *Aquaculture* 274, 166-174.

833 Hemre, G.-I., Lock, E.-J., Olsvik, P.A., Hamre, K., Espe, M., Torstensen, B.E., Silva, J., Hansen, A.-
834 C., Waagbø, R., Johansen, J.S., Sanden, M., Sissener, N.H. 2016. Atlantic salmon (*Salmo salar*)
835 require increased dietary levels of B-vitamins when fed diets with high inclusion of plant based
836 ingredients. *PeerJ* 4:e2493.

837 Hevrøy E.M., El-Mowafi, A., Taylor, R., Olsvik, P.A., Norberg, B., Espe, M. 2007. Lysine intake
838 affects gene expression of anabolic hormones in Atlantic salmon, *Salmo salar*. *General and*
839 *Comparative Endocrinology*, 152, 39-46.

840 Hoar, W.S., 1988. The physiology of smolting salmonids. In: Hoar, W.S., Randall, D.J. (Eds.), *Fish*
841 *Physiology*, vol. 11B. Academic Press, London, 275–343.

842 Julshamn, K., Brenna, J., Holland, R., Tanner, S., 1999. Plasma source mass spectrometry- New
843 developments and applications. *Roy. Soc. Chem.* 241, 167-172.

844 Kadri., S., Metcalfe, N.B., Huntingford, F.A., Thorpe, J.E. 1997. Early morphological predictors of
845 maturity in one-sea-winter Atlantic salmon. *Aquacult. Int.* 5, 41-50.

846 Kanehisa M, Goto S., 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acid Res.*
847 28, 27-30.

848 Kortner, T.M., Björkhem, I., Krasnov, A., Timmerhaus, G., Krogahl, A., 2014. Dietary cholesterol
849 supplementation to a plant-based diet suppresses the complete pathway of cholesterol synthesis
850 and induces bile acid production in Atlantic salmon (*Salmo salar* L.). *Br. J. Nutr.* 111, 2089-
851 2103.

852 Krossøy, C., Waagbø, R., Fjelldal, P.G., Wargelius, A., Lock, E.J., Graff, I.E, Ørnsrud, R. 2009.
853 Dietary menadione nicotinamide bisulphite (vitamin K3) does not affect growth or bone health
854 in first feeding fry of Atlantic salmon (*Salmo salar* L). *Aquac. Nutr.*, 15, 638–649.

855 Krossøy, C., Lock, E.-J., Ørnsrud, R., 2011. Vitamin K-dependent γ -glutamylcarboxylase in Atlantic
856 salmon (*Salmo salar* L.). *Fish Physiol. Biochem.* 36, 627-635.

857 Lall, S.P., Lewis-McCrea, L.M., 2007. Role of nutrients in skeletal metabolism and pathology in fish
858 - an overview. *Aquaculture* 267, 3-19.

859 Mæland, A., Rosenlund, G., Stoss, J., Waagbø, R. 1999. Weaning of Atlantic halibut (*Hippoglossus*
860 *hippoglossus* L.) using formulated diets with various levels of ascorbic acid. Aquacult. Nutr. 5,
861 211-219.

862 Mæland, A., Rønnestad, I., Fyhn, H.J., Berg, L., Waagbø, R. 2000. Water-soluble vitamins in natural
863 plankton (copepods) during two consecutive spring blooms compared to vitamins in *Artemia*
864 *franciscana* nauplii and metanauplii. Mar. Biol. 136, 765–772.

865 Maage, A., Julshamn, K. 1993. Assessment of zinc status in juvenile Atlantic salmon (*Salmo salar*)
866 by measurement of whole body and tissue levels of zinc. Aquaculture 117, 179-191.

867 Mannack, L.V.J.C., Lane, J.D. 2014. The autophagosome: current understanding of formation and
868 maturation. Res. Rep. Biochem. 2015, 39-58.

869 National Research Council (NRC), 2011. Nutrient Requirements of Fish and Shrimp. The National
870 Academies Press, Washington DC.

871 Menoyo, D., Lopez-Bote, C.J., Diez, A., Obach, A., Bautista, J.M., 2007. Impact of n - 3 fatty acid
872 chain length and n - 3/n - 6 ratio in Atlantic salmon (*Salmo salar*) diets. Aquaculture 267, 248-
873 259.

874 Morais, S., Taggart, J.B., Guy, D.R., Bell, J.G., Tocher, D.R. 2012. Hepatic transcriptome analysis
875 of inter-family variability in flesh n-3 long-chain polyunsaturated fatty acid content in Atlantic
876 salmon. BMC Genomics 13, 410.

877 Moren, M., Opstad, I., Hamre, K. 2004. A comparison of retinol, retinal and retinyl ester
878 concentrations in larvae of Atlantic halibut (*Hippoglossus hippoglossus* L.) fed *Artemia* or
879 zooplankton. Aquacult. Nutr.10, 253-259.

880 Mundheim, H., Aksnes, A., Hope, B. 2004. Growth, feed efficiency and digestibility in salmon
881 (*Salmo salar* L.) fed different dietary proportions of vegetable protein sources in combination
882 with two fish meal qualities. Aquaculture 237, 315-331.

883 Olsvik, P.A., Hemre, G.-I., Waagbø, R. 2013. Exploring early micronutrient deficiencies in rainbow
884 trout (*Oncorhynchus mykiss*) by next-generation sequencing technology - from black box to
885 functional genomics. PLoS ONE 11, e0156668.

886 Opstvedt, J., Aksnes, A., Hope, B., Pike, I.H. 2003. Efficiency of feed utilization in Atlantic salmon
887 (*Salmo salar* L.) fed diets with increasing substitution of fish meal with vegetable proteins.
888 Aquaculture 221, 365-379.

889 Ørnsrud, R., Graff, I.E., Høie, S., Totland, G.K., Hemre, G.I. 2002. Hypervitaminosis A in first-
890 feeding fry of the Atlantic salmon (*Salmo salar* L.). Aquacult. Nutr. 8, 7-13.

891 Ørnsrud, R., Lock, E.-J., Waagbø, R., Krossøy, C., Fjelldal, P.-G. 2013. Establishing an upper level
892 of intake for vitamin A in Atlantic salmon (*Salmo salar* L.) postsmolts. Aquacult. Nutr. 19, 651-
893 664.

894 Oxley, A., Tocher, D.R., Torstensen, B.E., Olsen, R.E. 2005. Fatty acid utilisation and metabolism
895 in caecal enterocytes of rainbow trout (*Oncorhynchus mykiss*) fed dietary fish or copepod oil.
896 Biochim. Biophys. Acta Mol. Cell Biol. Lipids 1737, 119-129.

897 Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR.
898 Nucleic Acids Res. 29, e45.

899 Pietschmann, N., Rijntjes, E., Hoeg, A., Stodter, M., Schweizer, U., Seemann, P., Schomburg, L.
900 2014. Selenoprotein P is the essential selenium transporter for bones. Metallomics, 6, 1043-1049.

901 Prabhu, A.J.P., Schrama, J.W., Kaushik, S.J. 2016. Mineral requirements of fish: a systematic review.
902 Rev. Aquacult. 8, 172-219.

903 Prabhu, A.J.P., Schrama, J., Fontagné - Dicharry, S., Mariojouis, C., Surget, A., Bueno, M., Geurden,
904 I., Kaushik, S.J. 2018. Evaluating dietary supply of microminerals as a premix in a complete
905 plant ingredient - based diet to juvenile rainbow trout (*Oncorhynchus mykiss*). Aquacult. Nutr.
906 24, 539-547.

907 Prabhu, A.J.P., Lock, E.-J., Hemre, G.-I., Hamre, K., Espe, M., Olsvik, P.A., Silva, J., Hansen, A.-
908 C., Johansen, J., Sissener, N.H., Waagbø, R. 2019a. Recommendations for dietary level of micro-
909 minerals and vitamin D3 to Atlantic salmon (*Salmo salar*) parr and post-smolt when fed low fish
910 meal diets. PeerJ 7:e6996.

911 Prabhu, A.J.P., Silva, M.S., Kröeckel, S., Holme, M.-H., Ørnsrud, R., Amlund, H., Lock, E.-J.,
912 Waagbø, R. 2019b. Effect of levels and sources of dietary manganese on growth and mineral
913 composition of post-smolt Atlantic salmon fed low fish meal, plant-based ingredient diets.
914 Aquaculture 512, 734287.

915 Pratoomyot, J., Bendiksen, E.Å., Bell, J.G., Tocher, D.R. 2008. Comparison of effects of vegetable
916 oils blended with southern hemisphere fish oil and decontaminated northern hemisphere fish oil
917 on growth performance, composition and gene expression in Atlantic salmon (*Salmo salar* L.).
918 Aquaculture 280, 170-178

919 Remø, S., Hevrøy, E.M., Olsvik, P.A., Fontanillas, R., Breck, O., Waagbø, R. 2014. Dietary histidine
920 requirement to reduce the risk and severity of cataracts is higher than the requirement for growth
921 in Atlantic salmon smolts, independently of the dietary lipid source. Br. J. Nutr. 111, 1759-1772.

922 Seierstad, S.L., Poppe, T.T., Koppang, E.O., Svindland, A., Rosenlund, G., Frøyland, L., Larsen, S.,
923 2005. Influence of dietary lipid composition on cardiac pathology in farmed Atlantic salmon,
924 *Salmo salar* L. J. Fish Dis. 28, 677-690.

925 Seierstad, S.L., Haugland, Ø, Larsen, S., Waagbø, R., Evensen, Ø. 2009. Pro-inflammatory cytokine
926 expression and respiratory burst activity following replacement of fish oil with rapeseed oil in
927 the feed for Atlantic salmon (*Salmo salar* L.). Aquaculture 289, 212-218.

928 Shearer, K., Åsgård, T., Andorsdottir, G., Aas, G. 1994. Whole body elemental and proximate
 929 composition of Atlantic salmon (*Salmo salar*) during the life cycle. J. Fish Biol. 44, 785–797.

930 Shepherd C.J., Jackson. A.J. 2013. Global fishmeal and fish - oil supply: inputs, outputs and markets.
 931 J. Fish Biol. 83, 1046–1066.

932 Sissener, N.H., Julshamn, K., Espe, M., Lunestad, B.T., Hemre, G.-I., Waagbø, R., Måge, A. 2013.
 933 Surveillance of selected nutrients, additives and undesirables in commercial Norwegian fish
 934 feeds in the years 2000-2010. Aquacult. Nutr. 19, 555-572.

935 Taylor, J.F., Vera, L.M., De Santis, C., Lock, E.-J., Espe, M., Skjaerven, K.H., Leeming, D., del Pozo,
 936 J., Mota-Velasco, J., Migaud, H., Hamre, K., Tocher, D.R. 2019. The effect of micronutrient
 937 supplementation on growth and hepatic metabolism in diploid and triploid Atlantic salmon
 938 (*Salmo salar*) parr fed a low marine ingredient diet. Comp. Biochem. Physiol. B 227, 106-121.

939 Torstensen, B.E., Lie Ø., Frøyland, L. 2000. Lipid metabolism and tissue composition in Atlantic
 940 salmon (*Salmo salar* L.) - effects of capelin oil, palm oil, and oleic acid-enriched sunflower oil
 941 as dietary lipid sources. Lipids 35, 653-664.

942 Torstensen, B.E., Bell, J.G., Rosenlund, G., Henderson, R.J., Graff, I.E., Tocher, D.R., Lie, Ø,
 943 Sargent, J.R. 2005. Tailoring of Atlantic salmon (*Salmo salar* L.) flesh lipid composition and
 944 sensory quality by replacing fish oil with a vegetable oil blend. J. Agric. Food Chem. 53, 10166-
 945 10178.

946 Turchini, G.M., Francis, D.S., Senadheera, S.P.S.D., Thanuthong, T., De Silva, S.S. 2011. Fish oil
 947 replacement with different vegetable oils in Murray cod: Evidence of an "omega-3 sparing effect"
 948 by other dietary fatty acids. Aquaculture 315, 250-259.

949 Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., Rozen, S.G. 2012.
 950 Primer3-new capabilities and interfaces. Nucleic Acids Res. 40, e115.

951 Valente, L.P.M., Bower, N.I., Johnston, I.A. 2012. Postprandial expression of growth-related genes
 952 in Atlantic salmon (*Salmo salar* L) juveniles fasted for 1 week and fed a single meal to satiation.
 953 Br. J. Nutr., 108, 2148-2157

954 Vera, L.M., Lock, E.-J., Hamre, K., Migaud, H., Leeming, D., Tocher, D.R., Taylor, J.F. 2019.
 955 Enhanced micronutrient supplementation in low marine diets reduced vertebral malformation in
 956 diploid and triploid Atlantic salmon (*Salmo salar*) parr, and increased vertebral expression of
 957 bone biomarker genes in diploids. Comp. Biochem. Physiol. B. 237, 110327.

958 Waagbø, R. 2008. Reducing production-related diseases in farmed fish. In: Improving Farmed Fish
 959 Quality and Safety. Woodhead Publishing Ltd. 363-398.

960 Waagbø, R. 2006. Feeding and disease resistance in fish. In: Biology of Nutrition in Growing
 961 Animals. Elsevier. (Eds. Mosenthin, R., Zentek, J., Żebrowska, T.) 387-415.

- 962 Waagbø, R., Glette, J., Raa-Nilsen, E., Sandnes, K. 1993. Dietary vitamin C, immunity and disease
963 resistance in Atlantic salmon (*Salmo salar*). Fish Physiol. Biochem. 12, 61-73.
- 964 Waagbø, R., Berntssen, M.H.G., Danielsen, T., Helberg, H., Kleppa, A.L., Berg Lea, T., Rosenlund,
965 G., Tvenning, L., Susort, S., Vikeså, V., Breck, O. 2013. Feeding Atlantic salmon diets with plant
966 ingredients during the seawater phase - a full-scale net production of marine protein with focus
967 on biological performance, welfare, product quality and safety. Aquacult. Nutr. 19, 598-618.
- 968 Watanabe, T., Kiron, V., Satoh, S. 1997. Trace minerals in fish nutrition. Aquaculture 151, 185-207.
- 969 Witten. P. E., Gil-Martens. L., Huysseune. A., Takle. H., Hjelde. K. 2009. Towards a classification
970 and an understanding of developmental relationships of vertebral body malformations in Atlantic
971 salmon (*Salmo salar* L.). Aquaculture 295, 6–14.
- 972 Ytrestøyl, T., Aas, T.S., Åsgård, T. 2015. Utilisation of feed resources in production of Atlantic
973 salmon (*Salmo salar*) in Norway. Aquaculture 448, 365-374.
- 974 Zhang, Z., Zhang, J. Xiao, J. 2014. Selenoproteins and selenium status in bone physiology and
975 pathology Biochim. Biophys. Acta. 1840, 3246–3256

976 **Figure Legends**

977 **Figure 1.** Evolution of growth (increase in weight) through freshwater and seawater phases of Atlantic
978 salmon fed the three experimental diets (mean \pm SD, n=3). The feeding period for each dietary
979 formulation (% FM/FO) is shown in the blocks at the top of the figure. Superscripts denote significant
980 differences between diets at each respective time point.

981
982 **Figure 2.** Percentage fish showing at least one or more radiological spinal deformities ($\geq +1dV$)
983 recorded at smolt (solid bars) and harvest (hatched bars) in the dietary treatments (25 fish per tank /
984 pen, mean \pm SD, n=3). Superscripts denote significant differences between life stage and diet
985 (Kruskall-Wallis, $p < 0.05$).

986

987 **Figure 3.** Functional categories of genes differentially expressed in liver of Atlantic salmon fed diets
988 L2 (A) and L3 (B) compared to fish fed diet L1. Non-annotated genes and features corresponding to
989 the same gene are not represented.

990

991 **Figure 4.** KEGG pathway analysis of genes belonging to the metabolism category in liver of Atlantic
992 salmon fed diets L2 (A) and L3 (B) compared to fish fed diet L1. Bars represent number of up- and
993 down-regulated genes. Different colours indicate different nutrient groups.

994

995

996 **Table 1.** Formulation of the diets in terms of FM and FO inclusions used during the course of the
 997 study
 998

Pellet Size	Fish Size	L1	L2	L3	
2.0 mm	20-75g	FM-15% FO-8%	FM-15% FO-8%	FM-15% FO-8%	FW
3.5 mm	75-250g	FM-15% FO-5%	FM-15% FO-5%	FM-15% FO-5%	} SW
5.0 / 7.0 mm	250-1000g	FM-10% FO-4%	FM-10% FO-4%	FM-10% FO-4%	
9.0 mm	1000-2500g	FM-5% FO-3%	FM-5% FO-3%	FM-5% FO-3%	

999

1000

1001 **Table 2.** Raw material formulation (g.100g diet⁻¹) and proximate composition (analysed) of
1002 experimental diets. ARRAINA Nutrient Package and Amino Acid Premix inclusion for L1/L2/L3
1003 diets respectively.

INGREDIENTS	MARINE PELLET SIZE (mm)		
	3.5	5.0 / 7.0	9.0
Fish Meal ¹	13.00	8.00	3.00
Krill Meal ²	2.00	2.00	2.00
Soy Protein Concentrate ³	15.58	15.40	12.58
Wheat Gluten ⁴	12.46	13.09	10.70
Maize Gluten	3.12	3.08	2.52
Pea Protein Concentrate ⁵	12.46	13.09	10.70
Wheat ⁶	17.86	10.00	13.33
Sunflower Expeller	-	6.32	6.77
Fish Oil ⁷	5.00	4.00	3.00
Rapeseed oil ⁴	7.65	10.61	16.17
Linseed oil	1.91	2.55	3.84
Palm kernel oil	4.78	6.38	9.61
ARRAINA Nutrient Package ^{8†‡}	0.75 / 1.50 / 3.00	0.75 / 1.50 / 3.00	0.75 / 1.50 / 3.00
Monosodium Phosphate	2.03	-	-
Monocalcium phosphate	-	2.05	1.46
Amino acid Premix ^{9,*}	2.14 / 2.34 / 2.32	2.07 / 2.11 / 2.16	2.15 / 2.18 / 2.24
Yttrium	0.05	0.05	0.05
Lucantin Pink	0.06	0.06	0.06
Water change	-1.98	-0.52	0.28
Proximate Composition (Analysed)			
Moisture (%)	5.6	7.2	6.7
Crude lipid (%)	24.6	25.7	31.5
Crude protein (%)	43.8	41.4	41.2
Ash (%)	6.0	5.3	5.3
Energy (MJ / kg)	23.2	23.9	25.4

1004 ¹Feed Services, Bremen, Germany; ²Aker Biomarine, Norway; ³Caramuru, Brazil; ⁴Cargill,
1005 Germany; ⁵Agrient, Germany; ⁶WN Lindsey, UK; ⁷ED & F Man, Germany; ⁸DSM, Netherlands;
1006 ⁹Evonik, Germany; †Added as components of the nutrient package (NP), and times requirement based
1007 on NRC (2011) minimum requirement for Atlantic salmon and modified according to Hamre et al.,
1008 (2016), diet L1 achieving assumed 100 % minimum requirement; ‡ NP *Balanced for lysine,
1009 methionine, threonine and valine. Contains antioxidant.

1010
1011

1012 **Table 3.** Analysed micronutrient contents (mg.kg⁻¹) within the nutrient package (NP): selected
1013 amino acids (histidine and taurine), minerals, vitamins and cholesterol. * Footnote denotes vitamin
1014 and mineral sources.

PREMIX INGREDIENTS	NP		
	100%	200%	400%
Vitamin A	3.79	7.58	15.16
Vitamin D3	0.05	0.10	0.20
Vitamin E	102.44	204.88	409.76
Vitamin K3	9.82	19.64	39.28
Thiamine	2.67	5.34	10.68
Riboflavin	8.30	16.60	33.20
B6	4.77	9.54	19.08
B12	0.25	0.50	1.00
Niacin	24.80	49.60	99.20
Pantothenic Acid	17.15	34.30	68.60
Folic Acid	2.82	5.64	11.28
Biotin	0.14	0.28	0.56
Vitamin C	80	160	320
Calcium	0.4	0.8	1.6
Cobalt	0.94	1.88	3.76
Iodine	0.67	1.34	2.68
Selenium	0.23	0.46	0.92
Iron	32.64	65.28	130.56
Manganese	12.03	24.06	48.12
Copper	3.24	6.48	12.96
Zinc	66.92	133.84	267.68
Taurine	2450	4900	9800
Histidine	1400	2800	5600
Cholesterol	1100	2200	4400

1015 ***Vitamin & Mineral Source:** **Vit A**, Retinyl acetate; **Vit D₃**, Cholecalciferol; **Vit E**, all-rac-alpha-tocopheryl acetate;
1016 **Vit K**, menadione nicotinamide bisulphite; **Thiamine**, thiamine mononitrate; **Riboflavin**, Riboflavin; **B6**, Vitamin B6 /
1017 pyridoxine hydrochloride; **B12**, cyanocobalamin; **Niacin**, nicotinic acid; **Pantothenic acid**, Calcium-D-pantothenate;
1018 **Folic acid**, Folic acid; **Biotin**, Biotin; **Vit C**, L-ascorbic acid; **Co**: Cobalt carbonate; **I**, Calcium iodate anhydrous; **Se**,
1019 Sodium selenite; **Fe**, ferrous sulphate monohydrate; **Mn**, Manganous oxide; **Cu**, Cupric sulphate pentahydrate; **Zn**,
1020 Zinc oxide.
1021

1022 **Table 4.** Analysed concentrations of selected amino acids (taurine, histidine and methionine g.kg⁻¹)
1023 macro-minerals (calcium, magnesium and phosphorous, g.kg⁻¹) micro-minerals and vitamins
1024 (mg.kg⁻¹) of the experimental diets. Nutrients added at graded levels to the feeds are shown with an
1025 asterisk.

	DIET			
	L1	L2	L3	NRC 2011 [‡]
Vitamin A*	3.7	5.2	12.2	0.75 ^a
Vitamin D3*	0.15	0.19	0.19	0.04 ^a
Vitamin E*	241.5	364.0	436.5	60 ^b
Vitamin K3*	0.71	1.51	2.70	<10 ^b
Thiamin*	4.5	7.1	8.8	1 ^a ,
Riboflavin*	17.2	27.8	33.5	4 ^a ,
Vitamin B6*	12.8	16.8	21.3	5 ^b
Vitamin B12*	0.18	0.35	0.67	NT
Niacin*	73.0	112.0	148.0	10 ^a ,
Pantothenic acid*	24.0	58.0	44.0	20 ^a ,
Folic acid*	6.53	9.69	11.67	1 ^a
Biotin*	0.51	0.72	0.74	0.15 ^a
Vitamin C*	183.0	251.0	409.0	20 ^b
Cobalt*	0.18	0.22	0.32	NT
Iodine*	n.a	n.a	n.a	1.1 ^a
Selenium*	1.13	1.48	1.65	0.15 ^a
Iron*	330.0	358.0	403.0	30-60 ^b
Manganese*	42.0	53.0	86.0	10 ^b
Copper*	11.8	14.8	22.8	5 ^b
Zinc*	94.0	156.0	330.0	37 ^b
Taurine*	2.6	4.4	10.1	NR ^b
Methionine	9.7	9.9	10.3	7.0 ^b
Histidine*	11.4	13.1	17.1	8.0 ^b
Calcium*	6.7	7.1	8.2	NR ^{b*}
Magnesium	1.7	1.7	1.7	0.4 ^b
Phosphorus	12.7	12.5	12.5	8.0 ^b
Cholesterol*	n.a	n.a	n.a	NR

1026 [‡]Current NRC, 2011 minimum requirement recommendations determined in ^a rainbow trout, ^b
1027 Atlantic salmon are shown for comparison. n.a. not analysed; NR* no requirement freshwater; NT,
1028 not tested.
1029

Table 5. Growth and morphometric indices recorded at the end of the fresh and seawater phases of Atlantic salmon fed low marine ingredient diets (FM 15% / FO 8%) with differing micronutrient supplementation levels (L1 = 100 %, L2 = 200 % and L3 = 400 % NRC Premix). Superscripts denote significant differences between diets (mean \pm SD, n=3, $p < 0.05$, one-way ANOVA)

	DIET			<i>p</i> values
	L1	L2	L3	
FRESHWATER				
Initial Parr Wt. (g)	38.2 ± 0.6	38.0 ± 0.6	38.1 ± 0.5	0.946
Final Smolt Wt. (g)	68.0 ± 0.5 ^c	77.4 ± 0.5 ^a	73.8 ± 0.8 ^b	0.0001
Condition Factor (K)	1.29 ± 0.00 ^c	1.19 ± 0.00 ^b	1.18 ± 0.00 ^a	0.0001
TGC _{wt}	1.29 ± 0.01 ^c	1.59 ± 0.01 ^a	1.48 ± 0.02 ^b	0.0001
bFCR	0.90 ± 0.01 ^a	0.71 ± 0.01 ^c	0.76 ± 0.01 ^b	0.0001
VSI (%)	8.23 ± 0.10	8.21 ± 0.18	8.44 ± 0.35	0.456
HSI (%)	1.28 ± 0.06 ^a	1.04 ± 0.01 ^b	1.15 ± 0.06 ^b	0.003
Mortality (%)	0.2 ± 0.2	0.3 ± 0.1	0.0 ± 0.0	0.646
SEAWATER				
Final Wt. (g)	2127.0 ± 59.0 ^b	2381.0 ± 82.0 ^a	2385.0 ± 37.0 ^a	0.041
Condition Factor (K)	1.37 ± 0.01 ^b	1.43 ± 0.02 ^{ab}	1.46 ± 0.03 ^a	0.049
TGC _{wt}	2.87 ± 0.04 ^b	2.98 ± 0.05 ^{ab}	3.02 ± 0.02 ^a	0.045
bFCR	1.11 ± 0.01 ^a	1.08 ± 0.01 ^b	1.08 ± 0.01 ^b	0.049
VSI (%)	8.66 ± 0.10	8.98 ± 0.11	8.75 ± 0.35	0.106
HSI (%)	1.08 ± 0.06	1.02 ± 0.01	1.10 ± 0.04	0.426
% Mortality	0.95 ± 0.76	2.64 ± 2.25	0.54 ± 0.13	0.594
% Maturation	9.99 ± 1.64	5.94 ± 0.48	9.84 ± 0.74	0.063

Table 6. Protein, lipid, ash (g.kg⁻¹ wet wt.), mineral content (mg.kg⁻¹ wet wt.), retention (%) in whole body, and protein, lipid and fatty acid composition (%) in muscle (Norwegian Quality Cut, NQC) of Atlantic salmon at end of trial. Superscripts denote significant differences between diets when variances were homogenous (mean \pm SD, n=3, p<0.05, ANOVA), when non-homogenous variances: (mean \pm SD, n=3, $p < 0.5$, Kruskal-Wallis).

	DIET			<i>p</i> values
	L1	L2	L3	
WHOLE BODY‡				
Protein	16.6 ± 0.3	16.8 ± 0.2	16.6 ± 0.5	0.61
Lipids	19.1 ± 1.4	20.0 ± 0.7	19.7 ± 0.8	0.59
Ash	1.5 ± 0.11 ^b	1.5 ± 0.11 ^b	1.7 ± 0.11 ^a	0.045
Zn	9.0 ± 1.0 ^c	18.0 ± 3.0 ^b	29.0 ± 3.0 ^a	0.0003
Mn	1.5 ± 0.5	1.5 ± 0.3	1.3 ± 0.8	0.89
Fe	7.1 ± 2.5	8.1 ± 0.4	8.9 ± 1.9	0.53
Cu	2.7 ± 0.1	2.8 ± 0.2	2.8 ± 0.4	0.81
Se	0.18 ± 0.01 ^b	0.22 ± 0.02 ^b	0.31 ± 0.02 ^a	0.0003
Mg	273.0 ± 15.0	273.0 ± 6.0	273.0 ± 15.0	1.00
Ca	2500 ± 1000	2800 ± 400	2600 ± 1700	0.50
P	3000 ± 200	3400 ± 200	3400 ± 800	0.94
WHOLE BODY RETENTION				
Protein	37.0 ± 1.0	37.0 ± 1.0	37.0 ± 2.0	0.63
Lipids	62.0 ± 5.0	68.0 ± 2.0	67.0 ± 2.0	0.12
Ash	28.0 ± 1.6	28.7 ± 1.3	29.1 ± 1.4	0.83
Zn	8.3 ± 0.7	11.3 ± 2.2	7.7 ± 1.0	0.18
Mn	2.8 ± 0.9	2.6 ± 0.6	1.3 ± 0.8	0.11
Fe	1.8 ± 0.6	1.8 ± 0.1	2.0 ± 0.4	0.77
Cu	19.7 ± 0.5 ^a	16.9 ± 1.0 ^{ab}	10.4 ± 1.5 ^b	0.0001
Se	12.5 ± 0.5 ^b	12.2 ± 1.0 ^b	17.9 ± 1.5 ^a	0.001
Mg	14.0 ± 1.0	14.9 ± 0.4	15.2 ± 1.1	0.29
P	25.0 ± 7.0	26 ± 1.0	26.0 ± 6.0	1.00
Ca	38.3 ± 23.0	33.4 ± 6.0	27.4 ± 19.0	0.92
NQC				
Protein	19.8 ± 0.5	20.2 ± 0.6	20.0 ± 0.4	0.62
Lipid	13.9 ± 0.3 ^b	13.5 ± 0.1 ^b	15.0 ± 0.5 ^a	0.004
Total PUFA	31.4 ± 0.4	31.4 ± 0.3	30.8 ± 0.2	0.40
n-6 LC PUFA	17.5 ± 0.2 ^a	17.1 ± 0.1 ^{ab}	16.9 ± 0.1 ^b	0.050
n-3 LC PUFA	13.7 ± 0.2	14.0 ± 0.2	13.7 ± 0.1	0.45
EPA	1.5 ± 0.00	1.6 ± 0.0	1.6 ± 0.0	0.056
DHA	1.82 ± 0.05 ^{ab}	1.90 ± 0.06 ^a	1.60 ± 0.07 ^b	0.036
Total Carotenoid	4.27 ± 0.17	3.87 ± 0.29	3.96 ± 0.33	0.59

[‡]Initial Body compositions at smolt sea transfer (L1 / L2 / L3):

Protein: (16.24 \pm 0.16 / 16.49 \pm 0.19 / 16.7 \pm 0.11); **Lipids:** (11.30 \pm 0.24 / 11.8 \pm 0.35 / 11.6 \pm 0.46); **Ash:** (2.23 \pm 0.12 / 2.25 \pm 0.14 / 2.15 \pm 0.07) ; **Zn:** (25 \pm 4.5 / 30 \pm 0.2 / 44 \pm 4.5); **Mn:** (2.2 \pm 0.2 / 2.1 \pm 0.6 / 1.9 \pm 0.4); **Fe:** (10 \pm 0.4 / 14 \pm 5.9 / 11 \pm 0.1); **Cu:** (1.2 \pm 0.1 / 1.4 \pm 0.1 / 1.5 \pm 0.1); **Se:** (0.29 \pm 0.01 / 0.32 \pm 0.02 / 0.37 \pm 0.02); **Mg:** (355 \pm 21 / 345 \pm 21 / 350 \pm 0); **Ca:** (3300 \pm 707 / 3050 \pm 1061 / 2900 \pm 849); **P:** (4750 \pm 354 / 4650 \pm 566 / 4600 \pm 354).

1048 **Table 7.** Amino acid retention (%) in Atlantic salmon fed the three plant-based diets. Superscripts
1049 denote significant differences between diets (mean \pm SD, n=3, p<0.05, ANOVA).

1050

	DIET			<i>p</i> values
	L1	L2	L3	
His	31.2 \pm 0.3 ^a	30.2 \pm 0.5 ^a	26.8 \pm 0.9 ^b	0.005
OH-pro	87.5 \pm 10.7	75.6 \pm 1.9	101.2 \pm 18.8	0.41
Arg	33.0 \pm 0.4	33.4 \pm 0.3	33.4 \pm 1.4	0.94
Ser	31.5 \pm 0.3	32.2 \pm 0.4	32.6 \pm 1.1	0.61
Gly	46.8 \pm 1.1	44.4 \pm 0.7	46.5 \pm 2.6	0.56
Asp	43.5 \pm 1.1	44.1 \pm 0.8	43.0 \pm 1.0	0.74
Glu	22.7 \pm 0.2	22.8 \pm 0.5	22.1 \pm 0.9	0.71
Thr	42.1 \pm 0.2	43.0 \pm 0.2	42.8 \pm 1.1	0.68
Ala	50.2 \pm 1.1	50.6 \pm 0.2	50.5 \pm 1.4	0.95
Pro	20.7 \pm 0.4	20.4 \pm 0.1	20.8 \pm 0.9	0.86
Tyr	36.6 \pm 0.4	37.7 \pm 0.3	36.4 \pm 1.1	0.55
Met	43.6 \pm 0.3	44.9 \pm 0.3	42.0 \pm 1.6	0.19
Lys	49.4 \pm 1.4 ^a	49.3 \pm 0.4 ^a	45.7 \pm 0.2 ^b	0.04
Val	41.0 \pm 0.5	41.7 \pm 1.1	39.4 \pm 0.7	0.18
Ile	38.4 \pm 0.5	39.2 \pm 1.1	37.0 \pm 0.6	0.21
Leu	35.5 \pm 0.1	36.0 \pm 0.1	35.5 \pm 0.6	0.53
Phe	27.6 \pm 0.3	28.5 \pm 0.3	28.3 \pm 1.2	0.71

1051
1052
1053

1054 **Table 8.** Non-protein N metabolites in muscle ($\mu\text{mol.g}^{-1}$ wet wt.). Superscripts denote significant
1055 differences between diets (mean \pm SD, n=3, p<0.05, ANOVA).

1056

	DIET			<i>p</i> values
	L1	L2	L3	
Taurine	0.94 \pm 0.01	1.13 \pm 0.22	1.00 \pm 0.02	0.58
Urea	0.53 \pm 0.02	0.52 \pm 0.02	0.55 \pm 0.06	0.88
Asp	0.09 \pm 0.00	0.10 \pm 0.00	0.12 \pm 0.00	0.15
OH-pro	0.27 \pm 0.02 ^a	0.25 \pm 0.01 ^{ab}	0.16 \pm 0.01 ^b	0.03
Thr	0.43 \pm 0.02	0.40 \pm 0.01	0.43 \pm 0.02	0.62
Ser	0.14 \pm 0.01 ^a	0.09 \pm 0.01 ^b	0.06 \pm 0.01 ^b	0.0005
Glu	0.99 \pm 0.06	0.93 \pm 0.02	0.87 \pm 0.11	0.51
Gln	0.14 \pm 0.02	0.23 \pm 0.03	0.20 \pm 0.02	0.08
Sarcosine	0.08 \pm 0.02	0.11 \pm 0.02	0.09 \pm 0.02	0.64
Pro	0.53 \pm 0.02	0.48 \pm 0.06	0.29 \pm 0.02	0.06
Gly	2.37 \pm 0.10 ^a	1.46 \pm 0.08 ^b	0.98 \pm 0.06 ^c	0.0006
Ala	2.83 \pm 0.08	2.78 \pm 0.11	2.47 \pm 0.12	0.11
Citruline	0.03 \pm 0.01	0.03 \pm 0.00	0.03 \pm 0.00	0.79
Alpha amino butyric acid	0.04 \pm 0.00	0.05 \pm 0.00	0.04 \pm 0.00	0.22
Val	0.28 \pm 0.00	0.29 \pm 0.01	0.29 \pm 0.01	0.81
Met	0.10 \pm 0.00	0.11 \pm 0.00	0.10 \pm 0.00	0.18
Cystathionine	0.01 \pm 0.00	0.01 \pm 0.00	0.01 \pm 0.00	0.42
Ile	0.13 \pm 0.00	0.14 \pm 0.00	0.13 \pm 0.00	0.73
Leu	0.23 \pm 0.01	0.25 \pm 0.01	0.24 \pm 0.01	0.30
Tyr	0.17 \pm 0.02	0.23 \pm 0.03	0.18 \pm 0.02	0.35
Beta Ala	0.35 \pm 0.02	0.39 \pm 0.04	0.30 \pm 0.01	0.10
Phe	0.07 \pm 0.00	0.09 \pm 0.00	0.07 \pm 0.00	0.05
Ammonia	4.94 \pm 0.15	4.80 \pm 0.07	4.66 \pm 0.05	0.22
Ornithine	0.02 \pm 0.00	0.01 \pm 0.00	0.01 \pm 0.00	0.30
Lys	0.81 \pm 0.12	0.86 \pm 0.04	0.95 \pm 0.13	0.62
1-methyl His	0.05 \pm 0.00	0.06 \pm 0.00	0.06 \pm 0.01	0.42
His	1.53 \pm 0.20 ^c	2.33 \pm 0.07 ^b	4.10 \pm 0.09 ^a	0.0001
Anserine	26.58 \pm 0.21	27.77 \pm 0.20	27.46 \pm 0.72	0.23
Carnosine	0.37 \pm 0.05	0.57 \pm 0.08	0.42 \pm 0.05	0.12
Arg	0.16 \pm 0.01	0.17 \pm 0.00	0.18 \pm 0.01	0.51

1057

1058

1059

1060 **Table 9.** Tissue concentrations (mg.kg⁻¹ wet wt.) and retention (%) of water soluble vitamins.
1061 Superscripts denote significant differences between diets when variances were homogenous (mean
1062 \pm SD, n=3, p<0.05, ANOVA), when non-homogenous variances: (mean \pm SD, n=3, $p < 0.5$,
1063 Kruskal-Wallis).

1064
1065

	DIET			<i>p</i> values
	L1	L2	L3	
WHOLE BODY				
Vitamin C	45.0 ± 4.0 ^b	60.0 ± 6.0 ^a	68.0 ± 5.0 ^a	0.004
Thiamin	1.2 ± 0.1	1.3 ± 0.1	1.2 ± 0.1	0.3
Riboflavin	2.3 ± 0.1	2.4 ± 0.21	2.3 ± 0.2	0.57
Pantothenic acid	9.3 ± 0.7 ^b	12.0 ± 0.0 ^a	12.3 ± 0.6 ^a	6*10 ⁻⁴
Niacin	32 ± 3 ^c	46 ± 2 ^b	56 ± 4 ^a	4*10 ⁻⁴
Folic acid	0.2 ± 0.0	0.3 ± 0.04	0.3 ± 0.02	0.25
Biotin	0.1 ± 0.0	0.2 ± 0.2	0.1 ± 0.01	0.25
Vitamin B6	0.3 ± 0.02	0.5 ± 0.03	0.5 ± 0.03	0.05
LIVER				
Vitamin C	53.0 ± 1.0	91 ± 28	44 ± 24	0.06
Biotin	2.8 ± 0.6	2.7 ± 0.5	3.1 ± 0.6	0.37
Vitamin B12	0.6 ± 0.0	0.5 ± 0.1	0.6 ± 0.1	0.28
Folic acid	16.8 ± 0.3	14.8 ± 1.2	15.9 ± 0.7	0.06
Niacin	70.0 ± 3.0	72.0 ± 6.0	79.0 ± 12.0	0.42
Riboflavine	12.7 ± 0.6	12.3 ± 0.6	12.3 ± 0.6	0.73
GILL				
Pantothenic acid	8.5 ± 1.3	10.9 ± 1.9	12.0 ± 2.1	0.17
MUSCLE				
Vitamin B6	5.8 ± 0.2	6.9 ± 0.8	6.9 ± 1.0	0.17
RETENTION IN WHOLE BODY (%)				
Thiamin	21.2 ± 2.2	18.3 ± 1.0	18.6 ± 1.0	0.113
Riboflavin	9.8 ± 0.1	7.1 ± 0.5	6.5 ± 0.5	0.051
Niacin	37.4 ± 3.3	39.2 ± 2.8	33.1 ± 2.2	0.079
Biotin	11.4 ± 0.0	9.0 ± 0.3	10.1 ± 1.5	0.148
Folic acid	3.5 ± 0.1	2.3 ± 0.4	1.6 ± 0.1	0.027

1066
1067

1068
1069

1070 **Table 10.** Tissue concentrations (mg.kg⁻¹ wet wt.) and retention (%) of lipid soluble vitamins.
1071 Superscripts denote significant differences between diets when variances were homogenous
1072 (p<0.05, ANOVA), when non-homogenous variances: (p < 0.5, Kruskal-Wallis) (mean ± SD, n=3
1073 tanks).

1074
1075

	DIET			<i>p</i> value
	L1	L2	L3	
WHOLE BODY				
Dry matter %	41.0 ± 1.0	40.0 ± 1.0	41.0 ± 1.0	0.40
Vitamin A ₁	0.20 ± 0.16	0.58 ± 0.13	2.23 ± 0.42	0.04
Vitamin A ₂	0.39 ± 0.06 ^c	0.83 ± 0.12 ^b	1.77 ± 0.25 ^a	0.0001
Vitamin K (K ₁)	30.0 ± 6.0	32.0 ± 1.0	30.0 ± 0.0	0.25
Vitamin K (MK ₄)	73.0 ± 6.0 ^b	84.0 ± 8.0 ^{ab}	100.0 ± 7.0 ^a	0.008
Vitamin E (α-TOH)	28.0 ± 5.0 ^c	41.0 ± 5.0 ^b	57.0 ± 2.0 ^a	0.0005
Vitamin E (γ-TOH)	12.5 ± 3.8	12.1 ± 1.4	11.1 ± 0.5	0.56
MUSCLE				
Vitamin D ₃	0.023 ± 0.006 ^c	0.030 ± 0.000 ^b	0.043 ± 0.006 ^a	0.01
Vitamin E (α-TOH)	17 ± 2 ^c	24 ± 1 ^b	36 ± 3 ^a	0.0001
Vitamin E (γ -TOH)	9.9 ± 1.7	9.8 ± 0.8	9.2 ± 0.5	ns
WHOLE BODY RETENTION				
Vitamin E (α-TOH)	8.8 ± 1.7 ^a	9.2 ± 0.9 ^{ab}	11.8 ± 0.6 ^b	0.03
Vitamin E (γ -TOH)	34.0 ± 10.0	30.0 ± 3.0	30.0 ± 1.0	0.60


1076
1077
1078

1079 **Table 11.** Relative expression of diets L2 and L3 to L1 for genes involved in lipid and fatty acid
1080 (FA) metabolism (liver) and protein turnover (muscle) in Atlantic salmon fed the experimental diets
1081 (mean \pm SEM, n=3). Superscripts denote significant differences between diets.

GENE	DIET			<i>p</i> value
	L1	L2	L3	
LIPID AND FA METABOLISM (Liver)				
Apolipoprotein B-100 (<i>apoB</i>)	1.00 ± 0.09	1.68 ± 0.10	1.68 ± 0.31	0.890
Acetyl-CoA carboxylase (<i>acc</i>)	1.00 ± 0.13	1.16 ± 0.22	1.23 ± 0.18	0.520
Cholesterol 7 alpha-hydroxylase (<i>Cyp7a1</i>)	1.00 ± 0.40	0.64 ± 0.15	1.14 ± 0.55	0.572
Fatty acyl elongase 2 (<i>elovl2</i>)	1.00 ± 0.26 ^{ab}	1.19 ± 0.20 ^a	0.56 ± 0.15 ^b	0.032
Fatty acyl elongase 5 isoform b (<i>elovl5</i>)	1.00 ± 0.07	1.07 ± 0.14	0.84 ± 0.16	0.454
Fatty acyl elongase 6 (<i>elovl6</i>)	1.00 ± 0.50	1.17 ± 0.78	0.91 ± 0.39	0.577
Delta-6 fatty acyl desaturase isoform a (<i>fads2d6a</i>)	1.00 ± 0.34 ^a	0.36 ± 0.09 ^b	0.44 ± 0.12 ^{ab}	0.035
S-acyl fatty acid synthase thioesterase (<i>fas</i>)	1.00 ± 0.14	0.90 ± 0.09	1.00 ± 0.10	0.671
1-CARBON METABOLISM (Liver)				
Cystathionine-β-synthase (<i>cbs</i>)	1.00 ± 0.26	0.91 ± 0.31	0.86 ± 0.26	0.854
Betaine-homocysteine S-methyltransferase (<i>bhmt</i>)	1.00 ± 0.13	0.91 ± 0.13	1.09 ± 0.13	0.534
Methionine adenosyltransferase1 (<i>mat1</i>)	1.00 ± 0.15	0.70 ± 0.15	1.10 ± 0.13	0.271
Growth hormone receptor (<i>ghr</i>)	1.00 ± 0.40 ^a	0.36 ± 0.11 ^b	0.79 ± 0.32 ^{ab}	0.043
IMMUNE FUNCTION (Liver)				
Interleukin 8 (<i>il8</i>)	1.00 ± 0.20	0.74 ± 0.45	0.94 ± 0.17	0.640
Interleukin 1 beta (<i>il1b</i>)	1.00 ± 0.20	1.10 ± 0.17	1.14 ± 0.12	0.663
PROTEIN TURNOVER (Muscle)				
Autophagy-related protein 4 homolog B (<i>atg4b</i>)	1.00 ± 0.09	1.28 ± 0.20	1.24 ± 0.11	0.401
Autophagy related 12 (<i>atg12</i>)	1.00 ± 0.08 ^b	1.04 ± 0.05 ^{ab}	1.29 ± 0.09 ^a	0.029
GABA(A) receptor-associated protein (<i>gabarap</i>)	1.00 ± 0.05 ^b	1.24 ± 0.06 ^{ab}	1.47 ± 0.09 ^a	0.001
Muscle-specific RING finger protein 1a (<i>murf1a</i>)	1.00 ± 0.04 ^b	1.11 ± 0.17 ^{ab}	1.19 ± 0.08 ^a	0.049
Muscle-specific RING finger protein 1b (<i>murf1b</i>)	1.00 ± 0.09	1.01 ± 0.18	1.51 ± 0.25	0.057
Microtubule-associated proteins 1A/1B light chain 3B precursor (<i>mlp3b</i>)	1.00 ± 0.06 ^b	1.20 ± 0.09 ^{ab}	1.48 ± 0.07 ^a	0.002
GROWTH & GLUCOSE TRANSPORT (Muscle)				
Growth hormone receptor (<i>ghr</i>)	1.00 ± 0.12	0.73 ± 0.16	0.98 ± 0.15	0.427
Insulin-like growth factor 1 (<i>igf1</i>)	1.00 ± 0.27	1.17 ± 0.33	1.43 ± 0.38	0.251
Glucose transporter type 4 (<i>glut4</i>)	1.00 ± 0.06	0.95 ± 0.14	0.98 ± 0.09	0.824
IMMUNE FUNCTION (Muscle)				
Interleukin 1 beta (<i>il1b</i>)	1.00 ± 0.33	1.04 ± 0.64	0.80 ± 0.11	0.488

1083
1084
1085
1086
1087

1088 Fig.1.



1089
1090
1091

Fig. 2.





Fig 3.

A



B





Fig 4.

A



B

