



OPEN

# The role of aspartic acid in reducing coral calcification under ocean acidification conditions

Celeste Kellock<sup>1,6</sup>, Catherine Cole<sup>1,7</sup>, Kirsty Penkman<sup>2</sup>, David Evans<sup>1,8</sup>, Roland Kröger<sup>3</sup>, Chris Hintz<sup>4</sup>, Ken Hintz<sup>5</sup>, Adrian Finch<sup>1</sup> & Nicola Allison<sup>1✉</sup>

Biomolecules play key roles in regulating the precipitation of  $\text{CaCO}_3$  biominerals but their response to ocean acidification is poorly understood. We analysed the skeletal intracrystalline amino acids of massive, tropical *Porites* spp. corals cultured over different seawater  $\text{pCO}_2$ . We find that concentrations of total amino acids, aspartic acid/asparagine (Asx), glutamic acid/glutamine and alanine are positively correlated with seawater  $\text{pCO}_2$  and inversely correlated with seawater pH. Almost all variance in calcification rates between corals can be explained by changes in the skeletal total amino acid, Asx, serine and alanine concentrations combined with the calcification media pH (a likely indicator of the dissolved inorganic carbon available to support calcification). We show that aspartic acid inhibits aragonite precipitation from seawater in vitro, at the pH, saturation state and approximate aspartic acid concentrations inferred to occur at the coral calcification site. Reducing seawater saturation state and increasing [aspartic acid], as occurs in some corals at high  $\text{pCO}_2$ , both serve to increase the degree of inhibition, indicating that biomolecules may contribute to reduced coral calcification rates under ocean acidification.

Tropical coral skeletons are major components of coral reefs and provide substrates and habitat spaces for fisheries and protection from wave erosion for coastal communities<sup>1</sup>. Ocean acidification, caused by rising seawater  $\text{pCO}_2$ , typically suppresses the calcification rates of tropical corals and threatens the formation of these structures<sup>2</sup>. Coral skeletons are formed from semi-isolated extracellular calcification media (ECM) contained between the base of the coral tissues and the underlying skeletons<sup>3</sup>. Ocean acidification reduces the pH of this ECM (termed  $\text{pH}_{\text{ECM}}$ ), shifts the dissolved inorganic carbon (DIC) equilibrium to the detriment of  $\text{CO}_3^{2-}$  and likely reduces the ECM saturation state ( $\Omega$ ), a measure of both the  $[\text{CO}_3^{2-}]$  and  $[\text{Ca}^{2+}]$  available for mineral precipitation<sup>4</sup>. Coral skeletons are composite materials, composed of aragonite (a calcium carbonate polymorph) and organic macromolecules which are disseminated in the mineral phase at the nanoscale<sup>5</sup>. This intracrystalline skeletal organic matrix (SOM) include proteins, sugars, polysaccharides and lipids and is implicated in the control of mineral precipitation<sup>6</sup>. For example, primary coral cell cultures produce extracellular organic materials on which aragonite precipitates<sup>7</sup> while SOM extracted from tropical, Mediterranean and deep sea coral skeletons affects the precipitation rate, morphology and polymorphism of  $\text{CaCO}_3$  precipitated in vitro<sup>8–10</sup>. Organic fibrils at the coral calcification site concentrate  $\text{Ca}^{2+}$ <sup>11</sup> and several proteins, lipids and macromolecules resolved from coral SOMs are capable of binding  $\text{Ca}^{2+}$ <sup>12,13</sup>. Aspartic acid is typically the most abundant amino acid in the coral SOM<sup>12,14</sup> and influences multiple stages of  $\text{CaCO}_3$  nucleation and precipitation<sup>15,16</sup>. L-aspartic acid forms complexes with  $\text{Ca}^{2+}$ , probably via the negatively charged carboxylic acid group ( $\text{COO}^-$ ) on the side chain, and this may provide the mechanism to control  $\text{CaCO}_3$  precipitation and morphology<sup>16</sup>.

Resolving how the organic component of the skeleton responds to increases in seawater  $\text{pCO}_2$  is critical to understand the effects of ocean acidification on coral accretion. Increasing seawater  $\text{pCO}_2$  increases the

<sup>1</sup>School of Earth and Environmental Sciences, University of St. Andrews, St. Andrews KY16 9AL, UK. <sup>2</sup>Department of Chemistry, University of York, York, UK. <sup>3</sup>Department of Physics, University of York, York, UK. <sup>4</sup>Department of Marine and Environmental Sciences, Savannah State University, Savannah, GA, USA. <sup>5</sup>Department of Electrical and Computer Engineering, George Mason University, Fairfax, VA, USA. <sup>6</sup>Present address: Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, Stirling FK9 4LA, Scotland, UK. <sup>7</sup>Present address: Centre of Science Communication, University of Otago, Dunedin 9016, New Zealand. <sup>8</sup>Present address: Institute of Geosciences, Goethe University Frankfurt, Frankfurt am Main, Germany. ✉email: na9@st-andrews.ac.uk

concentration of skeletal protein observed in coral skeletons<sup>17</sup> and is inferred to increase skeletal organic content<sup>18</sup>. Changes in  $\text{pH}_{\text{ECM}}$  in response to increasing seawater  $\text{pCO}_2$ , may alter the net negative charge of biomolecules and thereby influence their control of  $\text{CaCO}_3$  precipitation<sup>19</sup>. Here we identify large variations in the amino acid compositions of a suite of massive *Porites* spp. corals cultured over a range of seawater  $\text{pCO}_2$ <sup>20–22</sup>. We find that amino acid concentrations are significantly correlated with coral calcification rates and we explore how aspartic acid, the most prevalent skeletal amino acid, affects skeleton formation by precipitating aragonite in vitro, under fluid conditions analogous to those of the coral calcification site.

## Results and discussion

**Intracrystalline amino acids and coral calcification.** We analysed the intracrystalline amino acids of the skeletons of 3 genotypes of massive *Porites* spp. corals cultured at 25 °C and at 3 different seawater  $\text{pCO}_2$  conditions namely 180  $\mu\text{atm}$ , 400  $\mu\text{atm}$  and 750  $\mu\text{atm}$ <sup>20–22</sup> (Fig. 1, Table S1). Concentrations of total amino acids, aspartic acid/asparagine (termed Asx), glutamic acid/glutamine (termed Glx) and alanine exhibit significant positive correlations with seawater  $\text{pCO}_2$  and inverse correlations with seawater pH (Table S2). Strongest correlations are observed between these seawater parameters and Asx and are illustrated in Fig. 2a, b. No significant correlations were observed between concentrations of any skeletal amino acids and either  $\text{pH}_{\text{ECM}}$  or  $[\text{H}^+]_{\text{ECM}}$ .

Significant inverse linear correlations occur between coral calcification rate (reported previously<sup>20</sup>) and skeletal Asx (Fig. 2c), serine, alanine, Glx and total amino acids (Figure S1, Table S2). Multiple linear regression analysis indicate that a high degree of variance in calcification rate (the dependent variable) is correlated with these amino acid concentrations (the independent variables,  $p = 0.016$ ,  $r^2 = 0.90$ , see Table S3 for individual parameters). For comparison  $r^2 = 0.57–0.60$  for regressions between coral calcification rate and seawater  $\text{pCO}_2$  and seawater pH. Precipitation rates of inorganic aragonites are highly dependent on the seawater saturation state,  $\Omega$  (reflecting the availability of  $\text{CO}_3^{2-}$  for incorporation in the  $\text{CaCO}_3$  precipitate)<sup>23</sup>. The  $[\text{CO}_3^{2-}]$  of the calcification fluid is a function of total DIC and pH. Adding the  $\text{pH}_{\text{ECM}}$  for these corals (inferred from  $\delta^{11}\text{B}$  analysis of the coral skeletons<sup>21</sup>) into the regression increases both the significance and coefficient of determination ( $p = 0.0028$ ,  $r^2 = 0.98$ , Table S3, Fig. 3). We cultured and analysed two duplicate colonies of the *P. murrayensis* genotype at 400 and 750  $\mu\text{atm}$  seawater  $\text{pCO}_2$  and observed large variations in skeletal amino acid concentrations between one pair of duplicates (G3, 750  $\mu\text{atm}$ , Fig. 1). We also observe large variations in  $\text{pH}_{\text{ECM}}$  between these corals<sup>21</sup> (Fig. 1) but we are able to explain almost all of the variation in coral calcification between them (and the other colonies) on the basis of skeletal amino acids and  $\text{pH}_{\text{ECM}}$  in combination.


The concentrations of some amino acids are highly correlated over the dataset e.g. for a linear regression of [Asx] and [Glx]  $r^2 = 0.90$ . This multicollinearity does not affect the coefficient of determination of the multiple linear regression model (i.e. describing how well the independent variables predict the dependent variable) but does increase the likelihood that we underestimate the significance of one or more amino acids in the regression model and that we misidentify the amino acid which may be responsible for changes in calcification rate.

It is unclear if higher skeletal [amino acid] reflects variations in the coral production of amino acid or enhanced skeletal incorporation of the produced amino acid, but we do not observe consistent relationships between the total skeletal amino acid produced each day (calculated from skeletal concentration and calcification rate<sup>20</sup> and seawater  $\text{pCO}_2$ , Figure S2a). We do not observe a significant correlation between the total amount of Asx (or other amino acids) produced each day and coral calcification rate (Figure S2b) to support the interpretation that calcification is limited by an energy budget required to synthesise the skeletal organic matrix<sup>21</sup> (assuming that amino acid concentrations are reflective of the total SOM).


Asparagine (and glutamine) undergo deamination during the amino acid extraction process and we cannot separately quantify aspartic acid and asparagine or glutamic acid and glutamine by this method. Proteomic methods, suggest that aspartic acid is a major component of skeletal proteins<sup>24</sup> and it is likely that the asparagine contribution to Asx is small. We observe inverse correlations between Asx and both glycine and leucine (Figure S3) which reflect changes in the composition of intracrystalline proteins but the reason for this is unclear.

**Interactions of aspartic acid and aragonite precipitation in vitro.** The coral skeleton analyses demonstrate that Asx (assumed to be predominantly composed of aspartic acid) is the most prevalent amino acid in the skeletons (Fig. 1) and is the amino acid most strongly correlated with coral calcification rate (Table S2). To explore the potential roles of aspartic acid in coral calcification we precipitated synthetic aragonites from seawater at the approximate pH<sup>21</sup> and soluble amino acid concentrations (see “Methods” section) inferred to occur at the coral calcification site. In our initial experiments (at seawater  $\text{pCO}_2 = 400 \mu\text{atm}$  and with pH and omega ( $\Omega$ ) co-varying) aspartic acid inhibited aragonite precipitation and inhibition was more pronounced at higher [aspartic acid] and at lower pH/ $\Omega$  (Fig. 4). We conducted a second series of experiments, under varying seawater  $\text{pCO}_2$  with either constant pH (varying  $\Omega$ ) or constant  $\Omega$  (varying pH) to separate the interactions of pH and  $\Omega$  with aspartic acid during aragonite precipitation (Fig. 5a, b). A multiple linear regression model of the entire dataset (Table S4) indicates that aragonite precipitation rates are significantly affected by both [aspartic acid] ( $p = 5.1 \times 10^{-20}$ , Fig. 5c) and  $\Omega$  ( $p = 3.2 \times 10^{-58}$ , Fig. 5a) but are independent of pH ( $p = 0.40$ , Fig. 5b). As far as we are aware, decoupling the omega and pH, and finding omega to be the principal driver of aspartic acid induced aragonite precipitation inhibition, is a unique observation.

The interaction of aspartic acid in aragonite precipitation is not fully understood. The precipitation of  $\text{CaCO}_3$  from a solution can occur via multiple stages, and organic additives have the ability to both promote and inhibit different stages<sup>15,16</sup>. In brief, homogenous nucleation of  $\text{CaCO}_3$  occurs in the absence of nucleation sites and likely proceeds via the formation of pre-nucleation clusters of the constituent ions of the solid (or other chemical species) which aggregate and dehydrate to form amorphous solids<sup>25</sup>. Biomolecules can act as templates for the aggregation of these amorphous nanoparticles which then develop into crystal domains after reaching a critical




**Figure 1.** Skeletal amino acid concentrations of each coral genotype (G1, G2 and G3) cultured at 25 °C and over a range of seawater pCO<sub>2</sub>. Two replicate colonies of G3 were cultured at 400 and 750 μatm. Error bars indicate the mean standard deviation of analyses of duplicate drilled samples. Calcification rates<sup>20</sup> (error bar shows typical standard deviation of 3–4 measurements per colony) and calcification site pH<sup>21</sup> (adjusted to the pH<sub>NBS</sub> scale, error bars show 95% confidence limits) are shown for reference. Asx aspartic acid + asparagine, Glx glutamic acid + glutamine.




**Figure 2.** Regressions between skeletal Asx concentration and (a) seawater pCO<sub>2</sub>, (b) seawater pH and (c) coral calcification rate<sup>20</sup>.


size<sup>26</sup>. Heterogeneous nucleation occurs in the presence of existing nucleation surfaces e.g. a mineral seed or an organic material, and requires the formation and subsequent growth of a nucleus on the existing surface. It is not known if ions or clusters of species are involved in these processes<sup>25,27</sup>. Much of the existing literature on biomolecule interactions during CaCO<sub>3</sub> precipitation focuses on calcite and vaterite rather than aragonite and there are contradictory reports of biomolecule effects. Aspartic acid accelerated the nucleation of vaterite in seeded<sup>28</sup> and unseeded<sup>16</sup> precipitations but stabilised pre-nucleation clusters and delayed CaCO<sub>3</sub> nucleation in another report<sup>15</sup>, consistent with computations indicating that aspartic acid reduces ion dissolution and aggregation in solution<sup>29</sup>. Interactions of aspartic acid at the solution:solid interface may promote calcite precipitation rate by decreasing the energy barrier to the attachment of solutes<sup>19</sup>, or may inhibit precipitation by blocking subsequent ion attachment<sup>30</sup>. Discrepancies between studies may reflect the impact of varying environmental factors, e.g. ionic strength<sup>31</sup>, pH (affecting molecule charge<sup>19</sup>), biomolecule concentration<sup>19</sup> and the presence of other cations e.g. Mg<sup>2+</sup><sup>32</sup>. Acidic proteins (rich in aspartic acid) can bind to and inhibit extension of specific



**Figure 3.** Observed and predicted coral calcification rates. Predicted rates were modelled using a multiple linear regression model of concentrations of Asx, Glx, serine, alanine and total amino acid and  $pH_{ECM}$ .



**Figure 4.** Aragonite precipitation rates from seawater in vitro as a function of [aspartic acid], pH and  $\Omega$  at seawater  $pCO_2 = 400 \mu atm$ . The inset shows the points at pH 8.34 on an expanded axis. Error bars indicate standard deviations of replicate precipitations ( $n = 2-10$ ) and are usually smaller than the symbols.



**Figure 5.** The % inhibition of aragonite precipitation (calculated by comparing mean precipitation with aspartic acid with the mean rate observed with no added aspartic acid) as a function of (a) pH at 2 mM aspartic acid and various  $\Omega$ , (b)  $\Omega$  at 2 mM aspartic acid and at various pH and (c) [aspartic acid] and  $\Omega$ . Error bars were calculated by compounding the standard deviations of precipitations with and without aspartic acid.

faces of calcite crystals thereby regulating growth and crystal morphology<sup>33</sup> and also bind  $\text{Ca}^{2+}$  and catalyse the precipitation of aragonite in vitro<sup>34</sup>.

In this study, aragonite precipitation from seawater is inhibited by aspartic acid at the concentrations inferred to occur in the coral ECM. Calcite propagation rates were also inhibited at similar [aspartic acid] in lower ionic strength solutions<sup>19</sup>. Inhibition of precipitation likely reflects aspartic acid adsorption at the crystal:solution interface which impedes the attachment of ions to further  $\text{CaCO}_3$  growth<sup>30</sup>. This inhibitory effect is less pronounced at higher  $\Omega$  (Fig. 5a). Aragonite precipitation rates were constant throughout each precipitation (Figure S4), indicating that the surface area for  $\text{CaCO}_3$  growth did not change. Variations in precipitation rate between experiments likely reflect crystal growth (rather than pre-nucleation or nucleation events). A potential explanation for our observations is that the ability of aspartic acid to adsorb to aragonite is reduced at high  $\Omega$ . Increased  $\text{CaCO}_3$  lattice disorder is inferred in synthetic aragonites precipitated at high seawater  $\Omega$  in the absence of biomolecules<sup>35</sup>. This may hamper the ability of aspartic acid to adsorb to the solid:solution interface (and inhibit subsequent precipitation). Alternatively, faster aragonite growth (at high  $\Omega$ ) could itself disadvantage the adsorption of aspartic acid to the growing crystal. It is unlikely that variations in  $\Omega$  in this study (driven by changes in  $[\text{CO}_3^{2-}]$ ) significantly affect aspartic acid speciation. Between pH 8 and 9, aspartic acid speciation in seawater is dominated by  $\text{Mg}(\text{asp})\text{H}^+$ ,  $\text{Mg}(\text{asp})^0$  and  $(\text{asp})\text{H}^-$  with Ca complexes playing a minor role ( $< 10\%$  of aspartic acid)<sup>36</sup>. We observe no significant impact of pH on aragonite precipitation by aspartic acid (Fig. 5b). This likely reflects the large difference between aspartic acid pKa ( $\sim 1.99$ ,  $3.90$  and  $9.90$  at  $25^\circ\text{C}$ ) and the pH range studied here ( $\text{pH}_{\text{NBS}} 8.34\text{--}8.69$ ).

We note that the free amino acid used in the precipitations here is not observed in high concentrations in extracts of coral SOM<sup>37</sup>, where aspartic acid is combined with other amino acids into peptides and proteins. Small aspartic acid-rich peptides inhibit calcite precipitation at lower concentrations than Asp-dipeptides or free aspartic acid<sup>19</sup> and adsorbing the amino acid to a template in small pore spaces may influence its effect on  $\text{CaCO}_3$  precipitation<sup>38</sup>. However, the interaction of aspartic acid with precipitating  $\text{CaCO}_3$  is likely due to the acidic carboxylic acid side chain which is present in both the free and protein-bound forms of the amino acid. We interpret our data to indicate an increase in aspartic acid (in whatever form) in the coral ECM under ocean acidification conditions and our in vitro precipitation experiments indicate that an increase in ECM [aspartic acid] could suppress coral calcification at high seawater  $\text{pCO}_2$ .

**Implications for coral response to ocean acidification.** Our experiments provide insights into the potential role of aspartic acid in coral biomineralisation under ocean acidification scenarios. Corals cultured at  $750\ \mu\text{atm}$   $\text{pCO}_2$  have higher skeletal [Asx] compared to their genotype analogues cultured at  $180\ \mu\text{atm}$  and  $400\ \mu\text{atm}$  (by 99% and 61% respectively). ECM  $\Omega$  in the branching coral *Stylophora pistillata* cultured at  $400\ \mu\text{atm}$  and  $25^\circ\text{C}$  is  $\sim 12$ <sup>39</sup>. At this  $\Omega$ , raising seawater [aspartic acid] from  $0.1$  to  $0.4\ \text{mM}$  (our current best estimate of the ECM concentration in corals cultured at  $180$  and  $750\ \mu\text{atm}$  respectively) increases the biomolecule inhibition of aragonite precipitation from  $\sim 3$  to  $\sim 11\%$  (Fig. 5c). In reality, the ECM  $\Omega$  is likely to be reduced at high seawater  $\text{pCO}_2$ , as evidenced by reductions in  $\text{pH}_{\text{ECM}}$ <sup>4,21</sup> and coral calcification rate<sup>2</sup>. This  $\Omega$  decrease will further exacerbate the inhibition of aragonite precipitation by aspartic acid (Fig. 5c). For the colonies analysed here, calcification at  $750\ \mu\text{atm}$  is reduced by 49% and 28% compared to the corals cultured at  $180\ \mu\text{atm}$  and  $400\ \mu\text{atm}$   $\text{pCO}_2$  respectively<sup>20</sup>. We conclude that the inferred change in ECM [aspartic acid] may be responsible for a substantial proportion of this calcification decrease.

Our study demonstrates that coral calcification is affected by both the concentration of organic molecules at the calcification site, and the DIC chemistry in the ECM<sup>21</sup>. Further research on the response and interactions of ECM organic materials and DIC chemistry to increases in seawater  $\text{pCO}_2$  and temperature is vital to understanding the response of coral calcification to future climate change.

## Methods

**Coral culturing experiments.** Massive *Porites* spp. corals were cultured at  $25^\circ\text{C}$  and at 3 different seawater  $\text{pCO}_2$  conditions ( $180\ \mu\text{atm}$ ,  $400\ \mu\text{atm}$  and  $750\ \mu\text{atm}$ <sup>20–22</sup>). Heads were imported from Fiji, identified to species level based on corallite morphology and were considered to represent different genotypes when they were collected from spatially separate (non-adjointing) colonies. Heads were sawn into multiple pieces (each  $\sim 12\ \text{cm}$  in diameter) so that at least one piece of each genotype could be cultured in each seawater  $\text{pCO}_2$  treatment. Corals were maintained at the different  $\text{pCO}_2$  treatments for 5 months, stained with Alizarin Red S to create a time horizon in the skeletons and then cultured for another 5 weeks before sacrifice. Corals were immersed in 3–4% sodium hypochlorite for 24–48 h with intermittent ultrasonic agitation to remove the tissues, then rinsed in distilled water. A slice was cut through the centre of each coral skeleton, parallel to the maximum growth axis using a water lubricated rock saw, cleaned in an ultrasonic bath and dried. Samples for amino acid analysis were obtained by drilling the skeleton deposited between the skeleton surface and the stain line i.e. representing the final 5 weeks of skeletal growth.

**Organic analysis.** This method follows the organics analysis in Tomiak et al.<sup>40</sup>. To isolate the intracrystalline fraction of amino acids  $< 20\ \text{mg}$  of skeletal sample was weighed in to a microcentrifuge tube and bleached using  $50\ \mu\text{L}$  12%  $\text{NaOCl}$  per mg, the fraction was below  $100\ \mu\text{m}$ . The sample was re-agitated over 48 h before the bleach was removed. The remaining sample was rinsed five times by centrifuging with  $18.2\ \text{M}\Omega\text{m}$  milli-Q  $200\ \mu\text{L}$  MeOH was added to the Eppendorf and pipetted off after several minutes. Samples were dried overnight. To hydrolyse samples,  $< 10\ \text{mg}$  was weighed in to a  $2\ \text{mL}$  sterile vial and  $20\ \mu\text{L}/\text{mg}$   $7\ \text{M}$   $\text{HCl}$  was added. Vials were flushed with nitrogen and placed in an oven at  $110^\circ\text{C}$ , vial caps were then retightened to prevent drying. After 24 h, the samples were removed and dried in a centrifugal evaporator. Samples were rehydrated with  $10\ \mu\text{L}/$



mg rehydration fluid (0.01 M HCl + 1.5 mM NaN<sub>3</sub>, spiked with L-homo-arginine). Samples were analysed using reverse-phase HPLC with fluorescence detection, following the method of Kaufman and Manley<sup>41</sup>. This enables quantification of L and D isomers of 12 amino acids. Asparagine and glutamine undergo deamination during the preparation process and may therefore contribute to observed concentrations of aspartic acid and glutamic acid. However other methods e.g. proteomics, suggest that skeletal proteins are dominated by aspartic acid<sup>24</sup> and we consider the asparagine contribution to be small. Blanks and standards were run throughout; all samples were run in duplicate.

The precision of amino acid characterisation was estimated by splitting drilled powders before amino acid extraction and by drilling skeletal samples from different sections of the same coral head (Table S1). To estimate the impact of inadvertent inclusion of the Alizarin Red stain in the drilled sample, we compared the amino acid compositions of a sample drilled along the stain line and a powder drilled along a plane parallel to the stain and immediately above it. The amino acid concentrations agree within replicate precision for total, and each of the amino acids with the exceptions of serine and phenylalanine, which both occur in low levels in the skeletons (Table S1). Inclusion of the stain did not affect the mol% of different amino acid groups.

**In-vitro aragonite precipitation experiments.** Synthetic aragonite was precipitated from seawater using a pH stat titrator (Metrohm Titrand 902) coupled with a gas system designed to produce air with a range of CO<sub>2</sub> concentrations<sup>20–22</sup>. We adjusted the pH and [DIC] chemistry of the seawater and added amino acids to explore the impacts of these variables on precipitation. Precipitations were conducted over a pH<sub>NBS</sub> of 8.34–8.69 (similar to that observed in the coral ECM, Fig. 1) and a [DIC]<sub>seawater</sub> of 3,000–8,000 μmol kg<sup>-1</sup>. The coral [DIC]<sub>ECM</sub> in *Stylophora pistillata* cultured at 400 μatm and 25 °C is ~3,000 μmol kg<sup>-139</sup>.

Precipitations were conducted in natural seawater (salinity 35) collected from Crail, Fife, filtered (0.45 μm cellulose nitrate filter) and stored in a blacked-out 1000 L high density polyethylene (HDPE) container, under air with a CO<sub>2</sub> atmosphere of ~400 μatm. For each precipitation, 320 mL seawater was decanted into a HDPE plastic beaker (total volume ~360 mL) capped with an ethylene tetrafluoroethylene lid with multiple ports and immersed in a water bath at 25 °C. A high precision combined pH/temperature sensor (Metrohm Aquatrod PT1000), a propeller stirrer, a gas tube and 2 titrant dosing tubes were inserted through the lid and into the headspace (gas tube) or seawater (all others). To avoid invasion or outgassing of CO<sub>2</sub>, the precipitating solution was maintained under a headspace with an air gas stream with pCO<sub>2</sub> set to a value at equilibrium with the solution. Prior to the addition of the seed the gas stream was set to ~400 μatm CO<sub>2</sub>. At the point of seed addition and thereafter the gas stream [CO<sub>2</sub>] was altered to that in equilibrium with the seawater (calculated using CO<sub>2</sub> sys v2.1 from seawater [DIC] and pH<sub>NBS</sub> using the equilibrium constants for carbonic acid<sup>42</sup>, for KHSO<sub>4</sub><sup>43</sup> and for [B]<sup>44</sup>). For each precipitation the amino acid (if used) was added first by suspending the amino acid in 1 mL of seawater and then pipetting this into the reaction vessel. The pH of the seawater was adjusted back to the starting value with the addition of 1.0 M NaOH. [DIC] was increased with the addition of 0.6 M Na<sub>2</sub>CO<sub>3</sub> and the pH adjusted to the required value by addition of 1.0 M HCl or NaOH. Once pH was stable 400 mg of an aragonite seed (from 2 batches of ground coral skeleton with surface areas of 3.45 and 4.41 m<sup>2</sup> g<sup>-1</sup> as analysed by the Brunauer–Emmett–Teller technique<sup>45</sup>) was added to the reaction vessel to act as a surface for mineral growth. Precipitation of CaCO<sub>3</sub> consumes DIC and Ca<sup>2+</sup> and reduces solution pH. The associated pH decrease triggered the titrator to add equal volumes of 0.45 M Na<sub>2</sub>CO<sub>3</sub> and 0.45 M CaCl<sub>2</sub>. The titration continued until 5 mLs of each titrant had been added resulting in the precipitation of ~225 mg of CaCO<sub>3</sub>. Seawater temperatures within and between precipitations varied by <0.2 °C. XRD and/or Raman spectroscopy of a least one precipitate produced under each set of conditions confirmed that the original seeds and precipitates were aragonite. Post-precipitation the sensor, beaker and propeller stirrer were submerged in 0.1 M HCl to dissolve any precipitate and then rinsed thoroughly with deionized water.

The pH sensor was calibrated with NIST buffers daily and are reported on the NBS scale. Buffers were replaced each week and the maximum difference observed between old and fresh buffers in a single session was 0.004 pH units. Seawater [DIC] was measured at the start and end of a subset of experiments using an Apollo Sci Tech AS-C3 DIC analyser to ensure that seawater [DIC] was as expected and to monitor for any CO<sub>2</sub> invasion or outgas during precipitations. Measured seawater [DIC] agreed with predicted seawater [DIC] within instrumental error<sup>46</sup> and [DIC] variations within single precipitations were typically 3% and always <15%. Seawater [Ca<sup>2+</sup>] was 10.1 mM as analysed by ICP-OES after 1:50 dilution with 3% HNO<sub>3</sub> and spiking with 1 ppm Y as the internal standard. Seawater [Ca<sup>2+</sup>] was measured using a Ca<sup>2+</sup> ion selective electrode (Metrohm) in filtered, acidified (pH 4) samples collected at the start and end of a subset of precipitations. Samples were maintained at constant temperature and measured consecutively to reduce electrode drift. Reproducibility of standards measured before and after samples was ≤5% and seawater [Ca<sup>2+</sup>] at the start and end of the precipitations always agreed within this value.

A sample titration is illustrated in Figure S4 indicating the tight control of pH during precipitation. The aragonite precipitation rate was estimated from the linear fit between time and titrant dosed and was normalized to the surface area of the starting seed. Control experiments, conducted under each set of precipitation conditions but without the addition of seed, demonstrated that no homogenous nucleation occurred in the solutions over the time scale of the precipitation i.e. no dosing of titrants occurred. Each set of precipitation conditions was tested multiple times (n = 2–10) with the exception of the precipitations at Ω = 35.4 which were not replicated. Data for individual precipitations are summarized in the Supplementary information (Table S5).

**Identification of amino acid concentrations for precipitation experiments.** We precipitated an estimated 225 mg of aragonite from a seawater solution with 2 mM aspartic acid onto 400 mg of inorganic aragonite seed. We estimate that the aragonite precipitated in vitro incorporated [aspartic acid] of 8,554 pmol mg<sup>-1</sup> (based

on RP-HPLC analysis of the final precipitate and starting seed respectively ([aspartic acid] = 7,700 pmol mg<sup>-1</sup> and 17 pmol mg<sup>-1</sup> respectively). We estimate an aspartic acid<sub>aragonite</sub>/aspartic acid<sub>seawater</sub> partition coefficient of ~0.004 (mmol L<sup>-1</sup>/mmol g<sup>-1</sup>) indicating that the incorporation of the amino acid in the aragonite has a minimal effect (<0.5%) on seawater [aspartic acid]. We estimate an [aspartic acid] of the coral calcification media of ~0.1–0.4 mM to yield the coral skeletal [aspartic acid] observed in the corals analysed here.

Received: 20 April 2020; Accepted: 17 June 2020

Published online: 30 July 2020

## References

- Hoegh-Guldberg, O. *et al.* Coral reefs under rapid climate change and ocean acidification. *Science* **318**, 1737–1742 (2007).
- Erez, J., Reynaud, S., Silverman, J., Schneider, K. & Allemand, D. Coral calcification under ocean acidification and global change. In *Coral Reefs: An Ecosystem in Transition* (eds Dubinsky, Z. & Stambler, N.) 151–176 (Springer, New York, 2011).
- Allemand, D., Tambutte, E., Zoccola, D. & Tambutte, S. Coral calcification, cells to reefs. In *Coral Reefs: An Ecosystem in Transition* (eds Dubinsky, Z. & Stambler, N.) 119–150 (Springer, New York, 2011).
- Venn, A. *et al.* Impact of seawater acidification on pH at the tissue-skeleton interface and calcification in reef corals. *Proc. Natl. Acad. Sci.* **110**, 1634–1639 (2012).
- Mass, T., Drake, J. L., Peters, E. C., Jiang, W. & Falkowski, P. G. Immunolocalization of skeletal matrix proteins in tissue and mineral of the coral *Stylophora pistillata*. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12728–12733 (2014).
- Falini, G., Fermani, S. & Goffredo, S. Coral biomineralization: a focus on intra-skeletal organic matrix and calcification. *Semin. Cell Dev. Biol.* **46**, 17–26 (2015).
- Helman, Y. *et al.* Extracellular matrix production and calcium carbonate precipitation by coral cells in vitro. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 54–58 (2008).
- Falini, G. *et al.* Control of aragonite deposition in colonial corals by intra-skeletal macromolecules. *J. Struct. Biol.* **183**, 226–238 (2013).
- Reggi, M. *et al.* Biomineralization in mediterranean corals: the role of the intraskeletal organic matrix. *Cryst. Growth Des.* **14**, 4310–4320 (2014).
- Reggi, M. *et al.* Influence of intra-skeletal coral lipids on calcium carbonate precipitation. *CrystEngComm* **18**, 8829–8833 (2016).
- Clode, P. L. & Marshall, A. T. Calcium associated with a fibrillar organic matrix in the scleractinian coral *Galaxea fascicularis*. *Protoplasma* **220**, 153–161 (2003).
- Puvel, S. *et al.* Soluble organic matrix of two Scleractinian corals: partial and comparative analysis. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* **141**, 480–487 (2005).
- Isa, Y. & Okazaki, M. Some observations on the Ca<sup>2+</sup>-binding phospholipid from scleractinian coral skeletons. *Comp. Biochem. Physiol. B* **87**, 507–512 (1987).
- Cuif, J. P., Dauphin, Y., Freiwald, A., Gautret, P. & Zibrowius, H. Biochemical markers of zooxanthellae symbiosis in soluble matrices of skeleton of 24 Scleractinia species. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **123**, 269–278 (1999).
- Picker, A., Kellermeier, M., Seto, J., Gebauer, D. & Cölfen, H. The multiple effects of amino acids on the early stages of calcium carbonate crystallization. *Zeitschrift für Krist.* **227**, 744–757 (2012).
- Tong, H. *et al.* Control over the crystal phase, shape, size and aggregation of calcium carbonate via a L-aspartic acid inducing process. *Biomaterials* **25**, 3923–3929 (2004).
- Tambutté, E. *et al.* Morphological plasticity of the coral skeleton under CO<sub>2</sub>-driven seawater acidification. *Nat. Commun.* **6**, 1–9 (2015).
- Coronado, I., Fine, M., Bosellini, F. R. & Stolarski, J. Impact of ocean acidification on crystallographic vital effect of the coral skeleton. *Nat. Commun.* **10**, 1–9 (2019).
- Elhadj, S., De Yoreo, J. J., Hoyer, J. R. & Dove, P. M. Role of molecular charge and hydrophilicity in regulating the kinetics of crystal growth. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 19237–19242 (2006).
- Cole, C., Finch, A. A., Hintz, C., Hintz, K. & Allison, N. Effects of seawater pCO<sub>2</sub> and temperature on calcification and productivity in the coral genus *Porites* spp.: an exploration of potential interaction mechanisms. *Coral Reefs* **37**, 471–481 (2018).
- Allison, N. *et al.* The effect of ocean acidification on tropical coral calcification: insights from calcification fluid DIC chemistry. *Chem. Geol.* **497**, 162–169 (2018).
- Cole, C., Finch, A., Hintz, C., Hintz, K. & Allison, N. Understanding cold bias: variable response of skeletal Sr/Ca to seawater pCO<sub>2</sub> in acclimated massive *Porites* corals. *Sci. Rep.* **6**, 1–8 (2016).
- Burton, E. A. & Walter, L. M. Relative precipitation rates of aragonite and Mg calcite from seawater: temperature or carbonate ion control. *Geology* **15**, 111–114 (1987).
- Ramos-Silva, P. *et al.* The skeletal proteome of the coral *Acropora millepora*: the evolution of calcification by co-option and domain shuffling. *Mol. Biol. Evol.* **30**, 2099–2112. <https://doi.org/10.1093/molbev/mst109> (2013).
- Gebauer, D., Kellermeier, M., Gale, J. D., Bergström, L. & Cölfen, H. Pre-nucleation clusters as solute precursors in crystallisation. *Chem. Soc. Rev.* **43**, 2348–2371 (2014).
- Pouget, E. *et al.* The initial stages of template-controlled CaCO<sub>3</sub> formation revealed by cryo-TEM. *Science* **323**, 1455–1458 (2009).
- Demichelis, R. *et al.* Simulation of crystallization of biominerals. *Annu. Rev. Mater. Res.* **48**, 327–352 (2018).
- Malkaj, P. & Dalas, E. Calcium carbonate crystallization in the presence of aspartic acid. *Cryst. Growth Des.* **4**, 721–723 (2004).
- Finney, A. R. & Rodger, P. M. Probing the structure and stability of calcium carbonate pre-nucleation clusters. *Faraday Discuss.* **159**, 47–60 (2012).
- Sikirić, M. D. & Füredi-Milhofer, H. The influence of surface active molecules on the crystallization of biominerals in solution. *Adv. Colloid Interface Sci.* **128**, 135–158 (2006).
- Tai, C. Y. & Chen, F. B. Polymorphism of CaCO<sub>3</sub> precipitated in a constant-composition environment. *AIChE J.* **44**, 1790–1798 (1998).
- Sand, K. K., Pedersen, C. S., Matthiesen, J., Dobberschütz, S. & Stipp, S. L. S. Controlling biomineralisation with cations. *Nanoscale* **9**, 12925–12933 (2017).
- Addadi, L. & Weiner, S. Interactions between acidic proteins and crystals: stereochemical requirements in biomineralization. *Proc. Natl. Acad. Sci.* **82**, 4110–4114 (1985).
- Mass, T. *et al.* Cloning and characterization of four novel coral acid-rich proteins that precipitate carbonates in vitro. *Curr. Biol.* **23**, 1126–1131 (2013).
- DeCarlo, T. M. *et al.* Coral calcifying fluid aragonite saturation states derived from Raman spectroscopy. *Biogeosciences* **14**, 5253–5269 (2017).
- De Stefano, C., Foti, C., Gianguzza, A., Rigano, C. & Sammartano, S. Chemical speciation of amino acids in electrolyte solutions containing major components of natural fluids. *Chem. Speciat. Bioavailab.* **7**, 1–8 (1995).
- Puvel, S. *et al.* Evidence of low molecular weight components in the organic matrix of the reef building coral, *Stylophora pistillata*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **147**, 850–856 (2007).

38. Cantaert, B., Beniash, E. & Meldrum, F. C. The role of poly(aspartic acid) in the precipitation of calcium phosphate in confinement. *J. Mater. Chem. B* **1**, 6586–6595 (2013).
39. Sevilgen, D. S. *et al.* Full in vivo characterization of carbonate chemistry at the site of calcification in corals. *Sci. Adv.* **5**, eaau7447 (2019).
40. Tomiak, P. J. *et al.* Testing the limitations of artificial protein degradation kinetics using known-age massive *Porites* coral skeletons. *Quat. Geochronol.* **16**, 87–109 (2013).
41. Kaufman, D. S. & Manley, W. F. A new procedure for determining DL amino acid ratios in fossils using reverse phase liquid chromatography. *Quat. Sci. Rev.* **17**, 987–1000 (1998).
42. Lueker, T. J., Dickson, A. G. & Keeling, C. D. Ocean pCO<sub>2</sub> calculated from dissolved inorganic carbon, alkalinity, and equations for K<sub>1</sub> and K<sub>2</sub>: validation based on laboratory measurements of CO<sub>2</sub> in gas and seawater at equilibrium. *Mar. Chem.* **70**, 105–119 (2000).
43. Dickson, A. Standard potential of the reaction: AgCl (s) + 12H<sub>2</sub> (g) = Ag (s) + HCl (aq) and the standard acidity constant of the ion HSO<sub>4</sub><sup>−</sup> in synthetic sea water from 273.15 to 318.15 K. *J. Chem. Thermodyn.* **22**, 113–127 (1990).
44. Uppström, L. The boron/chlorinity ratio of deep-sea water from the Pacific Ocean. *Deep Sea Res. Oceanogr. Abstr.* **21**, 161–162 (1974).
45. Brunauer, S., Deming, L. S., Deming, W. E. & Teller, E. On a theory of the van der Waals adsorption of gases. *J. Am. Chem. Soc.* **62**, 1723–1732 (1940).
46. Evans, D., Webb, P. B., Penkman, K., Kröger, R. & Allison, N. The characteristics and biological relevance of inorganic amorphous calcium carbonate (ACC) precipitated from seawater. *Cryst. Growth Des.* **19**, 4300–4313 (2019).

## Acknowledgements

This work was supported by the Leverhulme Trust (Research project Grant 2015-268 to NA, RK, and KP) and the UK Natural Environment Research Council (NE/G015791/1 to NA and AF).

## Author contributions

N.A., K.P., R.K. and A.F. designed the study. N.A., A.F., K.H. and C.H. built the culturing system, C.C. and N.A. cultured the corals, C.K. and N.A. conducted the precipitation experiments. D.E., C.K. and K.P. prepared and analysed the samples for amino acid analysis. C.K., N.A. and K.P. analysed the data and wrote the manuscript. All authors contributed to the analysis of the results and to the writing of the paper.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary information** is available for this paper at <https://doi.org/10.1038/s41598-020-69556-0>.

**Correspondence** and requests for materials should be addressed to N.A.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020