



A randomised trial to assess fluid and electrolyte balance responses following ingestion of different beverages in young and older men

Nidia Rodriguez-Sanchez¹ · Stuart D. R. Galloway¹

Received: 27 February 2023 / Accepted: 23 May 2023
© The Author(s) 2023

Abstract

Background Older adults are susceptible to dehydration and fluid overload due to a reduced ability to maintain homeostatic control of fluid and electrolyte balance.

Purpose To assess fluid and electrolyte balance responses in young and older men following ingestion of commonly consumed beverages differing in composition.

Methods 12 young and 11 older men were recruited. Euhydrated body mass was recorded. Participants consumed 1L (250 ml every 15 min) of water, fruit juice, a sports drink or low-fat milk in a randomized cross-over design. Urine and blood samples were obtained before and after the drinking period and every hour thereafter for 3-h. Samples were used to determine osmolality, electrolytes (Na⁺ and K⁺), water clearance, and glomerular filtration rate.

Results Free water clearance was significantly higher in Young than Older at 1 and 2 h after the ingestion of W and S ($p < 0.05$). Net Na⁺ and K⁺ balance were not different between Young and Older ($p = 0.91$ and $p = 0.65$) adults, respectively. At 3 h Na⁺ balance was negative after ingesting water and fruit juice, but neutral after sport drink and milk. Net K⁺ balance was neutral at 3 h after ingesting milk, but negative after water, fruit juice and sport drink.

Conclusions Milk was retained longer than other beverages in Young, but not in Older, despite similar net electrolyte balance responses. Older had higher fluid retention in the first 2 h after the ingestion of all beverages, except for milk when compared to Young, indicating an age-related loss of ability to regulate fluid balance under current study conditions.

Keywords Hydration · Dehydration · Macronutrients · Renal excretion · Urine · Creatinine

Abbreviations

ANOVA Analysis of variance
GFR Glomerular filtration rate
NFB Net fluid balance

Introduction

Ageing is accompanied with changes in the homeostatic control systems that regulate fluid and electrolyte balance (Rolls and Philips 1990) increasing the risk of older adults becoming dehydrated, but also predisposing them

to developing problems of fluid overload (Allison and Lobo 2004). Inadequate hydration status has been linked to urinary tract infections, impaired cognitive function, and delirium, and is related to a higher incidence of falls in older adults (Rowe and Juthani-Mehta 2013; Dubeau 2006). Whereas an impaired ability to excrete excess fluid promptly due to an age-related reduction in glomerular filtration rate [GFR (Dontas et al. 1972)] can lead to passive reabsorption of fluid with a higher risk of presenting water overload and hyponatraemia (Musso and Oreopoulos 2011). The consequences of overhydration can be life-threatening (El-Sharkawy et al. 2014). Of course, any impact of ageing on human health or function is complicated by the interaction of various factors including lifestyle choices, such as physical activity behaviours. For example, research on musculoskeletal function is often complicated by the interaction between ageing and sedentary behaviour (Moreno-Agostino et al. 2020; Paterson and Warburton 2010).

However, under free-living conditions, older adults are more prone to become dehydrated than their younger

Communicated by Michalis G Nikolaidis.

✉ Nidia Rodriguez-Sanchez
nidia.rodriguezsanchez@stir.ac.uk

¹ Physiology, Exercise and Nutrition Research Group,
Faculty of Health Sciences and Sport, University of Stirling,
Stirling FK9 4LA, UK

counterparts. Fluid deficits in older adults can occur through a variety of routes such as: a blunted thirst response, resulting from a decrease in the sensitivity of volume and osmoreceptors (Kenney and Chiu 2001); loss of lean mass leading to a reduction in total body water content by 10–15% (Allison and Lobo 2004, Adams et al. 2014); and/or reduced capacity of kidneys to concentrate urine (Sands 2012). Older adults also have a reduced capacity to handle sodium loads, making them more prone to over-expansion of the extracellular fluid compartment (Luckey and Parsa 2003). Ageing is associated with progressive tubular dysfunction, resulting in decreased sodium reabsorption and reduced potassium secretion (Musso and Oreopoulos 2011). These differences in electrolyte handling suggest that older adults cannot deal with sodium or potassium loads as efficiently as younger adults, making them prone to electrolyte abnormalities such as dysnatremias as older adults can experience a decrease in the ability to excrete or reabsorb sodium (Musso et al. 2006, Schlanger et al. 2010). Older adults have a decreased transtubular potassium gradient when compared with their young population (Musso and Jauregui 2014, Lindeman et al. 1985). Ageing confers a natural decline in several aspects of gastrointestinal physiology, including gastrointestinal peristalsis, enteric nervous system, intestinal mucosa, mucosal immunity, and gut absorption (Atillasoy and Holt 1993, Hooper et al. 2014; Russell 1992). The risk of dehydration in older adults is exacerbated by cognitive and behavioural changes such as a lack of knowledge, or misconceptions concerning the effects of drinking, or not drinking, sufficient fluid (Hooper et al. 2014, Bhanu et al. 2020). Furthermore, many older adults limit or avoid beverages to reduce the frequency of urination (Godfrey et al. 2012). All older adults are at risk of low-intake dehydration, and they should be encouraged to ingest an adequate amount of fluids (Volkert et al. 2019). Different beverages can contribute to the daily fluid intake. However, little is known about the renal handling of popular and commercially available beverages with different composition in the older adult population.

Data on young adults highlights that electrolyte composition, as well as certain macronutrients (i.e., protein/carbohydrate), are important in determining the fluid retention response to beverage ingestion (Maughan et al. 2016, 2019). However, to our knowledge, only a pair of companion papers from the same sample of older adults have investigated fluid and electrolyte balance responses to beverages with different macronutrient and electrolyte composition (Clarke 2019, Wolf et al. 2019). In their sample, an amino acid-based beverage (containing 5 amino acids, 30 mmol Na⁺ and 10 mmol K⁺) had the highest hydration potential compared to water in the older group at 2 h after ingestion. Therefore, we aimed to further investigate the fluid and electrolyte balance responses to a range of frequently consumed commercially available beverages differing in macronutrient

and electrolyte content, and to compare responses between young and older adult males. We hypothesized that beverages with a higher macronutrient/electrolyte content would lead to greater fluid retention than water in young adults, but that these responses would be blunted in an older adult group.

Methods

The fluid and electrolyte balance responses after ingestion of four commonly consumed and commercially available beverages were tested in young and older volunteers. The beverages were: still water (as control); fruit juice with a moderate carbohydrate and potassium content; a sport drink with moderate carbohydrate and sodium content (S); and a low fat 1% milk with moderate carbohydrate, protein, and sodium and potassium content. The test beverages were specifically chosen with an energy content in a range of 230–350 kcal/L (except for water) and provided either sodium, potassium, or a mix of electrolyte content (Table 2). The study was approved by the University of Stirling—School of Sport Research Ethics Committee (SSREC #753). All participants provided their informed consent for their participation.

Pre-trial standardization/exclusion criteria

Twenty-four healthy active volunteers were recruited into two groups: $n = 12$ were allocated to the young group (18–35y) and $n = 12$ to the older group ($> 55y$) with participants matched for stature and body mass. One participant in the older group dropped out (Table 1). Those participants with

Table 1 Participant anthropometric characteristics, hydration habits, caffeine and alcohol intake, and self-reported physical activity

	Age group	
	Young ($n = 12$)	Old ($n = 11$)
Age (years)	24.5 (4.3)	63.8 (5.5)*
Body mass (kg)	76.7 (9.1)	78.2 (9.1)
Stature (m)	1.79 (0.07)	1.77 (0.06)
Body mass index (kg/m ²)	24.0 (2.3)	25.1 (3.6)
Self-reported fluid intake (ml/d)	1929 (310)	1780 (215)
Caffeine intake (mg/d)	211.3 (112.7)	230.5 (79.9)
Weekly alcohol intake (units)	4.6 (5.7)	7.5 (3.1)
Moderate to vigorous physical activity (hours per week)	5.9 (1.0)	6.7 (2.6)
Participants reporting to exercise 4 or more times per week (n)	9	8

Values are mean (SD)

*Indicates significant difference between young and older groups. ($p < 0.05$)

Table 2 Water, energy, macronutrient, sodium, and potassium content of tested beverages

Beverage	Water content (%)	Energy (kcal/100 ml)	CHO (g/100 ml)	Fat (g/100 ml)	Protein (g/100 ml)	Sodium (mmol)	Potassium (mmol)
Water-control	100	0	0.0	0.0	0.0	0	0
Fruit juice	95	23	4.4	0.0	0.3	0	28
Sport drink	94	28	6.4	0.0	0.0	22	3
Low fat 1% milk	91	35	5.0	0.1	3.4	19	23

Apart from water, test beverages were selected to provide similar carbohydrate (CHO) content, but with differing protein and electrolyte composition

a history of cardiovascular, renal, musculoskeletal, or metabolic disease determined from a health screen questionnaire were excluded. Participants following an energy restricted diet and/or exercise plan to lose or gain mass were excluded. Volunteers were asked to record and replicate their food and fluid intake two days before each experimental trial, and to refrain from alcohol ingestion and vigorous physical activity for 24 h before all trials.

Participant characteristics on entry into the study

The physical activity levels of the participants varied, with the older group reporting an average of 6.7 h of moderate/intense exercise per week, compared to the young group's reported average of 5.9 h per week. Participants engaged in a range of sports or activities, with jogging more than 5 k, swimming more than 600 m, and gym combining strength and aerobic exercises being the most frequently reported in the young group. In contrast, the older group reported cycling, gym combining strength and aerobic exercises, and hill walking as their most frequently reported activities, with some participants also reporting engaging in less common activities such as ice skating and skydiving (Table 1).

Experimental procedure

Participants attended the laboratory for four experimental trial days, each separated by 7-days. All trials were conducted in the morning after an overnight fast (> 8 h). Participants ingested 500 ml of still water (Highland Spring®) 1 h before attending the laboratory to ensure euhydration upon arrival. Euhydration was required to ensure that all participants could provide urine samples over the entire study monitoring period. When the participants arrived at the laboratory, they were asked to empty their bladder/bowel, urine was collected, urine mass was recorded, and a 5 ml aliquot retained for analysis. An intravenous cannula was inserted into an antecubital vein, and initial near nude body mass was recorded (in underwear only). Participants adopted a seated position for 15 min after which a baseline blood sample was drawn. Participants then consumed a fixed volume (1L, divided in four aliquots of 250 ml provided every

15 min) of water (Highland Spring®), fruit juice (Tropicana, Trop50®), sport drink (Lucozade Sport®) or milk (low fat 1% skimmed milk) (Table 2). Test beverages were administered in a randomised counterbalanced order. Standard commercial beverages were purchased as a single batch or from a single source (for products with a short shelf life) to be used for all trials. All beverages were stored at a standard refrigerated temperature (4–6 °C) until serving.

Immediately at the end of the 60-min beverage ingestion period, participants were asked to empty their bladder, and subsequently each hour for the next 3 h to monitor urine volume. Participants could urinate at any time they required but they were also asked to empty their bladder completely every 60-min to track hourly urine production. If volunteers needed to urinate before the 60-min period, the urine was collected and added to the urine produced at the next 60-min time point. Urine mass was recorded to determine cumulative urine mass every 60-min and a 5 ml aliquot was retained for analysis. Blood samples were drawn immediately after the drinking period, and every 60-min for the next 3 h. All blood samples were drawn from participants after they had remained in a seated position for at least 15-min (Supplementary Appendix 1).

Blood, serum, and urine analysis

Total urine mass (to nearest g) was measured over the 3 h post beverage ingestion. Samples of urine obtained each hour were analysed for urine osmolality, creatinine concentration and urine electrolyte content. From each urine void, a 5 ml aliquot was collected and stored at 4 °C for the analysis of urine osmolality, electrolyte content (sodium and potassium) and creatinine determination. Whole blood samples were dispensed into a serum tube for osmolality and electrolyte determinations, and an EDTA tube with plasma stored for later analysis of creatinine. Urine and serum osmolality were measured in duplicate using a freezing point depression method (Löser osmometer) on day of collection. Sodium and potassium concentration in urine and serum were measured in duplicate using flame-photometry (PFP7/C Clinical Flame Photometer, Jenway) within 5-days of collection.

Macronutrient data of test beverages were obtained from product nutrition information. Beverage electrolyte data were obtained by flame-photometry analysis.

Data calculations and statistical analysis

All the participants achieved a positive net fluid balance of 1000(g) following the fixed volume of the fluid ingested (1L). Net fluid balance each hour was then calculated from the total mass of urine (cumulative urine output) that had been collected to that point. Electrolyte balance on each trial was determined by subtracting the amount of sodium or potassium excreted through urine from the total sodium or potassium ingested within the test beverage.

Urine creatinine was used to estimate GFR (Levey et al. 2009, Devanand and Chithrapavai 2013). Free water clearance was calculated at 1 h, 2 h and 3 h after test beverage ingestion following the equations described by Wolf et al. (Wolf et al. 2019).

Cumulative urine outputs, by beverage and between groups, were compared using repeated measures ANOVA. Participant characteristics were analysed by paired t-tests. Data for net fluid balance, and electrolytes were analysed by 3-way ANOVA (beverage x time point x age group) to identify within and between group differences. All statistical analyses were completed using a statistical software package (IBM SPSS Statistics, V.23). Statistical significance was accepted at $P < 0.05$. Data are presented as mean (95% CI) or mean (SD) in the text, and as mean (SD) in Tables and Figures.

Results

Of the twenty-four participants recruited (12 young and 12 older men), $n = 1$ older adult participant dropped out of the study due to aversion with the blood sampling procedures. No participants experienced any adverse effect or gastrointestinal symptoms following beverage ingestion.

Mean urine and serum osmolality when participants arrived at the lab, prior to beverage ingestion, was not different between young and older adults, except for urine osmolality on trial M (Table 2). Mean serum and urine osmolality demonstrated that both groups began all trials in a euhydrated state.

Cumulative urine output and urine osmolality responses to beverage ingestion

Urine mass did not differ between trials immediately after the beverage ingestion period for Young and Older. However, 1 h after the ingestion of water, the cumulative urine output for Young was significantly higher than Older. This

was sustained at 2 h post drinking ($p < 0.05$). Urine osmolality was significantly lower in Young compared to Older at 1 h after ingestion of water; and significantly higher in Young than Older at 2 h after the ingestion of sport drink and milk, and at 3 h for water, fruit juice and sport drink (Table 2).

Net fluid balance (NFB)

At 1 and 2 h after ingesting water, there were significant differences between Young and Older. However, no significant difference was observed between age groups when comparing NFB at 3 h after ingestion of any of the different beverages. NFB after ingestion of milk was significantly different from water in Young, but not in Older (Fig. 1).

Estimated GFR and free water clearance

Mean (SD) estimated GFR was lower in older participants (74.1 mL/min/1.73 m²; 70.8, 77.5) when compared with the younger group (98.9 mL/min/1.73 m²; 94.8, 102.9) ($p < 0.001$). Estimated free water clearance rate was significantly higher in Young compared with Older, particularly at 1 h and 2 h ($p < 0.05$) after the ingestion of water and sport drink (Fig. 2). No other differences were observed at other time points between groups. When the test beverages were compared with water within the same age group, for young adults, milk was different from water at 1 h, 2 h, and 3 h post ingestion ($p < 0.05$), but this was not observed in the older adult group.

Electrolyte balance

Net sodium balance 3 h after test beverage ingestion was greater in both age groups when they ingested sport drink, leading to a more positive sodium balance in comparison with water. No other significant differences in sodium balance were observed between test beverages or between groups (Fig. 3). Fruit juice led to a more positive potassium balance 2 h after its ingestion in both groups when compared with water ($p < 0.05$) (Fig. 4). There was no significant difference between age groups in sodium or potassium concentration in serum at any time point. However, in the young group only the sodium concentration in serum was significantly higher when participants drank sport drink than when they ingested water, at 2 h and 3 h after ingestion. In the older group, there was a significant difference in serum potassium at 1 h, 2 h, and 3 h after ingesting milk in comparison with water (Table 3).

Fig. 1 Net fluid balance following the ingestion 1L of water (A), fruit juice (B), sports drink (C), and milk (D) in young and in older men. Values are mean (SD). # $p < 0.05$ between age groups





Fig. 2 Free water clearance responses after the ingestion of 1 L of water (A), fruit juice (B), sport drink (C) and milk (D) at 1, 2 and 3 h after ingestion of test beverage in young and older men. Values are mean (SD). # $p < 0.05$ between age groups



Discussion

We investigated fluid and electrolyte balance in young and older adults after the ingestion of beverages with different nutrient composition and assessed their retention over three hours post ingestion. In young adults, we replicated

our previous observations (Maughan et al. 2016, 2019) by demonstrating that milk led to greater net fluid balance than water over the follow-up period. However, this effect of milk was not observed in older adults. Our findings suggest that when older adults ingest beverages differing in electrolyte and macronutrient content the resulting fluid delivery, absorption and electrolyte balance appears

Fig. 3 Net sodium balance after the ingestion of 1 L of water (A), fruit juice (B), sport drink (C) and milk (D) at 1, 2 and 3 h after ingestion of test beverage in young and older subjects. Values are mean (SD)





Fig. 4 Net potassium balance after the ingestion of 1 L of water (A), fruit juice (B), sport drink (C) and milk (D) at 1, 2 and 3 h after ingestion of test beverage in young and older subjects. Values are mean (SD)



insufficient to impact upon overall net fluid balance, 3 h after ingestion. In the first 2 h after water ingestion, a lower free water clearance rate and more positive net fluid balance was observed in older adults. No differences in the electrolyte balance between age groups were observed. The different responses between older and young male adults are likely due to age-related changes in gastric emptying and absorption of fluid, declines in renal function.

Gastric emptying and intestinal absorption of fluid

In the present study, the difference in fluid balance response between young and older adult groups may be influenced mostly by gastric emptying rates. In older adults an age-related reduction in gastric acid secretion would potentially allow for a greater gastric emptying rate of milk by reducing casein clot formation (Huppertz and Chi 2021). It has been

Table 3 Cumulative urine mass, urine osmolality, serum osmolality and electrolytes (Na⁺ and K⁺) in serum prior and following the ingestion of the test beverages in young and older adult groups

	Age group	Test beverage	Time after ingestion of test beverage (min) (post ingestion time point)				
			– 60	0	60 (1 h)	120 (2 h)	180 (3 h)
Cumulative urine mass	Young	Water	–	–	825 (158) ⁺	1082(261) ⁺	1177 (288)
		Fruit Juice			773 (279)	1009 (303)	1082 (316)
		Sport drink			804 (243)	1034 (289)	1120 (311)
		Low fat 1% milk			635 (196)	822 (195)	956 (339)
	Old	Water	–	–	544 (285) ⁺	791 (342) ⁺	987 (331)
		Fruit Juice			599 (196)	878 (223)	1037 (227)
		Sport drink			610 (204)	861 (190)	994 (199)
		Low fat 1% milk			625 (196)	851 (184)	978 (174)
Urine osmolality (mOsmol/l)	Young	Water	545 (323)	276 (188)	86 (21) ⁺	194 (103)	403 (149) ⁺
		Fruit Juice	491 (305)	298 (260)	111 (34)	229 (94)	533 (165) ⁺
		Sport drink	523 (316)	315 (237)	106 (52)	255 (139) ⁺	519 (221) ⁺
		Low fat 1% milk	578 (290) ⁺	358 (267)	249 (159)	381 (104) ⁺	566 (187)
	Old	Water	387 (122)	221 (88)	146 (54) ⁺	161 (51)	222 (63) ⁺
		Fruit Juice	352 (111)	250 (109)	139 (41)	171 (42)	251 (85) ⁺
		Sport drink	434 (210)	310 (183)	111 (27)	158 (52) ⁺	287 (123) ⁺
		Low fat 1% milk	385 (139) ⁺	256 (151)	174 (64)	268 (106) ⁺	411 (160)
Serum osmolality (mOsmol/l)	Young	Water	295 (4)	295 (4)	294 (3)	295 (4)	295 (3)
		Fruit Juice	296 (3)	297 (3)	295 (3)	296 (3)	295 (3)
		Sport drink	296 (3)	297 (3)	296 (3)	295 (3)	295 (3)
		Low fat 1% milk	297 (3)	297 (2)	297 (3)	295 (2)	296 (2)
	Old	Water	296 (4)	294 (4)	293 (3)	294 (4)	295 (4)
		Fruit Juice	295 (3)	296 (4) *	294 (4)	294 (4)	295 (4)
		Sport drink	295 (3)	295 (3)	294 (3)	293 (3)	294 (3)
		Low fat 1% milk	294 (3)	296 (3)*	296 (3) *	295 (3)	295 (4)
Serum sodium (mmol)	Young	Water	142 (6)	140 (5)	141 (5)	139 (6)	139 (7)
		Fruit Juice	141 (5)	139 (4)	143 (4)	143 (5)	144 (5)
		Sport drink	144 (5)	142 (6)	143 (6)	147 (7)*	146 (6)*
		Low fat 1% milk	141 (7)	143 (6)	143 (6)	143 (6)	142 (5)
	Old	Water	142 (7)	139 (6)	139 (6)	139 (6)	142 (6)
		Fruit Juice	142 (6)	142 (6)	142 (7)	144 (5)	142 (5)
		Sport drink	141 (5)	143 (5)	143 (7)	144 (7)	142 (6)
		Low fat 1% milk	140 (6)	142 (8)	141 (8)	139 (6)	140 (7)
Serum potassium (mmol)	Young	Water	4.1 (0.7)	4.3 (0.6)	4.3 (0.6)	4.3 (0.5)	4.4 (0.6)
		Fruit Juice	4.1 (0.5)	4.2 (0.5)	4.4 (0.5)	4.4 (0.3)	4.5 (0.3)
		Sport drink	4.2 (0.5)	4.3 (0.6)	4.3 (0.7)	4.6 (0.8)	4.4 (0.5)
		Low fat 1% milk	4.1 (0.5)	4.3 (0.7)	4.5 (0.7)	4.6 (0.8)	4.4 (0.5)
	Old	Water	4.4 (0.5)	4.6 (0.5)	4.6 (0.5)	4.7 (0.5)	4.7 (0.5)
		Fruit Juice	4.6 (0.6)	4.7 (0.6)	4.7 (0.6)	4.8 (0.5)	4.8 (0.5)
		Sport drink	4.4 (0.4)	4.4 (0.5)	4.5 (0.6)	4.5 (0.5)	4.2 (0.5)
		Low fat 1% milk	4.5 (0.3)	4.9 (0.5)	5.0 (0.5)*	5.0 (0.4)*	5.0 (0.3)*

Values are mean (SD)

*Indicates significant difference from water within group and column

⁺Indicates significant difference between young and older groups. (p < 0.05)

demonstrated previously that basal and maximal gastric acid output both decrease in ageing humans (Holt et al. 1989). Dangin et al. (2003) observed that in young people casein

could be classified as a slow protein due to reduced gastric emptying rate and subsequent amino acid delivery, when compared against whey protein. However, in older adults

there is a faster casein-emptying rate due to an age-related decline in gastric acid secretion.

Most of the water (ingested and emptied into the gastrointestinal tract) is reabsorbed in the small intestine, and remaining water in the colon. The small intestine is highly permeable promoting a rapid balance of the digestive content through the absorption of electrolytes and nutrients (Leiper 1998). Water can pass through the intestinal epithelium via the paracellular or the transcellular route. The transcellular route consists of different mechanisms: passive diffusion, cotransport and the aquaporins (Laforenza 2012). To our knowledge, there are no studies documenting modifications of any of these mechanisms that could impair water bioavailability in older adults. However, the mucosal epithelium is also important for water and electrolyte absorption, and ageing impairs mucous secretion (Branca et al. 2019). Thus, it is not known if age-related changes in gastrointestinal physiology negatively impact upon water and electrolyte absorption (Luchette and Yelon 2017).

Renal function

The ability to concentrate urine declines with age. In the Baltimore Longitudinal Study of Ageing, individuals aged 60–79 years had a ~20% reduction in maximal urine osmolality, a 50% decrease in the ability to reabsorb sodium and urea, and reduced capacity to concentrate solutes, when compared with 20–39 years old group (Lindeman et al. 1985, Rowe et al. 1976). In the present study, it was observed that there was a 39% lower urine osmolality in the older group 3 h after beverage ingestion, regardless of beverage composition ingested, when compared with the younger group. This finding reinforces the decline in urinary concentrating capacity with ageing (Musso et al. 2012).

In the present study, the volume overload induced by ingesting 1 L of fluid in 1 h, when in an already euhydrated state, leads to a more prolonged fluid retention of water in older adults, likely due to a reduced homeostatic control response compared to young adults (Beck 2000). This is evidenced by lower total urine output and calculated free water clearance in the first 2 h after ingestion of water in older than in younger adults. Further, it is known that GFR is preserved until about the age of 40 years, after that it declines linearly at an average rate of about 8 mL/min/decade (Silva 2005). In the present study, we observed that mean estimated GFR was 25.7 mL/min lower in older than in younger men, demonstrating a modest reduction in estimated GFR in the older group (6.5 mL/min per decade). Thus, it seems that differences between young and older adults, particularly in the initial response to water ingestion, relate to age related declines in homeostatic control and renal function that result in delayed excretion of the fluid overload.

Electrolyte balance

Older people are prone to expansion of total body water when challenged with a volume or sodium overload. Older adults have been reported to have a diminished capacity for renal sodium excretion (Luckey and Parsa 2003, Luft et al. 1979, Stachenfeld et al. 1996). However, as a rapid increase in GFR is required to deal with acute salt overload it is natural to consider that older people (with a normally reduced GFR) will present a limited ability to manage sodium loads (Hall 2016). A reduced capacity to handle sodium loads would predispose older adults to overexpansion of the extracellular fluid compartment. In the present study, the commercially available beverages that were provided were in the range of 0–22 mmol/L sodium. This range is likely insufficient to challenge homeostatic controls and observe any significant difference between groups or between beverages. Indeed, no differences were noted in net sodium balance response to the sodium containing sport drink between young and older adults. The mean loss of sodium in urine appeared to be less in older than younger adults after milk ingestion; however, this did not reach statistical significance. Future research is needed to examine electrolyte balance responses in older adults following larger sodium loads. Potassium excretion is reported to be reduced in the aged kidney and is attributed to the combination of low potassium secretion (Musso et al. 2006) and high potassium reabsorption (Eaton and Pooler 2018). As a result, healthy older people have a reduction in the transtubular potassium gradient compared with younger adults, leading to reduced excretion of potassium load in older adults (Musso et al. 2006). In the present study no differences were observed between age groups suggesting no changes in handling of potassium across the age range and potassium loads examined.

Conclusion

Further investigation is necessary to examine older adults under a fluid deficit condition to establish how older adults deal with fluid replacement and fluid overload situations. In addition, future research could aim to investigate more significant challenges to homeostasis using test drinks with greater electrolyte and macronutrient composition differences. However, under the conditions of the present study, it is evident that free water clearance is impaired in the first two hours following a fluid overload in older adults when compared with responses in young.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00421-023-05241-0>.

Author contributions The authors' responsibilities were as follows: NR-S and SDRG conceived the project, NR-S and SDRG developed the overall research plan, NR-S conducted the research and analysed the samples, NRS performed the statistical analysis, NR-S and SDRG wrote the manuscript, SDRG had primary responsibility for the final content; and both authors read and approved the final manuscript. The authors would like to thank Mr Chris Grigson for his technical support in the samples analysis and all the volunteers for taking part in the study.

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Conflict of interest Nidia Rodriguez-Sanchez, no conflicts of interest to declare. Stuart DR Galloway, no conflicts of interest to declare.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adams KF, Leitzmann MF, Ballard-Barbash R, Albanes D, Harris TB, Hollenbeck A et al (2014) Body mass and weight change in adults in relation to mortality risk. *Am J Epidemiol* 179:135–144. <https://doi.org/10.1093/aje/kwt254>
- Allison SP, Lobo DN (2004) Fluid and electrolytes in the elderly. *Curr Opin Clin Nutr Metab Care* 7:27–33. <https://doi.org/10.1097/00075197-200401000-00006>
- Atilasoy E, Holt PR (1993) Gastrointestinal proliferation and aging. *J Gerontol* 48:B43–49. <https://doi.org/10.1093/geronj/48.2.b43>
- Beck LH (2000) The aging kidney. Defending a delicate balance of fluid and electrolytes. *Geriatrics* 55:26–28
- Bhanu C, Avgerinou C, Kharicha K, Bauernfreund Y, Croker H, Liljas A et al (2020) 'I've never drunk very much water and I still don't, and I see no reason to do so': a qualitative study of the views of community-dwelling older people and carers on hydration in later life. *Age Ageing* 49:111–118. <https://doi.org/10.1093/ageing/afz141>
- Branca JJV, Gulisano M, Nicoletti C (2019) Intestinal epithelial barrier functions in ageing. *Ageing Res Rev* 54:100938. <https://doi.org/10.1016/j.arr.2019.100938>
- Clarke MM, Stanhewicz AE, Wolf ST, Cheuvront SN, Kenefick RW, Kenney WL (2019) A randomized trial to assess beverage hydration index in healthy older adults. *Am J Clin Nutr* 109:1640–1647. <https://doi.org/10.1093/ajcn/nqz009>
- Dangin M, Guillet C, Garcia-Rodenas C, Gachon P, Bouteloup-Demange C, Reiffers-Magnani K et al (2003) The rate of protein digestion affects protein gain differently during aging in humans. *J Physiol* 549:635–644. <https://doi.org/10.1113/jphysiol.2002.036897>
- Devanand V, Chithrapavai SU (2013) Correlation of 2 hours and 24 hours creatinine clearance in renal donors after unilateral nephrectomy. *J Clin Diagn Res* 7:2119–2121. <https://doi.org/10.7860/JCDR/2013/5696.3447>
- Dontas AS, Marketos SG, Papanayiotou P (1972) Mechanisms of renal tubular defects in old age. *Postgrad Med J* 48:295–303. <https://doi.org/10.1136/pgmj.48.559.295>
- Dubeau CE (2006) The aging lower urinary tract. *J Urol* 175:S11–15. [https://doi.org/10.1016/S0022-5347\(05\)00311-3](https://doi.org/10.1016/S0022-5347(05)00311-3)
- Eaton DC, Pooler JP (2018) Basic Renal Processes for Sodium Chloride and Water. Vander's Renal Physiology, 9th edn. McGraw-Hill Education, New York, NY
- El-Sharkawy AM, Sahota O, Maughan RJ, Lobo DN (2014) The pathophysiology of fluid and electrolyte balance in the older adult surgical patient. *Clin Nutr* 33:6–13. <https://doi.org/10.1016/j.clnu.2013.11.010>
- Godfrey H, Cloete J, Dymond E, Long A (2012) An exploration of the hydration care of older people: a qualitative study. *Int J Nurs Stud* 49:1200–1211. <https://doi.org/10.1016/j.ijnurstu.2012.04.009>
- Hall JE (2016) Renal dysfunction, rather than nonrenal vascular dysfunction, mediates salt-induced hypertension. *Circulation* 133:894–906. <https://doi.org/10.1161/CIRCULATIONAHA.115.018526>
- Holt PR, Rosenberg IH, Russell RM (1989) Causes and consequences of hypochlorhydria in the elderly. *Dig Dis Sci* 34:933–937. <https://doi.org/10.1007/BF01540281>
- Hooper L, Bunn D, Jimoh FO, Fairweather-Tait SJ (2014) Water-loss dehydration and aging. *Mech Ageing Dev* 136–137:50–58. <https://doi.org/10.1016/j.mad.2013.11.009>
- Huppertz T, Chia LW (2021) Milk protein coagulation under gastric conditions: a review. *Int Dairy J* 113:104882. <https://doi.org/10.1016/j.idairyj.2020.104882>
- Kenney WL, Chiu P (2001) Influence of age on thirst and fluid intake. *Med Sci Sports Exerc* 33:1524–1532. <https://doi.org/10.1097/00005768-200109000-00016>
- Laforenza U (2012) Water channel proteins in the gastrointestinal tract. *Mol Aspects Med* 33:642–650. <https://doi.org/10.1016/j.mam.2012.03.001>
- Leiper JB (1998) Intestinal water absorption—implications for the formulation of rehydration solutions. *Int J Sports Med* 19(Suppl 2):S129–132. <https://doi.org/10.1055/s-2007-971977>
- Levey AS, Stevens LA, Schmid CH, Zhang (Lucy) Y, Castro AF, Feldman HI et al (2009) A new equation to estimate glomerular filtration rate. *Ann Intern Med* 150:604–612
- Lindeman RD, Tobin J, Shock NW (1985) Longitudinal studies on the rate of decline in renal function with age. *J Am Geriatr Soc* 33:278–285. <https://doi.org/10.1111/j.1532-5415.1985.tb07117.x>
- Luchette FA, Yelon JA (2017) Geriatric Trauma and Critical Care. Springer
- Luckey AE, Parsa CJ (2003) Fluid and electrolytes in the aged. *Arch Surg* 138:1055–1060. <https://doi.org/10.1001/archsurg.138.10.1055>
- Luft FC, Rankin LI, Bloch R, Weyman AE, Willis LR, Murray RH et al (1979) Cardiovascular and humoral responses to extremes of sodium intake in normal black and white men. *Circulation* 60:697–706. <https://doi.org/10.1161/01.CIR.60.3.697>
- Maughan RJ, Watson P, Cordery PA, Walsh NP, Oliver SJ, Dolci A et al (2016) A randomized trial to assess the potential of different beverages to affect hydration status: development of a beverage hydration index. *Am J Clin Nutr* 103:717–723. <https://doi.org/10.3945/ajcn.115.114769>
- Maughan RJ, Watson P, Cordery PAA, Walsh NP, Oliver SJ, Dolci A et al (2019) Sucrose and sodium but not caffeine content influence

- the retention of beverages in humans under euhydrated conditions. *Int J Sport Nutr Exerc Metab* 29:51–60. <https://doi.org/10.1123/ijsnem.2018-0047>
- Moreno-Agostino D, Daskalopoulou C, Wu YT et al (2020) The impact of physical activity on healthy ageing trajectories: evidence from eight cohort studies. *Int J Behav Nutr Phys Act* 17:92. <https://doi.org/10.1186/s12966-020-00995-8>
- Musso CG, Jauregui JR (2014) Renin-angiotensin-aldosterone system and the aging kidney. *Expert Rev Endocrinol Metab* 9(6):543–546. <https://doi.org/10.1586/17446651.2014.956723>
- Musso C, Liakopoulos V, De Miguel R, Imperiali N, Algranati L (2006) Transtubular potassium concentration gradient: comparison between healthy old people and chronic renal failure patients. *Int Urol Nephrol* 38:387–390. <https://doi.org/10.1007/s11255-006-0059-5>
- Musso CG, Alvarez Gregori J, Jauregui JR, Macías Núñez JF (2012) Creatinine, urea, uric acid, water and electrolytes renal handling in the healthy oldest old. *World J Nephrol* 1:123–126. <https://doi.org/10.5527/wjn.v1.i5.123>
- Musso CG, Oreopoulos DG (2011) Aging and physiological changes of the kidneys including changes in glomerular filtration rate. *Nephron Physiol* 119(Suppl 1):p1-5. <https://doi.org/10.1159/000328010>
- Paterson DH, Warburton DE (2010) Physical activity and functional limitations in older adults: a systematic review related to Canada's physical activity guidelines. *Int J Behav Nutr Phys Act* 7:38. <https://doi.org/10.1186/1479-5868-7-38>
- Rolls BJ, Phillips PA (1990) Aging and disturbances of thirst and fluid balance. *Nutr Rev* 48:137–144. <https://doi.org/10.1111/j.1753-4887.1990.tb02915.x>
- Rowe JW, Andres R, Tobin JD, Norris AH, Shock NW (1976) The effect of age on creatinine clearance in men: a cross-sectional and longitudinal study. *J Gerontol* 31:155–163. <https://doi.org/10.1093/geronj/31.2.155>
- Rowe TA, Juthani-Mehta M (2013) Urinary tract infection in older adults. *Aging Health*. <https://doi.org/10.2217/ahe.13.38>
- Russell RM (1992) Changes in gastrointestinal function attributed to aging. *Am J Clin Nutr* 55:1203S–1207S. <https://doi.org/10.1093/ajcn/55.6.1203S>
- Sands JM (2012) Urine concentrating and diluting ability during aging. *J Gerontol A Biol Sci Med Sci* 67:1352–1357. <https://doi.org/10.1093/gerona/gls128>
- Schlanger LE, Bailey JL, Sands JM (2010) Electrolytes in the Aging. *Adv Chronic Kidney Dis* 17:308–319. <https://doi.org/10.1053/j.ackd.2010.03.008>
- Silva FG (2005) The aging kidney: a review – part I. *Int Urol Nephrol* 37:185–205. <https://doi.org/10.1007/s11255-004-0873-6>
- Stachenfeld NS, Mack GW, Takamata A, DiPietro L, Nadel ER (1996) Thirst and fluid regulatory responses to hypertonicity in older adults. *Am J Physiol* 271:R757–765. <https://doi.org/10.1152/ajpregu.1996.271.3.R757>
- Volkert D, Beck AM, Cederholm T, Cruz-Jentoft A, Goisser S, Hooper L et al (2019) ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr* 38:10–47. <https://doi.org/10.1016/j.clnu.2018.05.024>
- Wolf ST, Stanhewicz AE, Clarke MM, Cheuvront SN, Kenefick RW, Kenney WL (1985) Age-related differences in water and sodium handling after commercial hydration beverage ingestion. *J Appl Physiol* (1985) 2019(126):1042–1048. <https://doi.org/10.1152/jappphysiol.01094.2018>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.