

RESEARCH ARTICLE

Open Access



Sleep and BMI in South African urban and rural, high and low-income preschool children

Dale E. Rae^{1*} , Simone A. Tomaz^{1†}, Rachel A. Jones², Trina Hinkley³, Rhian Twine⁴, Kathleen Kahn^{4,5,6}, Shane A. Norris⁷ and Catherine E. Draper^{1,7}

Abstract

Background: The extent to which income setting or rural and urban environments modify the association between sleep and obesity in young children is unclear. The aims of this cross-sectional observational study were to (i) describe and compare sleep in South African preschool children from rural low-income (RL), urban low-income (UL) and urban high-income (UH) settings; and (ii) test for associations between sleep parameters and body mass index (BMI).

Methods: Participants were preschoolers (5.2 ± 0.7 y, 49.5% boys) from RL ($n = 111$), UL ($n = 65$) and UH ($n = 22$) settings. Height and weight were measured. Sleep, sedentary behaviour and physical activity were assessed using accelerometry.

Results: UL children had higher BMI z-scores (median: 0.39; interquartile range: $-0.27, 0.99$) than the UH ($-0.38; -0.88, 0.11$) and RL ($-0.08; -0.83, 0.53$) children ($p = 0.001$). The UL children had later bedtimes ($p < 0.001$) and wake-up times ($p < 0.001$) and shorter 24 h ($p < 0.001$) and nocturnal ($p < 0.001$) sleep durations than the RL and UH children. After adjusting for age, sex, setting, SB and PA, for every hour less sleep obtained (24 h and nocturnal), children were 2.28 (95% CI: 1.28–4.35) and 2.22 (95% CI: 1.27–3.85) more likely, respectively, to belong to a higher BMI z-score quartile.

Conclusions: Shorter sleep is associated with a higher BMI z-score in South African preschoolers, despite high levels of PA, with UL children appearing to be particularly vulnerable.

Keywords: Sleep, Adiposity, Early childhood, Low- and middle-income country

Background

Insufficient or short sleep has been associated with obesity and increased cardiometabolic risk in adults [1–3] and children [4, 5]. Duration of sleep in children has been shown to vary between countries, ethnic groups

and income settings. The International Study of Childhood Obesity, Lifestyle and the Environment affirms that sleep duration and timing in children aged 9–11 years differs between countries and that associations between sleep parameters and activity or sedentary parameters were stronger in the high- compared to low-income countries [6]. Within high-income countries, sleep duration and patterns vary between children from different ethnic groups, with non-white children or minority groups being more likely to have shorter sleep [7] or later bedtimes [8]. There is also evidence to suggest that

* Correspondence: Dale.Rae@uct.ac.za

[†]Dale E. Rae and Simone A. Tomaz contributed equally as co-first authors.

¹Health through Physical Activity, Lifestyle and Sport Research Centre & Division of Exercise Science and Sports Medicine, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

children from higher income families have longer sleep [9], earlier bedtimes [8] and better sleep efficiency [10] than those from lower income settings.

South Africa (SA) is an upper middle-income country characterised by large discrepancies in socioeconomic status (SES) with nearly half of the population (49.2%) living in poverty [11] in both rural and urban settings. It is undergoing an epidemiological transition; obesity is on the rise across all age groups [12] and non-communicable diseases are now responsible for more deaths annually than infectious diseases [13].

Preschool-aged children warrant investigation since early childhood is a period understood to predict obesity in adolescence [14] and 13% of preschool children in SA are overweight or obese [12]. For 3 to 5-year-old children, 10–13 h of sleep is recommended per 24 h period (comprising nocturnal sleep and daytime naps) to promote optimal health and development [15–18]. Only one study has been published on objectively measured sleep in SA preschool children. This study observed short nocturnal sleep (9.3 ± 0.8 h) in preschool-aged children in an urban, low-income setting in SA, where even after the inclusion of daytime naps, 24 h sleep duration was less than 10 h in 38% of the children [19]. Whether or not this observation of shorter sleep is also present in the rural setting or is a consequence of income status has not been established in this age group. The aims of this study were to (i) describe and compare sleep in preschool-aged South African children from rural low-income, urban low-income and urban high-income settings and (ii) test for associations between sleep and body mass index.

Methods

This cross-sectional, observational study is a secondary analysis of a previously published, separate study, investigating body composition, physical activity and gross motor skills in urban and rural preschoolers [20]. Only relevant methodological components are described in more detail below. The present study includes 198 of the original 268 participants for whom valid sleep datasets were available. Compliance within the urban high-income was very poor, resulting in a small group size.

Setting

Data were collected from children (5.19 ± 0.69 years) living in rural low-income (RL, $n = 111$), urban low-income (UL, $n = 65$) and urban high-income (UH, $n = 22$) settings. Data collection took place between July and September 2012 (UH and UL), and between June and August 2014 (RL). While a range of geographical settings was included, this sample was not nationally representative. The RL site was a rural village in north-east SA characterised by high unemployment [21], and

households with limited access to electricity and running water [22], and a high dependence on social grants. Access to this site was facilitated through the SAMRC/Wits Rural Public Health and Health Transitions Research Unit (Agincourt). Both urban settings were in south-west SA. The UL setting was a suburb comprising a mix of informal and low-cost housing in which common challenges are unemployment, overcrowding and crime. The UH setting comprised adjacent suburbs where the population density is approximately 15 times lower than the UL setting. The area has high-quality private and public schools, private health facilities and services, public parks and green spaces, expensive retailers and well-serviced amenities.

Participants and recruitment

UH and UL preschools were selected using convenience sampling and were intentionally economically diverse, taking into account geographical location and community-level factors indicating socioeconomic circumstances, such as those mentioned above. Recruitment of the RL preschool was coordinated through the Public Engagement Office of the SAMRC/Wits-Agincourt Unit. All parents were provided with written information about the study, as well as with instructions (in their home language) for the use of accelerometers. Where requested, parent meetings were arranged to assist with recruitment. All children between the ages of 3 and 5 years who were: attending preschool; willing and able to take part in the testing for this study; and for whom written informed parental consent was obtained, were included in this study.

Ethical approval was obtained from the University of Cape Town Human Research Ethics Committee (HREC Ref. No.: 237/2012), the University of the Witwatersrand Human Research Ethics Committee (Medical) (M140250), and the Mpumalanga Provincial Departments of Health and Education.

Detailed procedures

Anthropometry

Height (m) and weight (kg) were measured to calculate body mass index (BMI, kg/m^2). Height and weight measurements were conducted with the children's shoes and heavy clothing removed. A portable stadiometer (Leicester 214 Transportable Stadiometer; Seca GmbH & Co, Hamburg, Germany) was used to measure height to the nearest millimetre. To measure body weight, the children stood centred on a calibrated scale (Soehnle 7840 Mediscale Digital; Soehnle Industrial Solutions GmbH, Backnang, Germany). Height and weight measurements were done by a trained member of the research team.

Age standardised scores for BMI (BMI z-score), height (height-for-age z-score, HAZ) and weight (weight-for-

age z-score WAZ) were computed using the World Health Organisation AnthroPlus software (<http://www.who.int/growthref/tools/en/>). The International Obesity Task Force cut-offs were used to classify children based on the following BMI categories: thinness, normal weight, overweight, obese and morbidly obese [23]. For the regression analyses, since the BMI z-score data were not normally distributed, BMI z-score quartile groups were used: quartile 1: $n = 50$, median: -1.21 (interquartile range: $-1.48, -0.88$); quartile 2: $n = 49$, -0.28 ($-0.49, -0.11$); quartile 3: $n = 50$, 0.24 ($0.13, 0.45$); quartile 4: $n = 49$, 1.13 ($0.82, 1.48$).

Sleep, sedentary behaviour and physical activity

Actigraph GT3X+ accelerometers (Actigraph LLC, Pensacola, FL; USA) were used for the objective measurement of sleep, sedentary behaviour (SB) and physical activity (PA) over a 7-day period as previously described [19]. For a valid sleep dataset, a child needed to have at least 3 valid sleep nights (> 160 min), including at least 1 weekend night. Sunday to Thursday nights were taken to represent week nights, Friday and Saturday nights weekend nights, Monday to Friday mornings week mornings and Saturday and Sunday mornings weekend mornings. In the absence of sleep diaries or an accepted sleep-wake algorithm for hip- or waist-worn ActiGraph devices in preschool-aged children, visual inspection [24] was used to obtain nocturnal sleep variables (bedtime, hh:mm; wake-up time, hh:mm; time-in-bed, h), and daytime nap duration (h) using ActiLife version 6 (ActiLife software; Pensacola, FL; USA). Compliance with sleep guidelines for this age group [15–18] was determined using 24 h sleep duration (nocturnal time-in-bed plus daytime nap duration) and children were classified as meeting sleep guidelines if their average 24 h sleep duration was between 10 and 13 h.

Datasets for SB and PA were deemed valid if they had 7 h of valid wear-time (excluding sleep time) on 3 weekdays and 1 weekend day [25]. Non-wear time was defined as 20 min of consecutive zeroes and was removed from the data [26]. Cut-off points chosen to identify SB and moderate- to vigorous-intensity physical activity (MVPA) in this study were < 25 counts/15 s [27] and > 420 counts/15 s [28], respectively. Duration (min) of time spent in SB, light physical activity (LPA, > 25 and < 420 counts/15 s), MVPA and total PA (light- to vigorous-intensity physical activity, LMVPA) were calculated. Children were classified as meeting South African [18] and World Health Organisation [17] PA guidelines for this age group if they spent an average of ≥ 180 min/day in LMVPA, inclusive of 60 min/day of 'energetic play' (operationalised as MVPA).

Data and statistical analyses

Data are presented as mean \pm standard deviation, median (interquartile range) or percentage. The Shapiro-Wilks test was used to assess normality of data. Between-group analyses were conducted using an independent t-test, Mann-Whitney U test, one-way analysis of variance (ANOVA) or Kruskal-Wallis tests. Post hoc analyses were conducted using Bonferroni or Mann-Whitney U tests. Distributions were compared using Pearson's Chi-squared and correlations determined using Pearson's r or Spearman's ρ tests. Ordered logistic regression analyses were used to evaluate associations between sleep and BMI. The dependent variable was BMI z-score quartile group and the independent variables were 24 h sleep duration and nocturnal time-in-bed. Regression analyses were adjusted for age, sex, setting, SB and MVPA a priori since these variables have known associations with sleep and BMI. Data were analysed using Stata (v13, STATA Corp, College Station, TX) and significance accepted for $p < 0.05$.

Results

Descriptive characteristics

Descriptive characteristics of the participants stratified by sex and setting are presented in Table 1. The proportion of boys ($n = 98$) and girls ($n = 100$) in the study was similar ($p = 0.841$). The boys were older ($p = 0.037$), had higher BMI ($p = 0.009$) and BMI z-scores ($p = 0.017$) and accumulated more MVPA each day ($p < 0.001$) than the girls. Post hoc analyses indicate that the RL children were younger than the UL ($p < 0.001$) children, although all were preschool-aged. The UL children had higher BMI ($p = 0.001$) and BMI z-scores ($p = 0.017$) than the RL (BMI: $p = 0.001$, BMI z-score: $p = 0.001$) and UH (BMI: $p = 0.002$, BMI z-score: $p = 0.002$) children. More UL children were classified as overweight/obese compared to the RL and UH groups ($p = 0.012$) and fewer UL children were classified as thin compared to the other two groups ($p = 0.016$). The UH children had significantly higher HAZ scores than the UL ($p = 0.008$) and RL ($p = 0.010$) children. The RL children engaged in less sedentary behavior than the UL ($p < 0.001$) and UH ($p = 0.006$) children, while the UH children accumulate less LPA and LMVPA compared to the UL (LPA: $p < 0.001$, LMVPA: $p < 0.001$) and RL (LPA: $p < 0.001$, LMVPA: $p < 0.001$) children. The majority of this cohort met the PA guidelines [17, 29] for this age group (98%), and this distribution did not differ between the three settings (RL: 98.2%, UL: 98.5%, UH: 95.5%; $\chi^2 = 0.812$, $p = 0.666$).

Sleep characteristics

Average, week and weekend sleep characteristics of the children are presented in Table 2. Boys napped less

Table 1 Descriptive characteristics of participants stratified by sex and group

	All (n = 198)	Boys (n = 98)	Girls (n = 100)	P-value	Rural Low (n = 111)	Urban Low (n = 65)	Urban High (n = 22)	P-value
Age (y)	5.19 ± 0.69	5.29 ± 0.70	5.09 ± 0.67	0.037	5.05 ± 0.64 ^a	5.36 ± 0.71 ^a	5.38 ± 0.74	0.006
BMI (kg/m ²)	15.4 (14.3, 16.2)	15.5 (14.9, 16.2)	14.9 (14.1, 16.1)	0.009	15.1 (14.1, 16.0)	15.8 (14.9, 16.7) ^b	14.8 (14.3, 15.5)	0.001
BMI-z score	0.05 (−0.72, 0.68)	0.16 (−0.34, 0.73)	−0.23 (−0.85, 0.53)	0.017	−0.08 (−0.83, 0.53)	0.39 (−0.27, 0.99) ^b	−0.38 (−0.88, 0.11)	0.001
HAZ	−0.31 ± 1.01	−0.34 ± 0.98	−0.27 ± 1.04	0.658	−0.36 ± 0.97	−0.42 ± 1.03 ^b	0.32 ± 0.99	0.007
WAZ	−0.17 ± 1.02	−0.09 ± 0.90	−0.25 ± 1.13	0.271	−0.30 ± 0.96	−0.02 ± 1.15	−0.02 ± 0.87	0.167
Thinness (%)	20.7	14.0	27.0	0.062	26.1	10.8 ^b	22.7	
Normal (%)	71.2	77.0	64.0		68.5	73.8	77.3	0.017
Overweight/obese (%)	8.1	7.0	9.0		5.4	15.4 ^b	0	
SB (min/d)	308.8 (279.3, 340.3)	308.2 (272.8, 337.3)	310.0 (281.6, 341.7)	0.509	292.1 (263.8, 330.6) ^b	321.1 (298.2, 345.7)	321.3 (304.5, 347.8)	< 0.001
LPA (min/d)	337.8 ± 37.2	334.3 ± 37.3	341.3 ± 39.7	0.072	343.6 ± 35.6	339.1 ± 37.1	304.9 ± 29.0 ^b	< 0.001
MVPA (min/d)	123.6 (102.4, 146.9)	138.2 (110.3, 155.4)	112.6 (91.9, 134.4)	< 0.001	121.8 (97.5, 146.9)	132.6 (112.8, 154.0)	112.0 (94.3, 132.0)	0.125
LMVPA (min/d)	464.1 ± 56.3	471.5 ± 56.9	456.8 ± 55.1	0.065	469.9 ± 54.8	469.8 ± 55.2	417.5 ± 46.9 ^b	< 0.001

Data are presented as mean ± SD, median with interquartile range or percentage

Significance was determined using independent t-tests, Mann Whitney-U, one-way ANOVA, Kruskal-Wallis ANOVA or Chi² analyses

Significance was accepted at $p < 0.05$

BMI Body mass index, HAZ Height-for-age z-score, WAZ Weight-for-age z-score, SB Sedentary behaviour, LPA Light-intensity physical activity, MVPA Moderate- to vigorous-intensity physical activity, LMVPA Light- to vigorous-intensity physical activity

^aPost hoc analyses indicate significant differences between two marked groups

^bPost hoc analyses indicate that marked group is significantly different to the other two groups

during the day on average ($p = 0.008$) and on weekends ($p = 0.002$) and had later bedtimes during the week ($p = 0.049$) than girls. Post hoc analyses indicate that the UL children had later bedtimes than the RL (average: $p < 0.001$, week: $p < 0.001$ and weekend: $p < 0.001$) and UH (average: $p < 0.001$, week: $p < 0.001$ and weekend: $p < 0.001$) children; later wake-up times than the RL (average: $p < 0.001$, week: $p < 0.001$ and weekend: $p < 0.001$) and UH (average: $p = 0.001$, week: $p = 0.027$ and weekend: $p < 0.001$) children; and shorter nocturnal time-in-bed durations than the RL (average: $p < 0.001$, week: $p < 0.001$, weekend: $p < 0.001$) and UH (average: $p < 0.001$, week: $p < 0.001$, weekend: $p < 0.001$) children. The RL children had significantly longer weekend nocturnal time-in-bed durations than the UH ($p < 0.001$) children. Daytime naps were identified in 55% ($n = 109$) of the children in this sample. There were no differences in daytime nap duration between the three groups, but more napping took place during the week (94%) than on weekends (20%, $p < 0.001$).

Figure 1 illustrates 24 h sleep durations of children in the RL, UL and UH groups on average (A), during the week (B) and on weekends (C). The UL group had shorter 24 h sleep durations compared to the RL (average: $p < 0.001$ and week: $p < 0.001$) and UH (average: $p < 0.001$ and week: $p < 0.001$) groups. On weekends, the RL children had longer 24 h sleep durations than the UH ($p < 0.001$) and UL ($p < 0.001$) children. Of the entire cohort, 80.8% met current sleep guidelines [15–18]. Fewer children in the UL group (55%) met the guidelines compared to those in the RL (92%) and UH (100%) groups ($p < 0.001$). Removing daytime naps (i.e. evaluating nocturnal time-in-bed) reduced compliance in all three groups to 87% (RL), 42% (UL) and 91% (UH).

0.001 and week: $p < 0.001$) groups. On weekends, the RL children had longer 24 h sleep durations than the UH ($p < 0.001$) and UL ($p < 0.001$) children. Of the entire cohort, 80.8% met current sleep guidelines [15–18]. Fewer children in the UL group (55%) met the guidelines compared to those in the RL (92%) and UH (100%) groups ($p < 0.001$). Removing daytime naps (i.e. evaluating nocturnal time-in-bed) reduced compliance in all three groups to 87% (RL), 42% (UL) and 91% (UH).

Correlation analyses

Within-group correlations between average 24 h sleep duration and nocturnal time-in-bed with BMI z-score are presented in Additional Figure A1. No significant relationships were observed, although there was a trend for 24 h sleep duration to be negatively correlated with nocturnal time-in-bed in the UL group ($n = 65$, $r = -0.241$, $p = 0.053$, Additional Figure A1B). No within-group correlations were observed between average bedtime, wake-up time and BMI z-score (Additional Figure A2). In all three groups, average nocturnal time-in-bed was negatively correlated with average bedtime and positively correlated with average wake-up time (Additional Figure A3).

Table 2 Sleep characteristics of participants stratified by sex and setting group

	All (n = 198)	Boys (n = 98)	Girls (n = 100)	P-value	Rural Low (n = 111)	Urban Low (n = 65)	Urban High (n = 22)	P-value
Average								
Bedtime (hh:mm)	20:08 (19:32, 21:02)	20:12 (19:38, 21:12)	20:01 (19:22, 21:00)	0.085	19:38 (19:11, 20:08)	21:20 (21:01, 21:41) ^c	19:56 (19:35, 20:22)	< 0.001
Wake-up time (hh:mm)	06:40 (06:22, 07:09)	06:42 (06:22, 07:18)	06:39 (06:22, 07:00)	0.242	06:26 (06:11, 06:44)	07:11 (06:53, 07:32) ^c	06:43 (06:21, 07:19)	< 0.001
Nocturnal time-in-bed (h)	10.48 ± 0.77	10.44 ± 0.07	10.52 ± 0.08	0.478	10.76 ± 0.68	9.92 ± 0.68 ^c	10.76 ± 0.61	< 0.001
Nap duration (h) ^a	0.97 (0.80, 1.24)	0.86 (0.77, 1.13)	1.07 (0.90, 1.41)	0.008	0.94 (0.77, 1.22)	1.03 (0.83, 1.33)	0.89 (0.81, 1.02)	0.367
Week								
Bedtime (hh:mm)	20:02 (19:29, 21:01)	20:13 (19:37, 21:11)	19:54 (19:19, 20:53)	0.049	19:40 (19:10, 20:06)	21:18 (20:59, 21:41) ^c	19:49 (19:29, 20:13)	< 0.001
Wake-up time (hh:mm)	06:34 (06:11, 06:59)	06:37 (06:11, 07:11)	06:31 (06:11, 06:52)	0.268	06:17 (06:04, 06:34)	07:04 (06:40, 07:24) ^c	06:44 (06:23, 07:05)	< 0.001
Nocturnal time-in-bed (h)	10.39 ± 0.80	10.33 ± 0.07	10.45 ± 0.09	0.310	10.63 ± 0.68	9.81 ± 0.70 ^c	10.90 ± 0.70	< 0.001
Nap duration (h) ^a	1.00 (0.80, 1.30)	0.86 (0.77, 1.12)	1.11 (0.91, 1.48)	0.002	1.01 (0.79, 1.31)	1.03 (0.85, 1.45)	0.83 (0.74, 0.95)	0.182
Weekend								
Bedtime (hh:mm)	20:13 (19:30, 21:11)	20:19 (19:33, 21:14)	20:05 (19:26, 21:02)	0.350	20:21 (19:58, 20:49)	21:31 (20:55, 22:03) ^c	19:35 (19:13, 20:13)	< 0.001
Wake-up time (hh:mm)	06:59 (06:31, 07:32)	07:04 (06:30, 07:39)	06:58 (06:32, 07:23)	0.276	06:37 (06:17, 07:21)	07:41 (07:05, 07:59) ^c	06:46 (06:20, 07:10)	< 0.001
Nocturnal time-in-bed (h)	10.72 (10.04, 11.41)	10.57 (10.02, 11.44)	10.81 (10.05, 11.37)	0.995	10.32 (10.04, 10.88) ^b	10.10 (9.63, 10.58) ^c	11.14 (10.52, 11.54) ^b	< 0.001
Nap duration (h) ^a	0.90 ± 0.32	0.83 ± 0.34	0.99 ± 0.30	0.132	0.89 ± 0.30	0.85 ± 0.37	0.95 ± 0.30	0.644

Data are presented as mean ± SD or median with interquartile range

Significance was determined using independent t-tests, Mann Whitney-U, one-way ANOVA or Kruskal-Wallis ANOVA tests

^aSample size for naps: Average: n = 103, weekdays: n = 103 and weekend days: n = 40


^bPost hoc analyses indicate significant differences between two marked groups


^cPost hoc analyses indicate that marked group is significantly different to the other two groups. Significance was accepted at p < 0.05

Regression analyses

The association between average 24 h sleep duration and BMI z-score quartiles for all participants is displayed in Fig. 2. There was a linear relationship between the two variables with the lowest BMI z-score quartile comprising longer sleeping children and the highest quartile comprising shorter-sleeping children (p < 0.001). Table 3

summarises the ordered logistic regression results for BMI-z quartile (dependent variable) and average 24 h sleep duration (independent variable, model 1) and average nocturnal time-in-bed (independent variable, model 2). For every hour less sleep obtained in a 24 h period, children were 2.38 times more likely to belong to a higher BMI z-quartile (95% confidence interval (CI):





1.28, 4.35, $p = 0.006$) when adjusting for sex, age, setting, sedentary behaviour and LMVPA. Additionally, UL children were 2.91 times more likely to fall into a higher BMI z-quartile than UH children (95% CI: 1.15, 7.40; $p = 0.025$). For every hour less nocturnal sleep obtained, children were 2.22 times more likely to fall into a higher BMI z-quartile (95% CI: 1.27, 3.85; $p = 0.004$). Regression analyses were also run with average bedtime (model 3), average wake-up time (model 4) and average nap duration (model 5) as independent variables and BMI-z quartile (dependent variable), but no associations were observed (Additional Table A1).

Discussion

This is the first study to compare sleep (objectively measured or parent-report) in preschool children from a range of income settings in South Africa, and to

investigate the relationship between sleep and obesity in these groups. This study found that for every hour less 24 h or nocturnal sleep obtained, these preschool children were more than twice as likely to have a higher BMI, after adjusting for age, sex, setting, sedentary behaviour and physical activity. This is similar to what has been reported in two other studies in preschoolers from other countries [30, 31] and highlights the need to address sleep duration as a risk factor for obesity in South African children as young as 3–5 years in age. This is particularly relevant in South Africa where 13% of children under 5 years are overweight or obese [12]. Although not significant, the trend for average 24 h sleep duration to be negatively correlated with BMI z-score in the UL children suggests that the urban low-income setting is particularly vulnerable. Just over half of the children in the UL group (55%) met the sleep guidelines (i.e. 10–13 h in a 24 h period) [15–18] compared to most of those in the RL (92%) and UH (100%) groups. Despite high levels of physical activity, children from the UL setting had higher BMI z-scores, slept less in a 24 h period and at night, and had later bedtimes and wake-up times than children from the RL and UH settings.

The shorter sleep observed in the low- compared to high-income children within the urban setting is consistent with what has been reported in children [32, 33] and adults [34] in other countries. Low-income settings are typically characterised by an adverse social environment (i.e. high disorder and low safety and social cohesion), which in turn has been associated with shorter [35], worse quality [36] sleep in adults. Furthermore, in low-income homes, housing type, crowding, number of sleeping rooms and people per bed may influence sleep duration and quality. For example, the association between lower SES and poorer subjective ratings of sleep and higher daytime sleepiness in 11-year-old children was mediated by a higher exposure to poor sleeping conditions (noisy, light, uncomfortable, too hot or cold)

Table 3 Ordered logistic regression models associating BMI-z quartile (dependent variable) and average 24-h sleep duration (Model 1) and nocturnal time-in-bed (Model 2)

Model 1				Model 2					
		OR	95% CI	p-value			OR	95% CI	p-value
Ave 24-h sleep duration		0.42	0.23, 0.78	0.006	Ave nocturnal time-in-bed		0.45	0.26, 0.79	0.005
Covariates	Sex: Boy v girl	0.61	0.36, 1.03	0.066	Covariates	Sex: Boy v girl	0.57	0.33, 0.97	0.037
	Age	0.78	0.51, 1.20	0.261		Age	0.72	0.46, 1.13	0.153
	Group: UL v UH	2.91	1.15, 7.40	0.025		Group: UL v UH	2.48	0.96, 6.38	0.060
	RL v UH	1.14	0.48, 2.69	0.773		RL v UH	1.15	0.49, 2.73	0.747
	Sedentary Behaviour	0.99	0.98, 1.00	0.009		Sedentary Behaviour	0.99	0.98, 1.00	0.018
	LMVPA	0.99	0.98, 1.00	0.128		LMVPA	1.00	0.99, 1.00	0.254
Model: n = 198, LR chi2 = 31.14, p < 0.001				Model: n = 198, LR chi2=31.62, p < 0.001					

OR Odds ratio, CI Confidence interval, UL Urban low-income group, UH Urban high-income group, RL Rural low-income group, LMVPA Low- to vigorous-intensity physical activity

[37]. Additionally, 1-year-olds from families of lower SES and poor home environment quality are more likely to have shorter, less consistent sleep timing at the age of 8 years [38].

Within low-income settings, we observed shorter sleep in the urban (9.90 ± 0.68 h) compared to rural (10.76 ± 0.68 h) environment. Parent-reported sleep of a group of preschoolers comprising both urban and rural American Indian preschool has been described (10.15 ± 0.97 h), but urban and rural children were not compared [31]. Similar to our results, but in slightly older children (6–10 year-olds), shorter sleep duration was reported in Thai children living in the city compared to rural areas [39]. No difference in self-reported sleep duration, however, was observed between urban and rural 12-year-old Ugandan children [40]. One possible explanation for the RL children obtaining longer sleep in the present study might relate to their being fewer environmental barriers to good sleep (noise, light, crowding, safety) in a rural compared to urban home, despite other disadvantages of low SES.

Nocturnal time-in-bed was driven by both later bedtimes and wake-up times in all three groups, and later wake-up times observed in the UL children were unable to offset the shortened sleep opportunity caused by later bedtimes. This raises two important points. First, later bedtimes might reflect limited sleep health knowledge amongst the parents or caregivers of this group and, as such, less enforcement of an earlier bedtime or bedtime routine. Indeed it has been shown that preschool children in lower SES homes are less likely to have regular bedtime routines [8]. A longitudinal study observed that 5–9 year olds with no bedtime routine or age-appropriate bedtimes were more likely to have higher BMIs in adolescence [41]. Second, the later sleep timing (both later bedtime and wake-up time) seen in the UL group may entrain a later circadian rhythm in these children. Not only would this make it more difficult to obtain longer sleep when earlier wake-up times are required for school or extramural activities, later sleep timing has also been associated with less healthy food choices (fewer vegetables and healthy proteins and more processed or fried foods) in preschool children in the USA [42].

Daytime napping rescued 24 h sleep duration to some extent in the UL group. This has also been observed in another SA study of low-income preschool children from a different urban low-income setting [19]. The concern is that while napping may well be typical at preschools, once children enter the primary school phase, daytime napping usually falls away. If these UL children are unable to lengthen their nocturnal sleep in the absence of daytime napping, they may be at risk for chronic sleep deprivation in their crucial foundation years of school.

Finally, there has been a recent trend to observe movement patterns encompassing physical activity, sedentary behaviour, screen time and sleep over a 24 h period in relation to health [17, 43, 44]. The findings from this study compliment the sleep message of the new SA 24 h movement guidelines [18] in which parents, caregivers and teachers at early childhood development facilities are encouraged to help their children attain the 10–13 h of sleep recommended for preschool children.

A strength of this study is that sleep was measured objectively and we are among the first to describe and compare sleep in an economically diverse sample of young children. One limitation relates to light exposure, since the urban and rural sites were located in different parts of the country with varying day lengths. We do not believe this to have had a major effect on sleep timing in the present study since the day length difference between the two sites was only 30 min. We also acknowledge that since our data were collected during the winter months (June to August), the results may be open to seasonality influences. Again, however, we estimate this effect to be minimal since seasonal difference between daylight hours in summer and winter (10–14 h) and even daytime temperatures are moderate. Thus the expected impact on sleep timing would certainly be less compared to countries at more extreme latitudes. A further limitation relates to the convenience sampling method used, which means that our data are not necessarily generalizable or representative of South Africa, and that there is a risk for selection bias. We also acknowledge the small sample size in the urban high-income group, for whom compliance with wearing the accelerometer at night significantly reduced the availability of valid sleep datasets in this group. Finally, future studies exploring the relationship between movement behaviours (physical activity, sedentary behaviour), sleep and obesity would benefit from including a dietary component. The use of a randomised sampling technique to ensure representation of the context in question should also be considered in future studies.

Conclusions

Our findings suggest that nearly half of the preschool aged children in this study living in an urban low-income setting in South Africa may not be obtaining sufficient sleep and, despite high levels of physical activity, may be at risk for weight gain. Given the importance of maintaining a healthy weight in this age group to protect them from obesity in adolescence, they may well benefit from interventions designed to promote healthy sleep. These might include education of parents and caregivers regarding the importance of sleep for the physical, emotional and cognitive well-being of preschool-aged children, as well as guidelines regarding age-appropriate

sleep durations and bedtime routines. While these data relate to South African preschoolers conveniently sampled from three small but diverse communities, they may be applicable to any country with social disparity or with existing rural communities.

Abbreviations

ANOVA: Analysis of variance; BMI: Body mass index; BMI z-score: Body mass index-for-age score; CI: Confidence interval; HAZ: Height-for-age score; PA: Physical activity; LPA: Light-intensity physical activity; LMVPA: Light- to vigorous-intensity physical activity; MVPA: Moderate- to vigorous-intensity physical activity; OR: Odds ratio; SA: South Africa; SB: Sedentary behaviour; SD: Standard deviation; SES: Socioeconomic status; RL: Rural low-income; UH: Urban high-income; UL: Urban low-income; WAZ: Weight-for-age score

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12889-021-10591-5>.

Additional file 1: Figure A1. Correlations between average 24h sleep duration (A, B, C), average nocturnal time-in-bed (D, E, F) and BMI z-score for the three groups. BMI z-score: body mass index age-standardised score. Analyses were performed using Pearson's correlation test.

Additional file 2: Figure A2. Correlations between average bedtime (A, B, C), average wake-up time (D, E, F) and BMI z-score. BMI z-score: body mass index age-standardised score. Analyses were performed using Pearson's correlation or Spearman's rho tests.

Additional file 3: Figure A3. Correlations between average nocturnal time-in-bed and average bedtime (A, B, C) with average wake-up time (D, E, F). Analyses were performed using Pearson's or Spearman's correlation tests.

Additional file 4: Table A1. Ordered logistic regression models associating BMI z-score quartile (dependent variable) with average bedtime (Model 3), wake-up time (Model 4) and nap duration (Model 5).

Acknowledgements

The authors would like to thank local fieldworkers Rose Mnisi and the late Vivian Nzima. We would also like to thank all preschools who participated in this research, as well as the SAMRC/Wits-Agincourt Unit for facilitating research in Bushbuckridge.

Authors' contributions

DR carried out the analyses, drafted, reviewed and revised the manuscript. ST conceptualized and designed the study, designed the data collection instruments and collected data, carried out analyses, reviewed and revised the manuscript. TH and RJ contributed to study design, supervised data collection, provided input on the management and analysis of data, reviewed and revised the manuscript. RT, KK and SN – contributed to the study design, recruitment and data collection, provided strategic research oversight reviewed and revised the manuscript. CD conceptualized and designed the study, designed the data collection instruments, collected data and supervised data collection, and critically reviewed the manuscript for important intellectual content. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Funding

Funding support for this research was received from the South African National Research Foundation (CPRR13090934186) and the University of Cape Town University Research Committee. For ST, the support of the DST-NRF Centre of Excellence (CoE) in Human Development at the University of the Witwatersrand, Johannesburg in the Republic of South Africa towards this research is hereby acknowledged. The SAMRC/Wits-Agincourt Unit is supported by the University of the Witwatersrand, South African Medical Research Council and the Wellcome Trust, UK (grants 058893/Z/99/A; 069683/Z/02/Z; 085477/Z/08/Z; 085477/B/08/Z). Opinions expressed and conclusions arrived at, are those of the author and are not to be attributed to the CoE in

Human Development. TH was funded by a National Health and Medical Research Council Early Career Fellowship (APP1070571).

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

All parents were provided with written information about the study and gave written informed consent for their children to take part in this study. Ethical approval was obtained from the University of Cape Town Human Research Ethics Committee (HREC Ref. No.: 237/2012), the University of the Witwatersrand Human Research Ethics Committee (Medical) (M140250), and the Mpumalanga Provincial Departments of Health and Education.

Consent for publication

Not applicable.

Competing interests

CD is a member of the editorial board for BMC Public Health. The other authors declare that they have no competing interests.

Author details

¹Health through Physical Activity, Lifestyle and Sport Research Centre & Division of Exercise Science and Sports Medicine, Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa. ²Early Start, Faculty of the Arts Social Science and Humanities, University of Wollongong, Wollongong, Australia. ³Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Deakin University, Geelong, Australia. ⁴MRC/Wits Rural Public Health and Health Transitions Research Unit (Agincourt), School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa. ⁵Umeå Centre for Global Health Research, Umeå University, Umeå, Sweden. ⁶INDEPTH Network, Accra, Ghana. ⁷SAMRC/Wits Developmental Pathways for Health Research Unit, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa.

Received: 18 March 2020 Accepted: 8 March 2021

Published online: 23 March 2021

References

- Buxton OM, Marcelli E. Short and long sleep are positively associated with obesity, diabetes, hypertension, and cardiovascular disease among adults in the United States. *Soc Sci Med*. 2010;71(5):1027–36. <https://doi.org/10.1016/j.socscimed.2010.05.041> Elsevier Ltd.
- Lucassen EA, Rother KI, Cizza G. Interacting epidemics? Sleep curtailment, insulin resistance, and obesity. *Ann N Y Acad Sci*. 2012;1264(1):110–34. <https://doi.org/10.1111/j.1749-6632.2012.06655.x>.
- Altman NG, Izci-Balserak B, Schopfer E, Jackson N, Rattanaumpawan P, Gehrman PR, Patel NP, Grandner MA. Sleep duration versus sleep insufficiency as predictors of cardiometabolic health outcomes. *Sleep Med*. 2012;13(10):1261–70. <https://doi.org/10.1016/j.sleep.2012.08.005>.
- Cespedes EM, Rifas-Shiman SL, Redline S, Gillman MW, Peña M-M, Taveras EM. Longitudinal associations of sleep curtailment with metabolic risk in mid-childhood. *Obesity*. 2014;22(12):2586–92 John Wiley & Sons, Ltd.
- Li L, Fu J, Yu XT, Li G, Xu L, Yin J, et al. Sleep duration and cardiometabolic risk among Chinese school-aged children: do adipokines play a mediating role? *Sleep*. 2017;40(5):2510.
- Chaput JP, Katzmarzyk PT, LeBlanc AG, Tremblay MS, Barreira TV, Broyles ST, et al. Associations between sleep patterns and lifestyle behaviors in children: An international comparison. *Int J Obesity Suppl*. 2015;5(S2):S59–65 Nature Publishing Group.
- Anujoo KO, Vrijkotte TGM, Stronks K, Jean-Louis G, Agyemang CO. Ethnic differences in sleep duration at 5 years, and its relationship with overweight and blood pressure. *Eur J Pub Health*. 2016;26(6):1001–6. <https://doi.org/10.1093/eurpub/ckw084>.
- Hale L, Berger LM, LeBourgeois MK, Brooks-Gunn J. Social and demographic predictors of preschoolers' bedtime routines. *J Dev Behav Pediatr*. 2009;30(5):394–402. <https://doi.org/10.1097/DBP.0b013e3181ba0e64>.

9. Dube N, Khan K, Loehr S, Chu Y, Veugelaers P. The use of entertainment and communication technologies before sleep could affect sleep and weight status: a population-based study among children. *Int J Behav Nutr Phys Act*. 2017;14(1):97–112 International Journal of Behavioral Nutrition and Physical Activity.
10. El-Sheikh M, Kelly RJ, Buckhalt JA, Benjamin HJ. Children's sleep and adjustment over time: the role of socioeconomic context. *Child Dev*. 2010; 81(3):870–83 Blackwell Publishing Ltd.
11. Statistics South Africa. Men, women and children: Findings of the living conditions survey 2014/2015. Pretoria: Statistics South Africa; 2018. p. 1–65.
12. National Department of Health (NDoH), Statistics South Africa (Stats SA), South African Medical Research Council (SAMRC), ICF. South Africa demographic and health survey 2016 - key indicator report. Pretoria and Rockville: NDoH, Stats SA, SAMRC, and ICF; 2018. p. 1–24.
13. Statistics South Africa. Mortality and causes of death in South Africa, 2016: Findings from death notification. Pretoria: Statistics South Africa; 2018. p. 1–146.
14. Lundeen EA, Norris SA, Adair LS, Richter LM, Stein AD. Sex differences in obesity incidence: 20-year prospective cohort in South Africa. *Pediatr Obes*. 2016;11(1):75–80. <https://doi.org/10.1111/ijpo.12039>.
15. Hirshkowitz M, Whitton K, Albert SM, Alessi C, Bruni O, DonCarlos L, Hazen N, Herman J, Katz ES, Kheirandish-Gozal L, Neubauer DN, O'Donnell AE, Ohayon M, Peever J, Rawding R, Sachdeva RC, Setters B, Vitiello MV, Ware JC, Adams Hillard PJ. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. 2015;1(1):40–3. <https://doi.org/10.1016/j.sleh.2014.12.010>.
16. Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, Malow BA, Maski K, Nichols C, Quan SF, Rosen CL, Troester MM, Wise MS. Recommended amount of sleep for pediatric populations: a consensus statement of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2016;12(6):785–6. <https://doi.org/10.5664/jcsm.5866>.
17. World Health Organization. WHO guidelines on physical activity, sedentary behaviour and sleep for children under 5 years of age. Geneva: World Health Organization; 2019.
18. Draper CE, Tomaz SA, Biersteker L, Cook CJ, Couper J, de Milander M, Flynn K, Giese S, Krog S, Lambert EV, Liebenberg T, Mendoza C, Nunes T, Pienaar A, Priorieschi A, Rae DE, Rahbeeni N, Reilly JJ, Reynolds L, Samuels ML, Siljeur R, Urion J, van Wyk M, Okely AD. The South African 24-hour movement guidelines for birth to five years: an integration of physical activity, sitting behaviour, screen time and sleep. *J Phys Act Health*. 2020;17(1):109–19. <https://doi.org/10.1123/jpah.2019-0187>.
19. Tomaz SA, Priorieschi A, Watson ED, McVeigh JA, Rae DE, Jones RA, et al. Body mass index, physical activity, sedentary behavior, sleep, and gross motor skill proficiency in preschool children from a low- to middle-income urban setting. *J Phys Act Health*. 2019;16(7):525–32. <https://doi.org/10.1123/jpah.2018-0133>.
20. Draper CE, Tomaz SA, Stone M, Hinkley T, Jones RA, Louw J, Twine R, Kahn K, Norris SA. Developing intervention strategies to optimise body composition in early childhood in South Africa. *BioMed Res Int*. 2017; 2017(1):1–13. <https://doi.org/10.1155/2017/5283457> Hindawi.
21. Collinson MA. Striving against adversity: the dynamics of migration, health and poverty in rural South Africa. *Glob Health Action*. 2010;3(1):5080. <https://doi.org/10.3402/gha.v3i0.5080>.
22. Kahn K, Collinson MA, Gomez-Olive FX, Mokoena O, Twine R, Mee P, Afolabi SA, Clark BD, Kabudula CW, Khosa A, Khoza S, Shabangu MG, Silaule B, Tibane JB, Wagner RG, Garenne ML, Clark SJ, Tollman SM. Profile: Agincourt health and socio-demographic surveillance system. *Int J Epidemiol*. 2012;41(4):988–1001. <https://doi.org/10.1093/ije/dys115>.
23. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes*. 2012;7(4):284–94 John Wiley & Sons, Ltd (10.1111).
24. Chow CM, Wong SN, Shin M, Maddox R, Feilds K-L, Paxton K, Hawke C, Hazell P, Steinbeck K. Defining the rest interval associated with the main sleep period in actigraph scoring. *Nat Sci Sleep*. 2016;8:321–8. <https://doi.org/10.2147/NSS.S114969> Dove Press.
25. Hinkley T, O'Connell E, Okely AD, Crawford D, Hesketh K, Salmon J. Assessing volume of accelerometry data for reliability in preschool children. *Med Sci Sport Exerc*. 2012;44(12):2436–41. <https://doi.org/10.1249/MSS.0b013e3182661478>.
26. Janssen X, Basterfield L, Parkinson KN, Pearce MS, Reilly JK, Adamson AJ, et al. Objective measurement of sedentary behavior: impact of non-wear time rules on changes in sedentary time. *BMC Public Health*. 2015;15(1):504. <https://doi.org/10.1186/s12889-015-1847-6> BioMed Central.
27. Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective measures of physical activity for children. *J Sport Sci*. 2008;26(14): 1557–65. <https://doi.org/10.1080/02640410802334196> Routledge.
28. Janssen X, Cliff DP, Reilly JJ, Hinkley T, Jones RA, Batterham M, Ekelund U, Brage S, Okely AD. Validation and calibration of the activPAL™ for estimating METs and physical activity in 4–6 year olds. *J Sci Med Sport*. 2014; 17(6):602–6. <https://doi.org/10.1016/j.jsams.2013.10.252>.
29. Draper CE, Tomaz SA, Biersteker L, Cook CJ, Couper J, de Milander M, et al. The south African 24-hour movement guidelines for birth to 5 years: an integration of physical activity, sitting behavior, screen time, and sleep. *J Phys Act Health*. 2019;17(1):109–19 Human Kinetics.
30. Park SH, Park CG, Bahorski JS, Cormier E. Factors influencing obesity among preschoolers: multilevel approach. *Int Nurs Rev*. 2019;10(34):10.
31. Ingram DG, Irish LA, Tomayko EJ, Prince RJ, Cronin KA, Parker T, Kim K, Carmichael L, Grant VM, Sheche JN, Adams AK. Overnight sleep duration and obesity in 2–5 year-old American Indian children. *Pediatr Obes*. 2018; 13(7):406–12. <https://doi.org/10.1111/ijpo.12281>.
32. El-Sheikh M, Bagley EJ, Keiley M, Elmore-Staton L, Chen E, Buckhalt JA. Economic adversity and children's sleep problems: multiple indicators and moderation of effects. *Health Psychol*. 2013;32(8):849–59. <https://doi.org/10.1037/a0030413>.
33. Marco CA, Wolfson AR, Sparling M, Azuaje A. Family socioeconomic status and sleep patterns of young adolescents. *Behav Sleep Med*. 2011;10(1):70–80 Taylor & Francis Group.
34. Whinnery J, Jackson N, Rattanaumpawan P, Grandner MA. Short and long sleep duration associated with race/ethnicity, sociodemographics, and socioeconomic position. *Sleep*. 2014;37(3):601–11. <https://doi.org/10.5665/sleep.3508>.
35. Desantis AS, Diez Roux AV, Moore K, Baron KG, Mujahid MS, Nieto FJ. Associations of neighborhood characteristics with sleep timing and quality: the multi-ethnic study of atherosclerosis. *Sleep*. 2013;36(10):1543–51. <https://doi.org/10.5665/sleep.3054>.
36. Hill TD, Burdette AM, Hale L. Neighborhood disorder, sleep quality, and psychological distress: testing a model of structural amplification. *Health Place*. 2009;15(4):1006–13. <https://doi.org/10.1016/j.healthplace.2009.04.001>.
37. Bagley EJ, Kelly RJ, Buckhalt JA, El-Sheikh M. What keeps low-SES children from sleeping well: the role of presleep worries and sleep environment. *Sleep Med*. 2015;16(4):496–502. <https://doi.org/10.1016/j.sleep.2014.10.008>.
38. Doane LD, Breitenstein RS, Beekman C, Clifford S, Smith TJ, Lemery-Chalfant K. Early life socioeconomic disparities in children's sleep: the mediating role of the current home environment. *J Youth Adolesc*. 2019;48(1):56–70. <https://doi.org/10.1007/s10964-018-0917-3>.
39. Sirirassamee F, Chonchaiya W, Pruksananonda C. Sleep behaviors and sleep problems in school-aged children in Thailand. *J Med Assoc Thai*. 2015; 98(Suppl 9):S71–7.
40. Christoph MJ, Grigsby-Toussaint DS, Baingana R, Ntambi JM. Physical activity, sleep, and BMI percentile in rural and urban Ugandan youth. *Ann Glob Health*. 2017;83(2):311–9. <https://doi.org/10.1016/j.aogh.2017.04.005>.
41. Lee S, Hale L, Chang A-M, Nahmod NG, Master L, Berger LM, et al. Longitudinal associations of childhood bedtime and sleep routines with adolescent body mass index. *Sleep*. 2019;42(1):2627.
42. Jansen EC, Peterson KE, Lumeng JC, Kaciroti N, LeBourgeois MK, Chen K, et al. Associations between sleep and dietary patterns among low-income children attending preschool. *J Acad Nutr Diet*. 2019;119(7):1176–87. <https://doi.org/10.1016/j.jand.2019.01.008>.
43. Tremblay MS, Chaput J-P, Adamo KB, Aubert S, Barnes JD, Choquette L, Duggan M, Faulkner G, Goldfield GS, Gray CE, Gruber R, Janson K, Janssen I, Janssen X, Jaramillo Garcia A, Kuzik N, LeBlanc C, MacLean J, Okely AD, Poitras VJ, Rayner ME, Reilly JJ, Sampson M, Spence JC, Timmons BW, Carson V. Canadian 24-hour movement guidelines for the early years (0–4 years): an integration of physical activity, sedentary behaviour, and sleep. *BMC Public Health*. 2017;17(Suppl 5): 874. <https://doi.org/10.1186/s12889-017-4859-6>.

44. Okely AD, Ghera D, Hesketh KD, Santos R, Loughran SP, Cliff DP, et al. A collaborative approach to adopting/adapting guidelines - The Australian 24-Hour Movement Guidelines for the early years (birth to 5 years): an integration of physical activity, sedentary behavior, and sleep. *BMC Public Health*. 2017;17(Suppl 5):869 2nd ed. BioMed Central.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com