

Pooled analysis of physical activity, sedentary behaviour and sleep among children from 33 countries

Kar Hau Chong, PhD, School of Health and Society, Early Start, Faculty of the Arts, Social Science and Humanities, University of Wollongong, NSW 2522, Australia.

Thomas Suesse, PhD, National Institute for Applied Statistics Research Australia, School of Mathematics and Applied Statistics, Faculty of Engineering and Information Sciences, University of Wollongong, Wollongong, NSW 2522, Australia.

Penny L. Cross, BEng (Mat) (Hons), Research Services Office, Research and Innovation Division, University of Wollongong, NSW 2522, Australia.

Sarah T. Ryan, BSc (Hons), School of Health and Society, Early Start, Faculty of the Arts, Social Science and Humanities, University of Wollongong, NSW 2522, Australia.

Eivind Aadland, PhD, Department of Sport, Food and Natural Sciences, Faculty of Education, Arts and Sports, Western Norway University of Applied Sciences, Sogndal, Norway.

Oluwayomi Aoko, MSc, Human Kinetics and Health Education Department, University of Lagos, Lagos, Nigeria.

Ankhmaa Byambaa, BSc (Hons), National Centre for Public Health, Ulaanbaatar 13381, Mongolia.

Valerie Carson, PhD, Faculty of Kinesiology, Sport, and Recreation, University of Alberta, Edmonton, AB, T6G 2H9, Canada.

Jean-Philippe Chaput, PhD, Healthy Active Living and Obesity Research Group, Children's Hospital of Eastern Ontario Research Institute, Ottawa, ON K1H 8L1, Canada.

29

30 Hayley Christian, PhD, Telethon Kids Institute, University of Western Australia, Nedlands WA 6009,
31 Australia.

32

33 Dylan P. Cliff, PhD, School of Education, Early Start, Faculty of the Arts, Social Sciences and Humanities,
34 University of Wollongong, Northfields Ave, Wollongong NSW 2522, Australia.

35

36 Marieke De Craemer, PhD, Department of Rehabilitation Sciences, Ghent University, Corneel Heymanslaan
37 10, 9000 Ghent, Belgium.

38

39 Clarice Maria de Lucena Martins, PhD, Research Centre in Physical Activity, Health and Leisure, and
40 Laboratory for Integrative and Translational Research in Population Health, University of Porto, Porto,
41 Portugal.

42

43 Christine Delisle Nyström, PhD, Department of Biosciences and Nutrition, Karolinska Institutet, Neo, 141
44 83, Huddinge, Sweden.

45

46 Catherine E. Draper, PhD, South African MRC/Wits Developmental Pathways for Health Research Unit,
47 Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa.

48

49 Asmaa El Hamdouchi, PhD, Unité de Recherche en Nutrition et Alimentation, CNESTEN, Rabat, Morocco.

50

51 Alex Antonio Florindo, PhD, School of Arts, Sciences, and Humanities, University of Sao Paulo, Brazil.

52

53 Hongyan Guan, PhD, Child Healthcare Centre, Nurturing Care Research and Guidance Center, Beijing
54 Municipal Key Laboratory of Child Development and Nutriomics, Capital Institute of Pediatrics, Beijing,
55 China.

56

57 Amy S. Ha, PhD, Department of Sports Science and Physical Education, Kwok Sports Building, The
58 Chinese University of Hong Kong, Shatin, Hong Kong, China.

59

60 Najmeh Hamzavi Zarghani, PhD, Department of Health Education and Health Promotion, Faculty of
61 Medical Science, Tarbiat Modares University, Tehran, Iran.

62

63 Kylie D. Hesketh, PhD, Institute for Physical Activity and Nutrition (IPAN), School of Exercise and
64 Nutrition Science, Deakin University, 1 Gheringhap Street, Geelong, VIC 3220, Australia.

65

66 Mohammad Sorowar Hossain, PhD, Department of Emerging and Neglected Diseases, Biomedical Research
67 Foundation, Dhaka, Bangladesh.

68

69 Jajat Jajat, PhD, Faculty of Sport and Health Education, Universitas Pendidikan Indonesia, Bandung, Jawa
70 Barat, Indonesia.

71

72 Thanh Kim, MD, Department of Epidemiology, Faculty of Public Health, Pham Ngoc Thach University of
73 Medicine, Ho Chi Minh City, Vietnam.

74

75 Denise Koh, PhD, Centre for Education and Community Well-being, Faculty of Education,
76 Universiti Kebangsaan Malaysia, 43600 UKM Bangi, Malaysia.

77

78 Anna V. Kontsevaya, PhD, National Medical Research Center for Therapy and Preventive Medicine, 101990
79 Moscow, Russia.

80

81 Nicholas Kuzik, PhD, Children's Hospital of Eastern Ontario Research Institute, Ottawa, ON, Canada.

82

83 Marja H. Leppänen, PhD, Institute of Biomedicine, School of Medicine, University of Eastern Finland,
84 Kuopio, Finland.

85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112

Marie Löf, PhD, Department of Biosciences and Nutrition, Karolinska Institutet, Neo, 141 83, Huddinge, Sweden.

Himangi Lubree, MSc, Vadu Rural Health Program, KEM Hospital Research Centre, Rasta Peth, Pune, India.

Kim Meredith-Jones, PhD, Department of Medicine, University of Otago, PO Box 56, Dunedin, New Zealand.

Tawonga W. Mwase-Vuma, MPH, Centre for Social Research, University of Malawi, Zomba, Malawi.

Johan Y. Y. Ng, PhD, Department of Sports Science and Physical Education, Kwok Sports Building, The Chinese University of Hong Kong, Shatin, Hong Kong, China.

Rachel Novotny, PhD, Department of Human Nutrition, Food and Animal Sciences, College of Tropical Agriculture and Human Resources, University of Hawaii at Manoa, Honolulu, HI, 96822, USA.

Jackline Jema Nusurupia, MSc, Tanzania Food and Nutrition Centre, 22 Barack Obama Road, Dar es salaam, Tanzania.

Bang N. Pham, PhD, Papua New Guinea Institute of Medical Research, Goroka, Papua New Guinea.

Bee Koon Poh, PhD, Centre for Community Health Studies (ReaCH), Faculty of Health Sciences, Universiti Kebangsaan Malaysia, 50300 Kuala Lumpur, Malaysia.

John J. Reilly, PhD, Department of Psychological Sciences & Health, University of Strathclyde, Glasgow, Scotland.

113

114 Amanda E. Staiano, PhD, Division of Population and Public Health Science, Pennington Biomedical
115 Research Center, Baton Rouge, LA United States.

116

117 Kuston Sultoni, MPd, Faculty of Sport and Health Education, Universitas Pendidikan Indonesia, Bandung,
118 Jawa Barat, Indonesia.

119

120 Chiaki Tanaka, PhD, Department of Human Nutrition, Tokyo Kasei Gakuin University, 22 Sanbancho,
121 Chiyoda-ku, Tokyo, 102-8341 Japan.

122

123 Hong K. Tang, PhD, Department of Epidemiology, Faculty of Public Health, Pham Ngoc Thach University
124 of Medicine, Ho Chi Minh City, Vietnam.

125

126 Rachael Taylor, PhD, Department of Medicine, University of Otago, PO Box 56, Dunedin, New Zealand.

127

128 Simone A. Tomaz, PhD, Faculty of Health Sciences and Sport, University of Stirling, UK.

129

130 Mark S. Tremblay, PhD, Children's Hospital of Eastern Ontario Research Institute and Department of
131 Pediatrics, University of Ottawa, Ottawa, Canada.

132

133 Stewart G. Trost, PhD, School of Human Movement and Nutrition Sciences, University of Queensland,
134 Brisbane QLD 4001, Australia

135

136 Ali Turab, MPH, PHC Global, Karachi, Pakistan.

137

138 Susana Vale, PhD, ESE, Polytechnic of Porto, 4200-465 Porto, Portugal.

139

140 V. Pujitha Wickramasinghe, MD, Department of Paediatrics, Faculty of Medicine, University of Colombo,
141 Sri Lanka.

142

143 Anthony D. Okely, EdD, School of Health and Society, Early Start, Faculty of the Arts, Social Science and
144 Humanities, University of Wollongong, NSW 2522, Australia.

145

146 **Corresponding author:**

147 Kar Hau Chong, PhD, School of Health and Society, Early Start, Faculty of the Arts, Social Science and
148 Humanities, University of Wollongong, NSW 2522, Australia.

149 Email: khchong@uow.edu.au; Telephone: +614 3240 6560

150

151 **Date of the revision:** 28 June 2024

152 **Manuscript word count:** 3169

153 **Key Points**

154

155 **Question:** What is the global proportion of children aged 3- to 4-year-old who met the World Health
156 Organization guidelines for physical activity, sedentary behaviour and sleep?

157

158 **Findings:** Our pooled analyses of 7,017 children across 33 countries showed that only 14.3% of children met
159 the recommendations for physical activity, screen time and sleep duration. While differences were noted
160 between sexes, regions and country income levels, the compliance with the overall guidelines was
161 universally low.

162

163 **Meaning:** Identifying key factors and implementing contextually appropriate, effective programs and
164 policies is essential to tackle movement behaviour inequalities among 3- to 4-year-old children worldwide.

165 **Abstract**

166 **Importance:** The prevalence estimates of physical activity, sedentary behaviour and sleep (collectively
167 known as movement behaviours) in 3- and 4-year-old children worldwide remains uncertain.

168 **Objective:** To report the proportion of 3- and 4-year-old children who met the World Health Organization
169 guidelines for physical activity, sedentary behaviour and sleep across 33 countries.

170 **Design:** Pooled analysis of data from 14 cross-sectional studies (2008-2022) identified through systematic
171 reviews and personal networks.

172 **Setting:** Thirty-three countries of varying income levels across six geographical regions.

173 **Participants:** Each study site needed to have at least 40 children aged 3.0 to 4.9 years with valid
174 accelerometry and parent-/caregiver-reported screen time and sleep duration data.

175 **Exposures:** Time spent in physical activity was assessed by re-analysing accelerometry data using a
176 harmonised data processing protocol. Screen time and sleep duration were proxy-reported by parents or
177 caregivers.

178 **Main Outcomes and Measures:** The proportion of children who met the World Health Organization
179 guidelines for physical activity (≥ 180 minutes/day of total physical activity and ≥ 60 minutes/day of
180 moderate- to vigorous-intensity physical activity), screen time (≤ 1 hour/day), and sleep duration (10-13
181 hours/day) was estimated across countries and by World Bank income group and geographical region using
182 meta-analysis.

183 **Results:** Of the 7,017 children (51.1% boys) in this pooled analysis, 14.3% (95% CI, 9.7%-20.7%) met the
184 overall guidelines for physical activity, screen time and sleep duration. There was no clear pattern according
185 to income group: the proportion meeting the guidelines was 16.6% (95% CI, 10.4%-25.3%) in low- and
186 lower-middle-income countries, 11.9% (95% CI, 5.9%-22.5%) in upper-middle-income countries, and
187 14.4% (95% CI, 9.6%-21.1%) in high-income countries. The region with the highest proportion meeting the
188 guidelines was Africa (23.9%; 95% CI, 11.6%-43.0%), while the lowest proportion was the Americas (7.7%,
189 95% CI, 3.6%-15.8%).

190 **Conclusions and Relevance:** Most 3- and 4-year-old children do not meet the current World Health
191 Organization guidelines for physical activity, sedentary behaviour and sleep. Priority must be given to
192 understanding factors that influence these behaviours in this age group and to implementing contextually

- 193 appropriate programs and policies proven to be effective in promoting healthy levels of movement
- 194 behaviours.

195 **Introduction**

196 In 2019, the World Health Organization (WHO) published global guidelines for physical activity (PA),
197 sedentary behaviour and sleep (collectively referred to as movement behaviours) for children under the age
198 of five.¹ These guidelines are based on an integrated movement behaviour paradigm,² acknowledging the co-
199 dependencies of these behaviours and their synergistic effects on health. For children aged 3- to 4-year-old,
200 the guidelines recommend participating in at least 180 minutes of PA (of which at least 60 minutes should be
201 of moderate- to vigorous-intensity), not spending more than one hour in sedentary screen time, and having
202 10-13 hours of good quality sleep in a 24-hour day.¹ Meeting these guidelines is associated with better
203 adiposity status,³ psychosocial health⁴ and motor skills^{5,6} in this age group. To increase awareness of the
204 importance of healthy levels of movement behaviours in early childhood, it is important to know the
205 proportion of children meeting the guidelines globally.

206

207 Building on the systematic review by Rollo and colleagues⁷, 33 articles (representing 21 studies) published
208 during January 2015–August 2022 have examined adherence to the WHO guidelines for children aged 3 and
209 4 years, with the reported proportions ranged from 0% to 37%. Notably, 17 of the 21 studies (85%) were
210 conducted exclusively in high-income countries, indicating the lack of evidence among lower-income
211 countries. A more recent meta-analysis, including data from 26 articles, reported the overall proportion
212 meeting the guidelines was 11% among children aged 3 to 5 years.⁸ These estimates should, however, be
213 interpreted cautiously given the methodological variations across studies, particularly in the accelerometer
214 data processing methods applied to obtain PA estimates. Pooling data from these studies using the same data
215 processing protocols while complementing it with new data from more low- and lower-middle-income
216 countries would allow, for the first time, the ability to report the global proportion of children who meet the
217 WHO guidelines. This will contribute to the limited knowledge base on global prevalence estimates of
218 movement behaviours in young children⁹ and inform global and regional policies to promote healthy
219 movement behaviours from an early age.

220

221 In this paper, we conducted a pooled analysis to determine the proportion of children aged 3- to 4-year-old
222 who met the WHO recommended levels of PA, sedentary behaviour (operationalised and hereinafter referred
223 to as screen time), and sleep across 33 countries using a harmonised data processing protocol.

224

225 **Methods**

226 **Study design and participants**

227 For the present analyses we collated data from published studies identified through systematic reviews and
228 personal networks. Our updated literature search (eMethods in Supplement), along with the published
229 reviews,^{7,8} identified 21 articles from 13 studies that met the inclusion criteria: 1) cross-sectional study that
230 involved children aged 3.0 to 4.9 years, and 2) provided valid accelerometer-measured PA and
231 parent/caregiver-reported screen time and sleep duration data for at least 40 children per study site. Four
232 additional studies were identified through personal communications with the lead investigators, resulting in a
233 total of 17 studies eligible for inclusion (eTable 1 in Supplement). The sample size was selected based on the
234 observation that many studies from low- and lower-middle-income countries were pilot studies and, as such,
235 recruited small, non-representative samples. Therefore, having a larger sample size as the inclusion criteria
236 would limit the scope of our analysis to provide a more global examination of movement behaviours. For the
237 types of measures, we considered only studies that used ActiGraph or Actical accelerometers as they are
238 most frequently used in research and have been validated for measuring PA in young children.¹⁰ We selected
239 only studies that provided parent/caregiver reports of screen time and sleep duration, in line with the body of
240 evidence that guided the development of WHO guidelines.¹

241

242 We approached lead authors and invited them to share their dataset, including accelerometer data files,
243 parent/caregiver-reported screen time and sleep duration, and child's sex and age in months. Following
244 confirmation of data availability and the establishment of data-sharing agreements with respective
245 institutions, we obtained datasets for 14 studies before September 2022 (eTable 2 in Supplement). This study
246 was approved by the University of Wollongong Human Research Ethics Committee (2018/044). All datasets
247 used had prior ethical approval including approval for data sharing and/or obtained additional approval for
248 the purposes of this study. As the analyses used only deidentified data, no additional consent from

249 parents/caregivers was required, as determined by the University of Wollongong Human Research Ethics
250 Committee. The study followed the Strengthening the Reporting of Observational Studies in Epidemiology
251 (STROBE) reporting guideline.

252

253 **Measurements**

254 PA: Data were collected using ActiGraph (ActiGraph LLC, Pensacola, FL, USA) or Actical (Philips
255 Respironics Inc., Murrysville, PA, USA) accelerometers, worn on the right hip or non-dominant wrist,
256 during waking hours only or 24 hours/day for at least four days (eTable 2 in Supplement). Data were
257 harmonised by re-processing the accelerometer files using ActiLife software (version 6.12.1) (Figure 1). This
258 was not possible with data from the Canadian Health Measures Survey due to data-sharing policies. In this
259 case, we asked the collaborator to re-analyse the data using the same procedure. Data files were re-integrated
260 into 15-second epochs for the analysis of PA. Non-wear time was defined as 20 minutes of consecutive zero
261 counts.¹¹ Time spent in total PA (TPA) and moderate- to vigorous-intensity PA (MVPA) were calculated
262 using the best available, device- and wear-site-specific cut-points.¹²⁻¹⁵ PA data were only included if children
263 had at least three days of accelerometry data, with at least six hours of wear time per day¹⁶ between 5AM and
264 11PM. Valid weekend day data were not required for inclusion as it does not substantially increase the
265 reliability of PA estimates in this age group.¹⁶

266

267 Screen time: Questionnaire items typically asked parents/caregivers to report the total time their child spent
268 using electronic media for recreational purposes on a typical day (eTable 2 in Supplement). For studies
269 collecting data separately for weekdays and weekend days, the weighted average of screen time was used.

270

271 Sleep duration: Questionnaire items typically asked parents/caregivers to report either their child's total
272 hours of sleep per night/24-hour period (including naps) or typical sleep schedule (eTable 2 in Supplement).
273 For studies collecting data separately for weekdays and weekend days, the weighted average of sleep
274 duration was used.

275

276 **Statistical analysis**

277 Individual-level data were pooled to estimate the proportion of children who met the WHO

278 recommendations for PA, screen time and sleep duration (individually and in combination)¹ across countries,

279 according to the World Bank income classification¹⁷ at the time the data were collected, and by WHO region

280 (eTable 3 in Supplement).

281

282 Prior to the analyses, accelerometer variables were adjusted to address discrepancies in accelerometer

283 protocol (waking-hour only vs. 24-hour wear) across studies. As our analyses focused only on PA during

284 waking hours, a linear mixed model¹⁸ was fitted using R package “lme4” (with participants as random

285 intercepts and countries as fixed intercepts) based on day-level data to adjust children with 24-hour-

286 measured accelerometer variables as if they were measured with a waking-hour protocol using the residual

287 method. PA variables were further adjusted for accelerometer wear time using the residual method by fitting

288 a linear mixed model (MVPA and TPA as response variables, wear time as covariate and children as random

289 effects) and using the same average wear time of 662 minutes for all children in all countries. Similar

290 methods were applied to adjust sleep data to account for discrepancies in measures used (nocturnal sleep

291 duration vs. total sleep duration [including naps]).

292

293 Using the adjusted data, we conducted a meta-analysis with the R package “meta” (‘metamean’ function for

294 means and ‘metaprop’ for proportions using the inverse method and the summary measure ‘plogit’)¹⁹ to

295 derive aggregated country-level estimates and confidence intervals for each movement behaviour variable.

296 These estimates were used in a subsequent meta-analysis to obtain average estimates and confidence

297 intervals for each income group and region, overall and separately for boys and girls. Following this, a meta-

298 analysis of the pooled estimates across regions was conducted to derive overall estimates. It was not possible

299 to produce valid survey-based estimates due to the lack of population-level data for all countries included.

300 All analyses were conducted using R version 4.3.0.

301

302 **Results**

303 The analytical sample included 7,017 children (51.1% boys; mean age 4.1±0.5 years) from 33 countries
304 across six regions (4.1% Africa, 21.2% Americas, 3.1% Eastern Mediterranean, 31.8% Europe, 3.4% South-
305 East Asia and 36.4% Western Pacific). Two-thirds (78.1%) of the sample came from high-income countries,
306 followed by low-income and lower-middle-income (12.8%) and upper-middle-income countries (9.1%). The
307 descriptive characteristics of participants are reported in eTable 4 in Supplement.

308
309 The proportion of children who met the WHO guidelines is presented in Table 1 and visualised in Figure 2.
310 The overall proportion of children who met all three recommendations was 14.3% (95% CI, 9.7%-20.7%),
311 with a lower proportion in upper-middle-income countries than in low- and lower-middle-income and high-
312 income countries. Higher proportions meeting the guidelines were observed in Africa and Europe, while the
313 lowest proportion was found in the Americas. Similar patterns of sex differences were noted across income
314 groups (except for upper-middle-income countries) and regions, with higher proportions of boys meeting the
315 guidelines. Country-level data on guideline adherence is reported in eTable 5 in Supplement.

316
317 The proportion of children who met the sleep duration recommendation was nearly two times higher than
318 that of PA and screen time recommendations (Table 1). Similar patterns were noted across income groups
319 and regions (except for the Americas, where the proportion meeting the PA recommendation was the
320 highest). Concerning sex differences, the proportions meeting the PA recommendation were higher in boys
321 than in girls across income groups and regions. For both screen time and sleep duration recommendations,
322 however, higher adherence was observed in girls than in boys (except for Africa and the Americas, where
323 higher rates were reported in boys for sleep duration and screen time recommendation, respectively).

324
325 **Discussion**

326 This study is the first to report pooled data on the proportion of 3- to 4-year-old children meeting the WHO
327 global guidelines¹ from a large number of countries of varying incomes across six geographical regions.
328 Overall, 14.3% of children from 33 countries met all three recommendations, with the lowest proportion in

329 upper-middle-income countries and the Americas region. The proportion was generally higher among boys
330 than girls, which appeared due to sex differences in PA.

331

332 Early childhood is recognised as a critical window of opportunity for establishing healthy movement
333 behaviour patterns that are important for lifelong health and wellbeing.⁹ Our finding of a low proportion of
334 children meeting the overall WHO guidelines across countries and regions poses important implications for
335 future population health if no further actions are taken to address this issue. We also found that the low
336 proportion of children meeting the WHO guidelines was primarily driven by the low adherence to PA or
337 screen time recommendations, which varied by income groupings and regions. This suggests inequalities in
338 movement behaviours worldwide, consistent with the results of a multi-country study involving older
339 children.²⁰ This finding emphasises the urgent need to address the surveillance and research gaps among
340 underrepresented populations to strengthen the accountability of global health metrics, and inform the
341 development of more inclusive strategies to tackle movement behaviour inequalities. There is also a critical
342 need for contextually relevant and scalable interventions capable of achieving population-wide impacts while
343 reducing inequalities within and between countries.⁹

344

345 Our study reinforces the urgency of increasing PA participation in young children worldwide, as less than
346 half of our sample are meeting the PA recommendations. This concern is particularly evident for low- and
347 middle-income countries, possibly due to ongoing rapid urbanisation,²¹ which often results in less supportive
348 environments for PA. For example, the decrease in size and availability of green spaces/parks and pedestrian
349 access is evident due to the increased demand for commercial and residential areas and road infrastructure.²²
350 This, along with parental concerns about child safety from strangers and traffic, reduces opportunities for
351 children to play actively outside.²³ To increase PA participation will require a systems-based approach,²⁴
352 involving all relevant stakeholders working together to use their expertise and resources to make changes to
353 systems, environments, and policies.

354

355 Less than half of the children in our sample met the screen time recommendation. Notably, adherence was
356 particularly low among children from high-income countries and the Americas region, likely due to their

357 high mobile digital accessibility and ownership.²⁵ A recent review on parental perceptions of their children's
358 screen time found that most parents acknowledged screen time as a "necessity" in this technological era, and
359 they often used screen-based devices as a distraction (e.g., to keep their child occupied while they are busy),
360 for educational purposes, and as a reward for their children's behaviours.²⁶ Additionally, the review
361 highlighted that parents expressed difficulty in regulating their child's screen use and knowing how much
362 screen time children should have.²⁶ The changes in routines and social disruption due to the COVID-19
363 pandemic may have further increased children's exposure to prolonged screen time,²⁵ underscoring the
364 importance of better activating the guidelines to assist parents/caregivers in understanding healthy screen use
365 and in establishing boundaries for children's screen usage.

366

367 Compared to the PA and screen time recommendations, a higher proportion of children met the sleep
368 duration recommendation – a trend consistent across different income groups and regions, except in the
369 Americas where fewer children met the sleep recommendation compared to PA. This discrepancy may be
370 attributed to the widespread use of screen-based devices in this region, which has been shown to adversely
371 affect sleep outcomes in this age group.²⁷ It is important to note that this finding was based on parent-
372 reported measures, which tend to overestimate actual sleep duration compared to device-based measures.²⁸
373 Further, our study did not assess other aspects of sleep (quality and consistency) due to a lack of available
374 data. Future studies are recommended to explore how adherence to sleep recommendations varies with
375 different measurement methods and examine their associations with health outcomes. This will inform future
376 updates to sleep recommendations, which are currently based predominantly on evidence synthesised from
377 studies using only parent-reported measures.¹

378

379 We found that a higher proportion of boys than girls met all three recommendations. Specifically, a
380 consistently higher proportion of boys met the PA recommendation across all income groupings and regions.
381 This disparity is largely due to environmental and social factors. In some countries, boys often dominate play
382 spaces in public playgrounds.²⁹ Parents typically grant boys greater freedom, allowing them to play outdoors
383 more frequently with less supervision compared to girls.²³ In contrast, girls tend to receive less social support
384 and encouragement from their parents to participate in outdoor play.²³ PA is also less socially reinforced

among girls in many cultures, particularly in the African and Asian context, where girls are encouraged to spend more time indoors and engage in more static types of play (e.g, playing with toys) or activities that are more nurturing or domestic in nature (e.g., doing household chores). This reinforces the need for a holistic approach targeting social and cultural environments to reduce sex inequalities in PA.

389

A higher proportion of girls met the screen time recommendation than boys. The observed sex differences in screen time may be attributed to the digital divide, especially in low- and lower-middle-income countries where girls are reported to be less likely than boys to own or access digital technologies, even within the same households.³⁰ This may be related to traditional gender norms in certain cultural contexts, where girls are expected to contribute more to household chores or other routine domestic tasks, leaving them with less leisure time. Because the correlates of screen time are different for boys and girls in this age group,³¹ there may be a need for sex-specific strategies to manage young children's screen use.

397

Consistent with observations in child and adolescent populations,³² we found a higher proportion of girls that met the sleep duration recommendation. This may be linked to higher screen time among boys, resulting in later bedtimes and shorter sleep compared to girls.^{27,32} It is noted that multiple factors, including environmental (e.g., sleeping arrangement) and social-cultural contexts (e.g., bedtime routine), may influence a child's sleep duration differently.³³ This complexity makes it challenging to determine the primary factor contributing to the observed sex differences in sleep duration. Further cross-cultural studies assessing sleep characteristics and associated factors are needed to better understand the mechanisms underlying sex differences in early childhood sleep.

406

This study has several limitations. First, the dataset used covers only a small proportion of countries globally (~17%), and the sample sizes in most of the included studies were small and not representative of the preschool-aged population in each country. This calls for more large-scale, international studies that employ standardised and culturally appropriate measurement protocols to provide stronger evidence on the global prevalence of movement behaviours in this age group. Second, it is acknowledged that pooling accelerometry data collected using different devices and protocols, even when re-processed with a

standardised method, may have introduced biases into the PA estimates. The intensity cutpoints used, though tailored to specific devices and wear-sites, have notable limitations, such as being derived from small calibration studies and lacking robust measurement properties.³⁴ The best available cutpoints for the wrist-worn Actical accelerometer were based on validation studies in older children (aged 6-11 years),¹⁵ which may have introduced errors in determining PA for the younger children in this study. Additionally, the use of absolute intensity-based cut-points could potentially lead to the misclassification of activity behaviours.³⁵ Nevertheless, we chose this approach to align with past literature from which the evidence guiding the development of the PA guidelines was derived. Similarly, variations in questions used to assess screen time and sleep duration may have led to varying estimates across studies, subsequently influencing our pooled estimates. More importantly, the evidence supporting the use of the existing measures of movement behaviours is largely based on studies conducted in the Western or high-income countries, with limited evidence on their cross-cultural validity.^{10,36} Therefore, our estimates may not accurately reflect the true disparities between countries or regions. Finally, our data were mostly collected more than five years ago and prior to the COVID-19 pandemic. As such, our estimates of movement behaviours may not be generalisable to contemporary young children.

428

429 **Conclusions**

We found that in a large multi-country sample of children, less than one in six met the overall WHO Guidelines. While differences were noted between boys and girls and among regions and income settings, the proportion meeting guidelines was universally low. The WHO recommends a systems-based approach to promote healthy levels of movement behaviours across all ages. At the country level, stakeholders from all sectors must work collaboratively to create an active society by changing social norms and attitudes and by providing places and spaces that support children to move more, be less sedentary, and have adequate sleep. These actions must not leave any country behind to ensure that current gaps in the evidence base are addressed equitably. In many low- and middle-income countries, movement behaviours need to be better anchored to other more salient outcomes, such as school readiness, and framed in the context of other priorities such as food insecurity and undernutrition. Finally, robust surveillance processes are essential for monitoring temporal changes and assessing interventions intended to elicit improvements.

441

442 **Acknowledgements**

443 Dr Kar Hau Chong and Professor Anthony D. Okely had full access to all the data in the study and take
444 responsibility for the integrity of the data and the accuracy of the data analysis.

445 Concept and design: Okely, Reilly, Tremblay, Chong, Suesse.

446 Acquisition, analysis, or interpretation of data: All authors.

447 Drafting of the manuscript: Chong, Okely, Suesse.

448 Critical review of the manuscript for important intellectual content: All authors.

449 Statistical analysis: Suesse, Chong.

450 Administrative, technical, or material support: All authors.

451 Supervision: Okely.

452 **Non-author contributions**

453 Cecilia H. S. Chan, PhD, Department of Sports Science and Physical Education, Kwok Sports Building, The
454 Chinese University of Hong Kong, Shatin, Hong Kong, China; Contribution: Data acquisition.

455 Fazlollah Ghofranipour, PhD, Department of Health Education and Health Promotion, Faculty of Medical
456 Science, Tarbiat Modares University, Tehran, Iran; Contribution: Data acquisition.

457 Adang Suherman, PhD, Faculty of Sport and Health Education, Universitas Pendidikan Indonesia, Bandung,
458 Jawa Barat, Indonesia; Contribution: Data acquisition.

459 **Conflict of Interest Disclosures:** Dr. Christian reports grants from Telethon Kids Institute, University of
460 Western Australia Western Australian Health Promotion Foundation (Healthway), grants from Telethon Kids
461 Institute, University of Western Australia Australian National Health & Medical Research Council
462 (NHMRC), and grants from Telethon Kids Institute, University of Western Australia Australian Research
463 Council (ARC) during the conduct of the study; grants from National Heart Foundation Future Leader
464 Fellowship outside the submitted work. Dr. Cliff reports grants from Australian Research Council during the
465 conduct of the study. Dr. De Craemer reports grants from Seventh Framework Programme (CORDIS FP7)
466 during the conduct of the study. Dr. El Hamdouchi reports grants from ISBNPA Pioneers grant. Dr.
467 Kontsevaya reports grants from WHO during the conduct of the study. Dr. Meredith-Jones reports grants
468 from Health Research Council New Zealand 08/374, grants from Health Research Council New Zealand

469 12/281, and grants from Health Research Council New Zealand 12/310 during the conduct of the study. Dr.
470 Suherman reports grants from Universitas Pendidikan Indonesia during the conduct of the study. Dr. Tanaka
471 reports grants from Sasakawa Sports Foundation Research Grant (grant number 190A2- 004) during the
472 conduct of the study. Dr. Taylor reports grants from Health Research Council of New Zealand during the
473 conduct of the study. Dr. Tremblay reports grants from Canadian Institutes of Health Research during the
474 conduct of the study.

475 **Funding/Support:** No funding was received for conducting this pooled analysis. The sources of funding
476 received for the conduct of each included study were as follows: ACTNOW (Research Council of Norway
477 Grant [grant number 287903]; the County Governor of Sogn og Fjordane; the Sparebanken Sogn og
478 Fjordane Foundation; Western Norway University of Applied Sciences); CHL (Agriculture and Food
479 Research Initiative Grant [grant number 2011-68001-30335]); CHMS (Canadian Research Data Centre
480 Network from the Social Sciences and Humanities Research Council; Canadian Institutes of Health
481 Research; Canadian Foundation for Innovation; Statistics Canada); DAGIS (Folkhälsan Research Center,
482 University of Helsinki; Ministry of Education and Culture in Finland; Ministry of Social Affairs and Health;
483 The Academy of Finland [grant number 285439, 287288, 288038, 350820]; The Juho Vainio Foundation;
484 The Signe and Ane Gyllenberg Foundation; The Finnish Cultural Foundation/South Ostrobothnia Regional
485 Fund; The Päivikki and Sakari Sohlberg Foundation; Medicinska Föreningen Liv och Hälsa, Finnish
486 Foundation for Nutrition Research; Finnish Food Research Foundation); HAPPY (Deakin University);
487 PACE (Palix Foundation; Alberta Family Wellness Initiative; PolicyWise for Children & Families; Women
488 and Children's Health Research Institute); PATH-ABC (Australian Research Council Discovery Early Career
489 Award [grant number DE230101174]); PLAYCE (Western Australian Health Promotion Foundation
490 (Healthway) [grant number 24219, 32925]; UKRI-NHMRC Built Environment Prevention Research Scheme
491 [grant number GNT1192764, MR/T039329/1]; Australian Research Council Centre of Excellence for
492 Children and Families over the Life Course [grant number CE200100025]); POI (Health Research Council
493 of New Zealand [grant number 08/374, 12/281, 12/310]; the Southern District Health Board); Parent-Child
494 Movement Behaviours and Pre-School Children's Development Study (Faculty of Kinesiology, Sport, &
495 Recreation at the University of Alberta); SUNRISE (National Health and Medical Research Council
496 Australia [grant number APP1176858]; Biomedical Research Foundation in Bangladesh [grant number

497 BRF/IRG2019/01]; Brazilian National Council for Scientific and Technological Development (CNPq) [grant
498 number 309301/2020-3]); Canadian Institutes of Health Research Planning and Dissemination (grant number
499 392396); The Talent Construction Plan of High Level Health Professional in Beijing Health System [grant
500 number 2015-3-008]); KEM Hospital Research Centre [grant number 1924]; Universitas Pendidikan
501 Indonesia Research Grant [grant number 293/UN40-D/PT/2019]; Sasakawa Sports Foundation Research
502 Grant (grant number 190A2-004); Sir Halley Stewart Trust [grant number 2674]; Universiti Kebangsaan
503 Malaysia Research University Grant [grant number GUP-2018-142]; International Society of Behavioral
504 Nutrition and Physical Activity Pioneers Program 2021; Comprehensive Health and Epidemiological
505 Surveillance System 2018-2022, funded by the Government of Papua New Guinea through the Department
506 of National Planning and Monitoring [grant number 02704]; University of the Witwatersrand, Faculty of
507 Health Sciences; Dr Stella de Silva Research Grant at the Sri Lanka College of Paediatricians; Region
508 Östergötland [grant number LIO-936921]; Pham Ngoc Thach University of Medicine's Fund for Science
509 [grant number 1320/HĐ-TĐHYKPNT]); The Movement's Cool Project (Brazilian Federal Foundation for
510 Support and Evaluation of Graduate Education [grant number CAPES-PRINT—88887.369625/2019-00]);
511 ToyBox-Study (Seventh Framework Programme of the European Commission [grant number 245200]). The
512 authors would like to thank all participating children, families, research staff and organisations for their
513 invaluable involvement in the respective studies.

References

1. World Health Organization. Guidelines on physical activity, sedentary behaviour and sleep for children under 5 years of age. <https://www.who.int/publications/i/item/9789241550536>. Accessed February 23, 2023.
2. Tremblay MS. Introducing 24-Hour Movement Guidelines for the Early Years: A New Paradigm Gaining Momentum. *J Phys Act Health*. 2020;17(1):92-95. doi: 10.1123/jpah.2019-0401.
3. Leppänen MH, Ray C, Wennman H, et al. Compliance with the 24-h movement guidelines and the relationship with anthropometry in Finnish preschoolers: the DAGIS study. *BMC Public Health*. 2019;19(1):1618. doi: 10.1186/s12889-019-7967-7.
4. Feng J, Zheng C, Sit CH, Reilly JJ, Huang WY. Associations between meeting 24-hour movement guidelines and health in the early years: A systematic review and meta-analysis. *J Sport Health Sci*. 2021;39(22):2545-2557. doi: 10.1080/02640414.2021.1945183.
5. Kracht CL, Webster EK, Staiano AE. Relationship between the 24-Hour Movement Guidelines and fundamental motor skills in preschoolers. *J Sci Med Sport*. 2020;23(12):1185-1190. doi: 10.1016/j.jsams.2020.06.021.
6. Byambaa A, Dechinjamts O, Jambaldorj B, Jones RA, Chong KH, Okely AD. Prevalence and Health Associations of Meeting the World Health Organization Guidelines for Physical Activity, Sedentary Behavior, and Sleep in Preschool-Aged Children: The SUNRISE Mongolia Pilot and Feasibility Study. *J Phys Act Health*. 2024;21(3):283-293. doi: 10.1123/jpah.2023-0511.
7. Rollo S, Antsygina O, Tremblay MS. The whole day matters: Understanding 24-hour movement guideline adherence and relationships with health indicators across the lifespan. *J Sport Health Sci*. 2020;9(6):493-510. doi: 10.1016/j.jshs.2020.07.004.
8. Tapia-Serrano MA, Sevil-Serrano J, Sánchez-Miguel PA, López-Gil JF, Tremblay MS, García-Hermoso A. Prevalence of meeting 24-Hour Movement Guidelines from pre-school to adolescence: A systematic review and meta-analysis including 387,437 participants and 23 countries. *J Sport Health Sci*. 2022;11(4):427-437. doi: 10.1016/j.jshs.2022.01.005.
9. Carson V, Draper CE, Okely A, Reilly JJ, Tremblay MS. Future Directions for Movement Behavior Research in the Early Years. *J Phys Act Health*. 2023;21(3):218-221. doi: 10.1123/jpah.2023-0679.

10. Phillips SM, Summerbell C, Hobbs M, et al. A systematic review of the validity, reliability, and feasibility of measurement tools used to assess the physical activity and sedentary behaviour of pre-school aged children. *Int J Behav Nutr Phys Act*. 2021;18(1):141. doi: 10.1186/s12966-021-01132-9.
11. Esliger DW, Copeland JL, Barnes JD, Tremblay MS. Standardizing and optimizing the use of accelerometer data for free-living physical activity monitoring. *J Phys Act Health*. 2005;2(3):366-383. doi: 10.1123/jpah.2.3.366.
12. Pate RR, Almeida MJ, McIver KL, Pfeiffer KA, Dowda M. Validation and calibration of an accelerometer in preschool children. *Obesity (Silver Spring)*. 2006;14(11):2000-2006. doi: 10.1038/oby.2006.234.
13. Pate RR, O'Neill JR, Brown WH, Pfeiffer KA, Dowda M, Addy CL. Prevalence of Compliance with a New Physical Activity Guideline for Preschool-Age Children. *Child Obes*. 2015;11(4):415-420. doi: 10.1089/chi.2014.0143.
14. Adolph AL, Puyau MR, Vohra FA, Nicklas TA, Zakeri IF, Butte NF. Validation of uniaxial and triaxial accelerometers for the assessment of physical activity in preschool children. *J Phys Act Health*. 2012;9(7):944-953. doi: 10.1123/jpah.9.7.944.
15. Schaefer CA, Nace H, Browning R. Establishing wrist-based cutpoints for the Actical accelerometer in elementary school-aged children. *J Phys Act Health*. 2014;11(3):604-613. doi: 10.1123/jpah.2011-0411.
16. Bingham DD, Costa S, Clemes SA, Routen AC, Moore HJ, Barber SE. Accelerometer data requirements for reliable estimation of habitual physical activity and sedentary time of children during the early years - a worked example following a stepped approach. *J Sports Sci*. 2016;34(20):2005-2010. doi: 10.1080/02640414.2016.1149605.
17. The World Bank. World Bank Country and Lending Groups – Country Classification. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519>. Accessed February 21, 2022.
18. Hox J, Moerbeek M, van de Schoot R, eds. *Multilevel Analysis: Techniques and Applications*. New York, NY: Routledge; 2017.
19. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. *Evid Based Ment Health*. 2019;22(4):153-160. doi: 10.1136/ebmental-2019-300117.

20. Chaput JP, Barnes JD, Tremblay MS, Fogelholm M, Hu G, Lambert EV, et al. Inequality in physical activity, sedentary behaviour, sleep duration and risk of obesity in children: a 12-country study. *Obes Sci Pract.* 2018;4(3):229-237. doi: 10.1002/osp4.271.
21. United Nations. World Urbanization Prospects: The 2018 Revision. <https://population.un.org/wup/Publications/Files/WUP2018-Report.pdf>. Accessed November 12, 2023.
22. Statistics Canada. Urban greenness, 2001, 2011 and 2019. <https://www150.statcan.gc.ca/n1/pub/16-002-x/2021001/article/00002-eng.htm>. Accessed November 12, 2023.
23. Lee EY, Bains A, Hunter S, Ament A, Brazo-Sayavera J, Carson V, et al. Systematic review of the correlates of outdoor play and time among children aged 3-12 years. *Int J Behav Nutr Phys Act.* 2021;18(1):41. doi: 10.1186/s12966-021-01097-9.
24. World Health Organization. Global action plan on physical activity 2018-2030: more active people for a healthier world. <https://www.who.int/publications/i/item/9789241514187>. Accessed November 10, 2023.
25. Ribner AD, Coulanges L, Friedman S, Libertus ME; I-FAM-Covid Consortium. Screen Time in the Coronavirus 2019 Era: International Trends of Increasing Use Among 3- to 7-Year-Old Children. *J Pediatr.* 2021;239:59-66. doi: 10.1016/j.jpeds.2021.08.068.
26. Chong SC, Teo WZ, Shorey S. Exploring the perception of parents on children's screentime: a systematic review and meta-synthesis of qualitative studies. *Pediatr Res.* 2023;94(3):915-925. doi: 10.1038/s41390-023-02555-9.
27. Janssen X, Martin A, Hughes AR, Hill CM, Kotronoulas G, Hesketh KR. Associations of screen time, sedentary time and physical activity with sleep in under 5s: A systematic review and meta-analysis. *Sleep Med Rev.* 2020;49:101226. doi: 10.1016/j.smr.2019.101226.
28. Perpétuo C, Fernandes M, Veríssimo M. Comparison Between Actigraphy Records and Parental Reports of Child's Sleep. *Front Pediatr.* 2020;8:567390. doi: 10.3389/fped.2020.567390.
29. Reimers AK, Schoeppe S, Demetriou Y, Knapp G. Physical Activity and Outdoor Play of Children in Public Playgrounds-Do Gender and Social Environment Matter? *Int J Environ Res Public Health.* 2018;15(7):1356. doi: 10.3390/ijerph15071356.

30. UNICEF. Bridging the Gender Digital Divide: Challenges and an Urgent Call for Action for Equitable Digital Skills Development. <https://data.unicef.org/resources/ictgenderdivide/>. Accessed November 29, 2023.
31. Downing KL, Hinkley T, Salmon J, Hnatiuk JA, Hesketh KD. Do the correlates of screen time and sedentary time differ in preschool children? *BMC Public Health*. 2017;17(1):285. doi: 10.1186/s12889-017-4195-x.
32. Olds T, Blunden S, Petkov J, Forchino F. The relationships between sex, age, geography and time in bed in adolescents: a meta-analysis of data from 23 countries. *Sleep Med Rev*. 2010;14(6):371-378. doi: 10.1016/j.smrv.2009.12.002.
33. Zhang Z, Sousa-Sá E, Pereira JR, Okely AD, Feng X, Santos R. Correlates of Sleep Duration in Early Childhood: A Systematic Review. *Behav Sleep Med*. 2021;19(3):407-425. doi: 10.1080/15402002.2020.1772264.
34. Lettink A, Altenburg TM, Arts J, van Hees VT, Chinapaw MJM. Systematic review of accelerometer-based methods for 24-h physical behavior assessment in young children (0-5 years old). *Int J Behav Nutr Phys Act*. 2022;19(1):116. doi: 10.1186/s12966-022-01296-y.
35. Trost SG. Population-level physical activity surveillance in young people: are accelerometer-based measures ready for prime time? *Int J Behav Nutr Phys Act*. 2020;17(1):28. doi: 10.1186/s12966-020-00929-4.
36. Arts J, Gubbels JS, Verhoeff AP, Chinapaw MJM, Lettink A, Altenburg TM. A systematic review of proxy-report questionnaires assessing physical activity, sedentary behavior and/or sleep in young children (aged 0-5 years). *Int J Behav Nutr Phys Act*. 2022;19(1):18. doi: 10.1186/s12966-022-01251-x.

Table 1. Proportion (95% CI) of children meeting the World Health Organization guidelines for physical activity, sedentary behaviour (screen time), and sleep duration


| | Physical activity ^a | | | Sedentary behaviour (Screen time) ^b | | | Sleep duration ^c | | | All three recommendations | | |
|---|--------------------------------|---------------------|---------------------|---|---------------------|---------------------|-----------------------------|---------------------|---------------------|---------------------------|---------------------|---------------------|
| | Boys | Girls | Total | Boys | Girls | Total | Boys | Girls | Total | Boys | Girls | Total |
| Global Estimate (n=7,017) | 55.6 (42.3,68.1) | 41.7 (29.3,55.1) | 49.2 (36.1,62.3) | 38.2 (27.3,50.4) | 45.1 (30.7,60.5) | 41.8 (29.0,55.9) | 78.8 (67.9,86.7) | 82.3 (71,89.8) | 81.0 (69.7,88.8) | 16.8 (12.0,22.9) | 12.8 (8.3,19.2) | 14.3 (9.7,20.7) |
| World Bank income groups | | | | | | | | | | | | |
| Low/Lower-middle income countries (n=900) | 47.2 (35.8,58.9) | 35.3 (21.9,51.5) | 43.3 (29.7,58.0) | 47.2 (33.7,61) | 53.3 (37.4,68.5) | 49.9 (35.9,64.0) | 76.3 (68.7,82.5) | 77.8 (70.3,83.8) | 77.9 (70.7,83.8) | 20.6 (13.6,30.1) | 14.9 (8.9,24.0) | 16.6 (10.4,25.3) |
| Upper-middle income countries (n=641) | 44.7 (23.9,67.5) | 36.6 (11.8,71.2) | 43.7 (16.2,75.7) | 45.2 (25.0,67.1) | 53.0 (29.8,75.0) | 50.5 (27.1,73.7) | 67.2 (47.9,82.1) | 72.1 (55.5,84.2) | 71.0 (52.4,84.5) | 12.3 (5.8,24.1) | 12.3 (6.0,23.6) | 11.9 (5.9,22.5) |
| High-income countries (n=5,476) | 71.1 (52.7,84.5) | 56.7 (36.4,75.0) | 65.4 (45.4,81.2) | 31.0 (22.3,41.3) | 32.1 (23.9,41.7) | 31.3 (22.8,41.2) | 84.7 (73.6,91.7) | 87.8 (76.9,94.0) | 86.7 (75.6,93.1) | 16.4 (10.9,23.9) | 12.5 (7.7,19.7) | 14.4 (9.6,21.1) |
| WHO Regions | | | | | | | | | | | | |
| Africa (n=286) | 52.0 (43.2,60.6) | 49.2 (34.2,64.4) | 50.1 (38.6,61.5) | 59.4 (26.7,85.4) | 64.8 (33.3,87.1) | 62.9 (31.6,86.2) | 81.5 (50.4,95) | 78.2 (57.1,90.6) | 80.3 (55.2,93.1) | 24.8 (12.0,44.3) | 24.7 (12.9,42.2) | 23.9 (11.6,43) |
| Americas (n=1,487) | 72.8 (27.2,95.0) | 61.3 (14.0,93.9) | 67.6 (20.4,94.4) | 17.5 (9.8,29.3) | 17.1 (8.4,31.8) | 17.0 (9.1,29.7) | 62.8 (40.5,80.8) | 64.4 (42.2,81.7) | 63.5 (41.5,81.0) | 9.2 (5.2,15.8) | 7.0 (2.4,18.3) | 7.7 (3.6,15.8) |
| Eastern Mediterranean (n=219) | 41.2 (25.6,58.8) | 28.8 (10.3,58.8) | 36.2 (19.0,57.8) | 39.7 (16.0,69.5) | 60.3 (32.5,82.7) | 47.7 (22.8,73.8) | 74.9 (53.0,88.7) | 84.0 (73.9,90.8) | 80.1 (61.8,91.0) | 18.5 (6.7,41.8) | 14.3 (4.2,39.2) | 15.8 (5.2,38.9) |
| Europe (n=2,232) | 62.8 (46.7,76.5) | 44.1 (33.0,55.9) | 53.5 (40.4,66.1) | 43.6 (29.9,58.3) | 48.6 (29.3,68.4) | 50.0 (27.5,72.5) | 93.1 (87.4,96.3) | 95.1 (89.7,97.7) | 94.7 (89.1,97.5) | 26.4 (17.5,37.8) | 19.8 (11.8,31.4) | 23.5 (14.8,35.1) |
| South-East Asia (n=240) | 33.1 (15.5,57.2) | 20.2 (8.5,40.9) | 26.0 (12.3,46.9) | 35.6 (24.5,48.4) | 46.8 (21.3,74.1) | 40.6 (22.1,62.2) | 75.3 (66.6,82.3) | 83.2 (75.2,89.1) | 79.7 (74.0,84.3) | 13.4 (7.6,22.3) | 7.3 (3.1,16.2) | 9.1 (4.8,16.6) |
| Western Pacific (n=2,553) | 69.4 (42.5,87.4) | 51.5 (24.9,77.3) | 62.9 (35.3,84.1) | 39.2 (27.6,52.2) | 39.3 (27.9,52.1) | 39.0 (27.9,51.5) | 74.1 (61.4,83.8) | 75.2 (62.6,84.6) | 75.0 (62.5,84.4) | 13.7 (7.9,22.7) | 11.0 (6.4,18.3) | 12.4 (7.7,19.3) |

^aDefined as meeting both the total physical activity (≥ 180 mins/day) and moderate- to vigorous-intensity (≥ 60 mins/day) recommendations.

^bNot more than one hour of screen time per day.


^c10 to 13 hours of sleep per day.

Figure 1. Accelerometry data processing and analysis procedures.



Abbreviation: MVPA moderate- to vigorous-intensity physical activity, TPA total physical activity

Figure 2. Proportion of children meeting World Health Organization guidelines, overall and by income group and region.



Note. Data presented as average estimates with 95% confidence intervals.