

This is the peer reviewed version of the following article: Mesa-Rodriguez A, Hernández-Cruz CM, Betancor MB, Fernández-Palacios H, Izquierdo MS, Roo J. Effect of increasing docosahexaenoic acid content in weaning diets on survival, growth and skeletal anomalies of longfin yellowtail (*Seriola rivoliana*, Valenciennes 1833). *Aquaculture Research* 2018;49:1200–1209, which has been published in final form at <https://doi.org/10.1111/are.13573>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

1       **Effect of increasing DHA content in weaning diets on survival,**  
2       **growth and skeletal anomalies of longfin yellowtail (*Seriola rivoliana*,**  
3                               **valenciennes 1833).**

4       A. Mesa-Rodríguez<sup>1\*</sup>, C.M. Hernández-Cruz<sup>1</sup>, M.B. Betancor<sup>2</sup>, H.  
5       Fernández-Palacios<sup>1</sup>, M.S. Izquierdo<sup>1</sup> and J. Roo<sup>1</sup>.

6

7       <sup>1</sup>*Grupo de Investigación en Acuicultura, Parque Científico Tecnológico*  
8       *Marino, Universidad de Las Palmas de Gran Canaria, Taliarte, 35200*  
9       *Telde, Canary Islands, Spain*

10      <sup>2</sup>*Institute of Aquaculture, Faculty of Natural Sciences, University of*  
11      *Stirling, Stirling FK9 4LA, United Kingdom*

12

13      \*Author to whom correspondence should be addressed.

14      Antonio Mesa Rodríguez

15      Tel.: +34928132900; Fax: +34928132908;

16

17      e-mail: [a.mesa.rodriguez@gmail.com](mailto:a.mesa.rodriguez@gmail.com); web page: [www.giaqua.org](http://www.giaqua.org)

## 18   **Abstract**

19   Five isoproteic (54.8%) and isolipidic (24.1%) microdiets, which varied  
20   in their docosahexaenoic acid (DHA) content (0.25, 0.75, 1.64, 1.99 and  
21   3.17%; dw), were manufactured to determine its effects on longfin  
22   yellowtail *Seriola rivoliana* larvae in terms of fish biological  
23   performance, whole body fatty acid profile and incidence of skeletal  
24   anomalies from 30 dah ( $11.31 \pm 1.79$  Total Length, TL) to 50 dah  
25   ( $19.80 \pm 0.58$  mm TL). The inclusion of dietary DHA up to 3.17% (dw)  
26   improved larval resistance to air exposure, although DHA did not  
27   significantly affect fish final growth or final survival. Indeed, high levels  
28   of dietary DHA (1.99% and 3.17%, dw) tended to increase the incidence  
29   of skeletal anomalies in *S. rivoliana* larvae, albeit no significant  
30   differences were observed. Furthermore, the occurrence of severe  
31   anomalies such as kyphosis and lordosis, was mainly associated to the  
32   larvae fed with the highest levels of dietary DHA. In terms of survival,  
33   increasing dietary DHA levels did not significantly affect longfin  
34   yellowtail survival rate, despite a tendency for enhanced survival. The  
35   results of the present study proved that the inclusion of dietary DHA in  
36   inert diets up to a 3.17% (dw) and a DHA/EPA ratio above 3.1 increased  
37   the final survival and stress resistance in *S. rivoliana* larvae.

38 **Keywords:** longfin yellowtail, fish larvae, docosahexaenoic acid,  
39 microdiets, skeletal anomalies.

## 40 **1. Introduction**

41 The recent interest on marine fast-growing teleost for aquaculture  
42 diversification has lead to research in fish species such as Atlantic bluefin  
43 tuna (*Thunnus thynnus*), greater amberjack (*Seriola dumerili*), yellowtail  
44 kingfish (*Seriola lalandi*), Japanese yellowtail (*Seriola quinqueradiata*) or  
45 meagre (*Argyrosomus regius*). Longfin yellowtail, (*Seriola rivoliana*,  
46 Valenciennes 1833) is a carangid with a high commercial interest due to  
47 its fast growth rate and worldwide distribution (Roo *et al.*, 2012; Mesa-  
48 Rodriguez *et al.*, 2014; Mesa-Rodriguez *et al.*, 2016). Moreover, *S.*  
49 *revoliana* is already commercially produced in Hawaii (Sims & Key,  
50 2011) and under pilot scale experimental production in Gran Canaria  
51 (Canary Islands; Spain) from 2010 (GIA, 2011).  
52 Nonetheless, very few studies have been performed in order to determine  
53 *S. rivoliana* nutritional requirements (Roo *et al.*, 2012; Fernández-  
54 Palacios, Schuchardt, Roo, Hernández-Cruz & Izquierdo, 2015). In this  
55 sense, several studies have been reported for other species from the same  
56 genus, such as *Seriola dumerili* (Garcia-Gomez, 2000; Tomas, de la  
57 Gandara, Garcia-Gomez, Perez & Jover, 2005; Takakuwa, Fukada,

58 Hosokawa & Masumoto, 2006; Papadakis, Chatzifotis, Divanach &  
59 Kentouri, 2007; Hamasaki, Tsuruoka, Teruya, Hashimoto & Hamada,  
60 2009; Matsunari *et al.*, 2012; Matsunari *et al.*, 2013), *Seriola lalandi*  
61 (Cobcroft, Pankhurst, Poortenaar & Tait, 2004) and *Seriola*  
62 *quinqueradiata* (Masuda *et al.*, 1998; Ishizaki *et al.*, 2001; Yamamoto *et*  
63 *al.*, 2008; Takeuchi, 2014).

64 Among the nutrients, long chain polyunsaturated fatty acids (LC-PUFAs)  
65 are determinant for the success of larvae rearing (Izquierdo, 2005).  
66 Moreover, the adequate culture performance of marine fish larvae is  
67 related to the inclusion of the omega 3 (n-3) LC-PUFA docosahexaenoic  
68 acid (DHA; 22:6n-3) in the diet, due to its direct relationship with tissues  
69 and cell functioning (Izquierdo & Koven, 2011). Not only DHA is an  
70 essential fatty acid (EFA) for larval rearing success, but also the  
71 importance of other n-3 LC-PUFA (eicosapentaenoic acid; EPA; 20:5n-3)  
72 as well as n-6 LC-PUFA (arachidonic acid; ARA; 20:4n-6) has been  
73 emphasized (Izquierdo, 1996). Besides, several studies indicated that  
74 DHA had a greater potential than EPA as an EFA for marine fish larvae  
75 (Watanabe, Izquierdo, Takeuchi, Satoh & Kitajima 1989; Takeuchi, 2001;  
76 Izquierdo & Koven, 2011), being the DHA requirements more limiting  
77 for growth, survival (Izquierdo, 1996) and development of schooling

78 behaviour (Masuda *et al.*, 1998; Ishizaki *et al.*, 2001) than EPA.  
79 Contrarily, some studies observed that high levels of dietary DHA may  
80 cause muscular dystrophy (Betancor *et al.*, 2011) or lead to the  
81 appearance of supernumerary vertebrae (Villeneuve, Gisbert, Moriceau,  
82 Cahu & Zambonino-Infante, 2006) in *Dicentrarchus labrax* larvae due to  
83 the peroxidation of DHA and the formation of toxic oxidized compounds.  
84 On the other hand, the effects of dietary DHA deficiency have been  
85 reported in a variety of marine fish species, being characterized by an  
86 increase in the incidence of skeletal deformities in larvae of *Sparus aurata*  
87 (Roo, Hernandez-Cruz, Socorro, Fernandez-Palacios & Izquierdo, 2010;  
88 Izquierdo *et al.*, 2013) and *Pagrus pagrus* (Roo *et al.*, 2009; Izquierdo,  
89 Socorro & Roo, 2010), as well as jaw anomalies in *Latris lineata*  
90 (Cobcroft, Pankhurst, Sadler & Hart, 2001). Additionally, the deficiency  
91 of DHA could lead to alteration in gut and liver in *Latris lineata*  
92 (Bransden, Battaglione, Morehead, Dunstan & Nichols, 2005), or to  
93 malpigmentation and irregular eye migration in flatfish (Bell, McEvoy,  
94 Estévez, Shields & Sargent, 2003) as well as reduced stress resistance in  
95 *Huso huso* (Jalali, Hosseini & Imanpour, 2008).  
96 Apart from all the negatives effects caused by inadequate dietary DHA  
97 levels in larval feeds previously described, the low culture performance

98 and survival has been identified as the main issue in different larval  
99 species (Watanabe, Izquierdo, Takeuchi, Satoh & Kitajima, 1989; Furuita,  
100 Takeuchi, Toyota & Watanabe, 1996a; Furuita *et al.*, 1996b; Copeman,  
101 Parrish, Brown & Harel, 2002; Rezek, Watanabe, Harel & Seaton, 2010).  
102 Due to the relevance of DHA as a main dietary fatty acid for larval marine  
103 finfish rearing success, the purpose of this study was to evaluate the effect  
104 of increasing dietary DHA levels on growth performance and larval  
105 quality of *S. rivoliana* with the intention to elucidate the adequate dietary  
106 DHA level for this species. In order to do so, five feeds containing  
107 increasing levels of DHA were fed to longfin yellowtail larvae from 30 to  
108 50 dah and larvae growth, final survival, survival after activity test, larvae  
109 fatty acid profile and incidence of skeletal anomalies evaluated.

## 110 **2. Materials and methods**

### 111 ***2.1 Experimental diets***

112 Five isoproteic and isolipidic diets were formulated to contain increasing  
113 DHA contents (Table 1). DHA, EPA (DHA-50 and EPA-50, Croda  
114 Chemicals Ltd. Goole, U.K.) and ARA (Vevodar DSM Food Specialities,  
115 Netherlands) oils were added in graded amounts in substitution of oleic  
116 acid to maintain a constant lipid content (~ 20%; Table 1). Diets were  
117 named according to their analysed DHA content (dw) as follows: DHA0

118 (0.25% DHA); DHA1 (0.75% DHA); DHA1.5 (1.64% DHA); DHA2  
119 (1.99% DHA) and DHA3 (3.17% DHA). Microdiets were manufactured  
120 according to Betancor *et al.*, 2012a,b by mixing squid meal and water-  
121 soluble components, then the lipid and fat soluble vitamins and, finally,  
122 gelatin dissolved in warm water. The paste was compressed pelleted  
123 (Severin, Suderm, Germany) and dried in an oven (Ako, Barcelona,  
124 Spain) at 38 °C for 24 h. Pellets were ground (Braun, Kronberg,  
125 Germany) and sieved (Filtru, Barcelona, Spain) to obtain two particle  
126 sizes, from 250 to 500 µm and from 500-710 µm. Formulated diets were  
127 analysed for proximate and fatty acid composition.

128

## 129 ***2.2 Broodstock and larval rearing***

130 *S. rivoliana* eggs were obtained from induced spawning of fifteen wild  
131 adults ( $1.76 \pm 0.25$  kg) adapted to captivity at GIA (Grupo de  
132 Investigación en Acuicultura) facilities 10 m<sup>3</sup> squared glass fiber tanks in  
133 land. Gonadotropin releasing hormone analogue (LHRHa, des-Gly10, [D-  
134 Ala6]; Sigma- Aldrich, St. Louis, MO, USA) was used at a dose of 20 µg  
135 kg<sup>-1</sup> body weight, based on the reported dosage for longfin yellowtail  
136 (Roo *et al.*, 2012). Larvae were reared under mesocosms rearing system



137 following the methodology described by Roo *et al.* (2012). In this way,  
138 4.5 eggs l<sup>-1</sup> were stocked in two 40 m<sup>3</sup> tanks up to 29 days after hatching  
139 (dah). At 30 dah (11.31 ± 1.79 total length, TL; 11.72 ± 0.97 mg), larvae  
140 were settled in 200 l fibreglass cylinder tanks with conical bottom and  
141 painted a light grey colour (90 larvae per tank, in triplicates). Filtered  
142 seawater was supplied (37 g l<sup>-1</sup> salinity) and water conditions were daily  
143 measured (temperature: 22.5 ± 0.6 °C; oxygen levels: 6.5 ± 0.3 g l<sup>-1</sup>;  
144 OxyGuard, Denmark). Photoperiod was kept at 12:12 (12 h light:12 h  
145 dark) by fluorescent daylights at 1700 lux (digital Lux Tester YF-1065;  
146 Powertech Rentals, Osborne, Australia).

### 147 **2.3 Growth, survival and activity test**

148 Larval growth was assessed by estimating the TL of the larvae using a  
149 profile projector (Nikon V-12A, NIKON™, Tokyo, Japan) at 30, 42 and  
150 50 dah. Final larvae survival was calculated by individually counting the  
151 larvae at the beginning and at the end of the trial. Additionally, an activity  
152 test was performed by subjecting fifteen larvae per tank to 30 seconds of  
153 air exposure at 42 and 50 dah and counting all the remaining alive larvae  
154 after 24 h as previously described (Izquierdo, Watanabe, Takeuchi,  
155 Arakawa & Kitajima, 1989).

#### 156 **2.4 Biochemical analyses of diets and larvae**

157 A sample of 50 dah larvae from each tank was washed with distilled water  
158 and kept at -80 °C for proximate analysis and fatty acid composition.  
159 Besides, 5 g of each diet was stored (-20 °C) at the beginning of the  
160 experimental trial in order to conduct the same analysis. Crude protein,  
161 moisture and ash content were analysed following A.O.A.C. methods  
162 (A.O.A.C., 2000). Total lipids were extracted (Folch, Lees & Sloane-  
163 Stanley, 1957) and fatty acids were prepared by trans-etherification  
164 (Christie, 1989). Separation and identification of the fatty acids was  
165 realized with gas chromatography (GC, THERMO FINNIGAN FUCUS  
166 GC, Milan, Italy) under the conditions reported in Izquierdo, Arakawa,  
167 Takeuchi, Haroun & Watanabe (1992).

168

#### 169 **2.5 Osteological studies**

170 For the characterisation of skeletal anomalies, a total of 15 larvae (50 dah)  
171 per tank were fixed in 10% buffered formalin and stained with alizarin red  
172 according to the methodology of Vandewalle, Gluckmann & Wagemans  
173 (1998). Terminology described by Mesa-Rodriguez *et al.* (2014) and  
174 Mesa-Rodriguez *et al.* (2016) was used for *S. rivoliana* bone structures

175 identification. The different regions of the axial column were divided and  
176 evaluated according to Boglione, Gagliardi, Scardi & Cataudella (2001).

## 177 **2.6 Statistical analysis**

178 All data were statistically treated using a SPSS Statistical Software  
179 System 15.0 (SPSS, [www.spss.com](http://www.spss.com)). The significant level for all the  
180 analysis was set at 5% and results are given as mean values and standard  
181 deviation. All values presented as percentage were arcsine transformed.  
182 All variables were checked for normality and homogeneity of variance,  
183 using the Kolmogorov-Smirnoff and the Levene tests, respectively. To  
184 compare means, the group data were statistically tested using one-way  
185 ANOVA. When variances were not homogenous, a non-parametric  
186 Kruskal-Wallis test was done. To evaluate the differences in skeletal  
187 frequency of deformities log linear statistical analysis were performed  
188 (Sokal & Rolf, 1995).

## 189 **3. Results**

190 *S. rivoliana* larvae survival was positively correlated with increasing  
191 dietary DHA levels ( $y=1.137x^2 - 4.121x + 73.48$ ;  $R^2 = 0.890$ ); with values  
192 ranging from 69.63% at 0.25% (dw) dietary DHA to 81.48% with 3.17%  
193 (dw) dietary DHA (Fig. 1). In addition, the increase of dietary DHA

194 significantly ( $P<0.05$ ) enhanced resistance to stress test (Fig. 2). On the  
195 other hand, no significant differences among treatments were observed in  
196 growth (Table 2) at middle ( $15.08\pm0.48$  mm TL) or final sampling points  
197 ( $19.80\pm0.58$  mm TL).

198 Fatty acid profiles of experimental fish was affected by increasing dietary  
199 DHA levels in weaning diets after 20 days of feeding the experimental  
200 feeds (Table 3). Total sum of saturated fatty acids (SFA) was highest in  
201 larvae fed the highest DHA levels (3.17%, dw; Diet 5), showing  
202 intermediate values in the larvae fed with a 2% of DHA (dw). Differences  
203 were also found in total monounsaturated fatty acid (MUFA) contents,  
204 finding the highest levels in larvae fed the lowest DHA levels (Diet  
205 DHA0), mainly due to increased contents of oleic acid (18:1n-9) in the  
206 feeds. The main SFA present in total body of *S. rivoliana* larvae were  
207 palmitic acid (16:0) and stearic acid (18:0). DHA contents in larval tissue  
208 showed a positive correlation with dietary DHA content, finding the  
209 lowest DHA levels in fish fed with DHA0 (0.25% DHA, dw) and the  
210 highest in DHA3 (3.17% DHA, dw; Table 3). ARA levels showed minor  
211 variations among dietary treatments, while a significant progressive  
212 decrease of EPA content was observed along with the increase in dietary  
213 DHA ( $P<0.05$ ). Total n-3 and total n-3 PUFA levels were positively

214 correlated with the DHA increase in the different dietary treatments. All  
215 the FA ratios were significantly ( $P<0.05$ ) affected by dietary treatment,  
216 thus ARA/EPA, DHA/EPA, DHA/ARA and n-3/n-6 ratios were increased  
217 according to the DHA contents in microdiets while the opposite trend was  
218 observed in oleic/DHA and oleic/n-3 PUFA ratios (Table 3).

219 Regarding the characterization of skeletal anomalies, scores showed no  
220 significant differences among dietary treatments ( $P>0.1$ ). The occurrence  
221 of cranial (jaw) abnormalities (6.7 - 4.4%) was only observed in larvae  
222 fed with the lowest dietary DHA treatments (Diets DHA0 and DHA1;  
223 0.25 and 0.75% DHA, respectively). However, a reduced incidence of  
224 skeletal deformities was observed in larvae fed the lowest dietary DHA  
225 treatment (DHA0), whereas increasing the dietary DHA content seemed  
226 to promote an increase in the number of total skeletal anomalies  
227 (kyphosis, lordosis, abnormal vertebra and cranial). In this sense, larvae  
228 fed with DHA2 (1.99% DHA, dw) showed the highest number of total  
229 anomalies. Furthermore, severe anomalies such as kyphosis and lordosis  
230 were absent in larvae fed DHA0 (0.25% DHA, dw). The occurrence of  
231 kyphosis and lordosis increased along with the dietary DHA contents  
232 (Fig. 3). Moreover, the occurrence of kyphosis was only observed in  
233 larvae fed with the highest dietary DHA treatments (Diets DHA2 and

234 DHA3; 1.99 and 3.17% DHA, respectively). Additionally, the incidence  
235 of abnormal vertebra centra was also in concordance with the increasing  
236 dietary DHA content.

## 237 **Discussion**

238 The inclusion of dietary DHA in inert diets up to 3.71 % (dw) increased  
239 the final survival in *S. rivoliana* larvae (81.5 %), being higher than  
240 previous studies with other marine finfish species such as 25% (Eryalçin  
241 et al., 2017), 45% (Saleh et al., 2013) and 48% (Hernández-Cruz et al.,  
242 2015) in *S. aurata* or 49% (Betancor et al., 2012b) and 73% in  
243 *Dicentrarchus labrax* (Cahu, Zambonino-Infante & Takeuchi, 2003). In  
244 agreement, larvae from species from the same genus fed with live preys  
245 enriched with DHA displayed enhanced final larval survival (Furuita et  
246 al., 1996b; Ishizaki, Takeuchi, Watanabe, Arimoto & Shimizu, 1998;  
247 Takeuchi, Ishizaki, Watanabe, Imaizumi & Shimizu, 1998; Yamamoto et  
248 al., 2008; Matsunari et al., 2012). For instance, *S. quinquerediata* larvae  
249 fed with *Artemia* sp. enriched with DHA (2.5 %, dw), showed enhanced  
250 final survival (88.5%) at 13 dah (Ishizaki et al., 1998). Another study in *S.*  
251 *dumerili* found the highest larval survival during the first 7 days (22%),  
252 when DHA contents increased up to 2.0% (dw; Matsunari et al., 2012).  
253 On the other hand, Yamamoto et al. (2008) stated that DHA contents

254 between 0.7-1.3 % (dw) in rotifers and 1.2-2.1 % (dw) in *Artemia* sp. did  
255 not satisfy DHA larval requirements for *S. dumerili*.

256 The increase of dietary DHA and EPA can improve, not only larval  
257 performance, but also stress resistance (Liu *et al.*, 2002; Izquierdo, 2005;  
258 Eryalçin *et al.*, 2013). In this sense, EFA play an important role as  
259 eicosanoids precursors (Ganga *et al.*, 2005) which play a pivotal role in  
260 stress response and immune system (Sargent, Bell, Bell, Henderson &  
261 Tocher, 1995). In the present study, *S. rivoliana* larvae fed increasing  
262 DHA levels from 0.25 % to 3.17 % (dw) showed improved resistance to  
263 air exposure along with the dietary increase of DHA. Similar results have  
264 been observed for *S. aurata* larvae fed with high DHA levels coming from  
265 marine phospholipids which showed better survival rate after handling  
266 (Saleh *et al.*, 2013; Saleh *et al.*, 2015). Additionally, the deficiency of  
267 DHA may reduce the tolerance to stressful conditions as observed in *Huso*  
268 *huso* larvae (Jalali *et al.*, 2008). It is known that deficiencies in structural  
269 components due to nutritional privation may produce a range of effects in  
270 the membrane of immune cells. These structural changes caused by  
271 component deficiencies in the membrane can alter eicosanoids production  
272 and membrane permeability. Moreover, cell membrane changes can also  
273 modulate the alternative complement pathway (ACP) activity as well as

274 the immune response in fish (Montero, Tort, Izquierdo, Robaina &  
275 Vergara, 1998).

276 On the other hand, inclusion of dietary DHA did not significantly affect *S.*  
277 *rivoliana* larval growth. Similar results have been reported in other marine  
278 teleost species, such as *Sparus aurata* (Izquierdo *et al.*, 2013; Hernández-  
279 Cruz *et al.*, 2015), *Pagrus pagrus* (Roo *et al.*, 2009), *Coryphaena*  
280 *hippurus* (Kraul, 1993) and *Centropomus parallelus* (Seifert, Cerqueira &  
281 Madureira, 2001), where fish performance was not influenced by  
282 increasing dietary levels of DHA. Contrarily to what could be expected  
283 taking into account other studies from the *Seriola* genus (Furuita *et al.*,  
284 1996b; Takeuchi *et al.*, 1998; Matsunari *et al.*, 2012), larval growth was  
285 slightly higher among the larvae fed the lowest DHA dietary content (Diet  
286 DHA0; 0.25% DHA, dw), albeit no significant differences were observed.  
287 This fact could indeed be related to larvae survival. Given that DHA0- fed  
288 larvae showed the lowest survival rate (although not significantly  
289 different), a higher amount of feed would be available per larvae.  
290 Moreover, an unbalanced DHA/EPA ratio seems to affect the growth in  
291 certain fish species (Izquierdo, 1996, 2005; Takeuchi, 1997; Shiozawa,  
292 Takeuchi & Hirokawa, 2003), indicating that not only the increasing  
293 levels of dietary DHA could promote the larvae final survival and growth,



294 but also an adequate ratio DHA to EPA. In this sense, Matsunari *et al.*  
295 (2012) observed the maximum total length in *S. dumerili* larvae fed a  
296 DHA/EPA ratio between 1.4 and 2.9, being this ratio much lower than the  
297 ones used in the present trial (up to 7.2).

298 The DHA/EPA ratio has been correlated with the dietary DHA  
299 supplementation. In the present study, an enhancement in survival after a  
300 challenge was observed when the DHA/EPA ratio was above 3.1  
301 (DHA1). This result is in agreement with the DHA/EPA ratio obtained in  
302 the tissues of wild specimens of the same genus such as *S. lalandi* and *S.*  
303 *dumerili* with DHA/EPA ratios of 3.5 and 5.6 respectively (O'Neill, Le  
304 Roux & Hoffman, 2015; Haouas, Zayene, Guerbej, Hammami & Achour,  
305 2010). Whitmore, *S. rivoliana* larvae fed with DHA0 and DHA1 with a  
306 DHA/EPA ratio lower than 1.4 showed significantly poor survival after  
307 activity test (Fig. 2), being in concordance with the minimum ratio  
308 suggested by Matsunari *et al.* (2012) of at least 1.4 for *S. dumerili* larvae.  
309 However, in other marine fish species, the optimum dietary DHA/EPA  
310 ratio during larval development seemed to be about 1.4 as it is the case for  
311 *Pagrus pagrus* (Hernández-Cruz *et al.*, 1999), 0.32 for *Dentex dentex*  
312 (Mourete, Tocher, Diaz-Salvago, Grau & Pastor, 1999), 1.2 for *S. aurata*  
313 (Rodríguez *et al.*, 1997) and 1.5 for *Lateolabrax japonicus* (Xu *et al.*,

314 2014). In these sense, it seems that *S. rivoliana* larvae needs higher  
315 DHA/EPA ratios than other commercially produced marine species,  
316 maybe related to the fast growth of this teleost.

317 As expected, the fatty acid compositions of the larvae mirrored the  
318 increasing dietary DHA levels. Therefore, larvae fed with high DHA  
319 contents consequently accumulated higher DHA and total n-3 LC-PUFA  
320 levels. Whitmore, the increase of MUFA levels, mainly oleic acid (18:1n-  
321 9) in larvae, was correlated with the low dietary DHA inclusion, given  
322 that olive oil, naturally rich in 18:1n-9, was used to equalize the lipid  
323 levels in the feeds. Contrarily, total body larvae fatty acid profile  
324 displayed increasing levels of total SFA when dietary DHA levels were  
325 increased, instead of decreasing its content with the minor amount of  
326 oleic. This is in agreement with other studies from species of the same  
327 genus, in which the comparison between wild and reared specimens  
328 showed that the main MUFA presented in muscle samples of both wild  
329 and reared fish was 18:1n-9, being the total amount of MUFA higher in  
330 wild specimens rather than in reared fish (*S. lalandi*; O'Neill *et al.*, 2015;  
331 *S. dumerili*; Rodriguez-Barreto *et al.*, 2012, 2014). In this sense, a  
332 comparison between reared and wild specimens of *S. quinquerradiata*  
333 determined that the triglycerides content observed in reared fish was

334 higher than in wild fish, as well as the amount of n-3 PUFA, particularly  
335 DHA (Arakawa *et al.*, 2002). Curiously, in other marine teleost species,  
336 increased DHA levels did not result in alterations in the total SFA content  
337 in larvae (Izquierdo *et al.*, 2013; Hernández-Cruz *et al.*, 2015).

338 Regarding skeletal abnormalities, the occurrence of cranial abnormalities  
339 in *Seriola sp.* has been previously reported (Cobcroft *et al.*, 2004). This  
340 author suggested that the inclusion of high DHA/EPA ratios, particularly  
341 around notochord flexion stages, and certain environmental factors such  
342 as light conditions may contribute to "wall-nosing" behaviour and the  
343 apparition of jaw malformations in yellowtail kingfish (*Seriola lalandi*).  
344 Conversely, in the present study, the reduction of cranial abnormalities  
345 was concomitant with the increased dietary DHA content. In previous  
346 studies, the appearance of skeletal muscle lesions (Betancor *et al.*, 2011)  
347 and the occurrence of skeleton anomalies (Villeneuve, Gisbert, Le  
348 Delliou, Cahu & Zambonino-Infante, 2005; Izquierdo *et al.*, 2010;  
349 Izquierdo *et al.*, 2013) were associated with increased dietary DHA levels.  
350 In this way, the incidence of skeletal anomalies in *S. rivoliana* larvae in  
351 the present study could be related with the high dietary DHA levels, albeit  
352 no significant differences were observed. Furthermore, the occurrence of  
353 severe anomalies such as kyphosis and lordosis, were mainly found in

larvae fed with the highest levels of DHA (Spearman correlation,  $p=0.9$ ). In this sense, severe deformities of the vertebral column always involve abnormalities over a relative wide range of vertebrae, which can appear fused and deformed, particularly in the region of the maximal axis curvature (Boglione *et al.*, 2001). This may explain the relationship between the numbers of severe abnormalities with abnormal vertebral bodies observed in the present study.

The relationship between n-3 LC-PUFA and the bone formation mechanism is still unknown. Previous studies in sea bream larvae indicated that DHA inclusion increased the n-3/n-6 ratio and could promote ossification (Izquierdo *et al.*, 2013), reduce vertebral fusion and cranial deformities in *P. pagrus* (Roo *et al.*, 2009) and decrease the incidence of opercular deformities in *Chanos chanos* (Gapasin & Duray, 2001). Moreover, low dietary DHA levels can delay early mineralization and increase the risk of cranial and axial skeletal deformities in sea bream larvae (Izquierdo *et al.*, 2013). Thus, high dietary DHA levels and adequate balance between pro and antioxidant nutrients seem to promote good skeletal health.

In summary, the results of the present study proved that the inclusion of dietary DHA in inert diets up to a 3.17% (dw) and a DHA/EPA ratio

374 above 3.1 increased the final survival and stress resistance in *S. rivoliana*  
375 larvae. Further studies on EFA requirements are required in order to  
376 enhance *S. rivoliana* larval production.

### 377 **Acknowledgements**

378 This study was funded by the Agencia Canaria de Investigación,  
379 Innovación y Sociedad de la Información (ACIISI) and Fondo Europeo  
380 de Desarrollo Regional (FEDER) through the program “Mejora de las  
381 técnicas de cría de larvas de (*Seriola rivoliana*): Determinación de  
382 requerimientos de ácidos grasos esenciales en su etapa larvaria y  
383 optimización de la secuencia alimentaria (METCSER- ProID20100094)”.

### 384 **References**

- 385 A.O.A.C. (2000). Official Methods of Analysis. Washington, DC:  
386 Association of Official Analytical Chemists
- 387 Arakawa, T., Ishizaki, Y., Chuda, H., Shimizu, K., Arimoto, M., &  
388 Takeuchi, T. (2002). Comparison of lipid classes and fatty acid  
389 compositions between hatchery reared and wild caught yellowtail *Seriola*  
390 *quinqueradiata* juvenile. Nippon Suissan Gakkaishi, 68, 374-381.
- 391 Bell, J.G., McEvoy, L.A., Estévez, A., Shields, R.J., & Sargent, J.R.

392 (2003). Optimising lipid nutrition in first-feeding at fish larvae.  
393 Aquaculture, 227, 211-220.

394 Betancor, M.B. (2012). Oxidative stress in sea bass, *Dicentrarchus labrax*  
395 L. larvae fed high DHA microdiets and its relation with several  
396 antioxidant nutrients. PhD Thesis, University of Las Palmas of Gran  
397 Canaria, Spain.

398 Betancor, M.B., Atalah, E., Caballero, M.J., Benitez-Santana, T., Roo, J.,  
399 Montero, D., & Izquierdo, M. (2011). Dietary a-tocopherol in weaning  
400 diets for European sea bass (*Dicentrarchus labrax*) improves survival and  
401 tissue damage caused by excess dietary DHA contents. Aquaculture  
402 Nutrition, 17, e112–e122.

403 Betancor, M.B., Caballero, M.J., Terova, G., Saleh, R., Atalah, E.,  
404 Benítez-Santana, T., Bell, J.G., & Izquierdo, M. (2012a). Selenium  
405 inclusion decreases oxidative stress indicators and muscle injuries in sea  
406 bass larvae fed high DHA microdiets. British Journal of Nutrition, 108,  
407 2115-2128.

408 Betancor, M.B., Caballero, M.J., Terova, G., Corà, S., Saleh, R., Benítez-  
409 Santana, T., Bell, J.G., Hernández-Cruz, C.M., & Izquierdo, M. (2012b).  
410 Vitamin C enhances vitamin E status and reduces oxidative stress

411 indicators in sea bass larvae fed high DHA microdiets. *Lipids*, 47, 1193-  
412 1207.

413 Boglione, C., Gagliardi, F., Scardi, M., & Cataudella., (2001). Skeletal  
414 descriptors and quality assessment in larvae and post-larvae of wild  
415 caught and hatchery reared gilthead sea bream (*Sparus aurata* L. 1758).  
416 *Aquaculture*, 192, 1 –22.

417 Bransden, M.P., Battaglione, S.C., Morehead, D.T., Dunstan, G.A., &  
418 Nichols, P.D. (2005). Effect of dietary 22:6n-3 on growth, survival and  
419 tissue fatty acid profile of striped trumpeter (*Latris lineata*) larvae fed  
420 enriched Artemia. *Aquaculture*, 243, 331-344.

421 Cahu, C., Zambonino-Infante, J., & Takeuchi, T. (2003). Nutritional  
422 components affecting skeletal development in fish larvae. *Aquaculture*,  
423 227, 245-258.

424 Christie, W. W. (2003). *Lipid Analysis*. (3rd Ed.). Bridgwater: Oily Press.

425 Cobcroft, J. M., Pankhurst, P. M., Sadler, J., & Hart, P. (2001). Jaw  
426 development and malformation in cultured striped trumpeter *Latris*  
427 *lineata*. *Aquaculture*, 199, 267-282.

428 Cobcroft, J.M., Pankhurst, P.M., Poortenaar, C., Hickman, B., & Tait, M.

429 (2004). Jaw malformation in cultured yellowtail kingfish (*Seriola lalandi*)  
430 larvae. New Zealand Journal of Marine and Freshwater Research, 38, 67-  
431 71.

432 Copeman, L.A., Parrish, C.C., Brown, J.A., & Harel, M. (2002). Effects  
433 of docosahexaenoic, eicosapentaenoic and arachidonic acids on the early  
434 growth, survival, lipid composition and pigmentation of yellowtail  
435 flounder (*Limanda ferruginea*): a live food enrichment experiment.  
436 Aquaculture, 210, 285–304.

437 Eryalçin, K.M., Roo, J., Saleh, R., Benitez, T., Betancor, M., Hernandez-  
438 Cruz, C.M., & Izquierdo, M. (2013). Fish oil replacement by different  
439 microalgal products in microdiets for early weaning of gilthead sea bream  
440 (*Sparus aurata*). Aquaculture Research, 44, 819-828.

441 Eryalçin, K.M., Torrecillas, S., Caballero, M.J., Hernández-Cruz, C.M.,  
442 Sweetman, J., & Izquierdo, M. (2017). Effects of dietary mannan  
443 oligosaccharides in early weaning diets on growth, survival, fatty acid  
444 composition and gut morphology of gilthead sea bream (*Sparus aurata*  
445 L.) larvae. Aquaculture Research, 48, 5041-5052.

446 Fernández-Palacios, H., Schuchardt, D., Roo, J., Hernández-Cruz, C.M.,  
447 & Izquierdo, M. (2015). Spawn quality and GnRH $\alpha$  induction efficiency



448 in longfin yellowtail (*Seriola rivoliana*) broodstock kept in captivity.  
449 Aquaculture, 435, 167-172.

450 Folch, J., Lees, N., & Sloane-Stanley, G.H. (1957). A simple method for  
451 the isolation and purification of total lipids from animal tissues. Journal of  
452 Biological Chemistry, 226, 497–509.

453 Furuita, H., Takeuchi, T., Toyota, M., & Watanabe, T. (1996a). EPA and  
454 DHA requirements in early juvenile red seabream using HUFA enriched  
455 *Artemia* nauplii. Fisheries Science, 62, 246–251.

456 Furuita, H., Takeuchi, T., Watanabe, T., Fujimoto, H., Sekiya, S., &  
457 Imaizumi, K. (1996b). Requirements of larval yellowtail for  
458 eicosapentaenoic acid, docosahexaenoic acid and highly unsaturated fatty  
459 acid. Fisheries Science, 62, 372–379.

460 Ganga, R., Bell, J.G, Montero, D., Robaina, L., Caballero, M.J., &  
461 Izquierdo, M. (2005). Effect of dietary lipids on plasma fatty acid profiles  
462 and prostaglandin and leptin production in gilthead seabream (*Sparus*  
463 *aurata*). Comparative Biochemistry and Physiology Part B, 142, 410-418.

464 Garcia-Gomez, A. (2000). Recent advances in nutritional aspects of  
465 *Seriola dumerili*. Paper presented at: Recent advances in Mediterranean

466 aquaculture finfish species diversification. Zaragoza: CIHEAM-IAMZ,  
467 2000. pp. 249–257: 6 graphs, 1 table, bibliography pp. 254–257,  
468 summaries (En, Fr) (Cahiers Options Méditerranéennes; v. 47). Seminar  
469 of the CIHEAM Network on Technology of Aquaculture in the  
470 Mediterranean on ‘Recent advances in Mediterranean aquaculture finfish  
471 species diversification’, 1999/05/24–28, Zaragoza (Spain).

472 Gapasin, R.S.J., & Duray, M.N. (2001). Effects of DHA-enriched live  
473 food on growth, survival and incidence of opercular deformities in  
474 milkfish (*Chanos chanos*). Aquaculture, 193, 49-63.

475 GIA (2011). Grupo de Investigación en Acuicultura (GIA). IU-  
476 ECOAQUA, Universidad de Las Palmas de Gran Canaria. Available  
477 online: [http://www.mispecies.com/nav/actualidad/noticias/noticia-](http://www.mispecies.com/nav/actualidad/noticias/noticia-detalle/Investigadores-del-GIA-de-la-ULPGC-avanzan-en-la-cra-larvaria-y-engorde-de-seriola/#.WAFOh-DhC00)  
478 [detalle/Investigadores-del-GIA-de-la-ULPGC-avanzan-en-la-cra-larvaria-](http://www.mispecies.com/nav/actualidad/noticias/noticia-detalle/Investigadores-del-GIA-de-la-ULPGC-avanzan-en-la-cra-larvaria-y-engorde-de-seriola/#.WAFOh-DhC00)  
479 [y-engorde-de-seriola/#.WAFOh-DhC00](http://www.mispecies.com/nav/actualidad/noticias/noticia-detalle/Investigadores-del-GIA-de-la-ULPGC-avanzan-en-la-cra-larvaria-y-engorde-de-seriola/#.WAFOh-DhC00)

480 Hamasaki, K., Tsuruoka, K., Teruya, K., Hashimoto, H., & Hamada, K.  
481 (2009). Feeding habits of hatchery-reared larvae of greater amberjack  
482 *Seriola dumerili*. Aquaculture, 288, 216–225.

483 Haouas, W.G., Zayene, N., Guerbej, H., Hammami, M., & Achour, L.  
484 (2010). Fatty acids distribution in different tissues of wild and reared

485 *Seriola dumerili*. International Journal of Food Science and Technology,  
486 45, 1478-1485.

487 Hernández-Cruz, C.M., Salhi, M., Bessonart, M., Izquierdo, M.,  
488 Gonzalez, M.M., & Fernández-Palacios, H. (1999). Rearing techniques  
489 for red porgy (*Pagrus pagrus*) during larval development. Aquaculture,  
490 179, 489-497.

491 Hernández-Cruz, C.M., Mesa-Rodríguez, A., Betancor, M., Haroun-  
492 Izquierdo, A., Izquierdo, M., Benítez-Santana, T, Torrecillas, S., & Roo,  
493 J. (2015). Growth performance and gene expression in gilthead seabream  
494 (*Sparus aurata*) fed microdiets with high docosahexaenoic acid and  
495 antioxidant levels. Aquaculture Nutrition, 21, 881-891.

496 Ishizaki, Y., Takeuchi, T., Watanabe, T., Arimoto, M., & Shimizu, K.  
497 (1998). A preliminary experiment on the effect of Artemia enriched with  
498 arachidonic acid on survival and growth of Yellowtail. Fisheries Science,  
499 64, 295-299.

500 Ishizaki, Y., Masuda, R., Uematsu, K., Shimizu, K., Arimoto, M., &  
501 Takeuchi, T. (2001). The effect of dietary docosahexaenoic acid on  
502 schooling behaviour and brain development in larval yellowtail. Journal  
503 of Fish Biology, 58, 1691–1703.

504 Izquierdo, M., Arakawa, T., Takeuchi, T, Haroun R., & Watanabe, T.,  
505 (1992). Effect of n-3 LC-PUFA levels in Artemia on growth of larval  
506 Japanese flounder (*Paralichthys olivaceous*). Aquaculture, 105, 73-82.

507 Izquierdo, M (1996). Essential fatty acid requirements of cultured marine  
508 fish larvae. Aquaculture Nutrition, 2, 183–191.

509 Izquierdo, M (2005). Essential fatty acid requirements in Mediterranean  
510 fish species. Cahier Options Mediterraneennes, 63, 91-102.

511 Izquierdo, M., & Koven, W.M. (2011). Lipids. In J. Holt (Ed), Larval  
512 Fish Nutrition (pp. 47–82). Oxford: Wiley-Blackwell.

513 Izquierdo, M, Watanabe, T., Takeuchi, T., Arakawa, T., & Kitajima, C.  
514 (1989). Requirement of larval red seabream *Pagrus major* for essential  
515 fatty acids. Nippon Suisan Gakkaishi, 55, 859–867.

516 Izquierdo, M, Socorro, J., & Roo, J. (2010). Studies on the appearance of  
517 skeletal anomalies in red porgy: effect of culture intensiveness, feeding  
518 habits and nutritional quality of live preys. Journal of Applied  
519 Ichthyology, 26, 320–326.

520 Izquierdo, M, Scolamacchia, M., Betancor, M.B., Roo, J., Caballero, M.J.,  
521 Terova, G., & Witten, P.E. (2013). Effects of dietary DHA and a-

522 tocopherol on bone development, early mineralisation and oxidative stress  
523 in *Sparus aurata* (Linnaeus, 1758) larvae. British Journal of Nutrition,  
524 109, 1796–1805.

525 Jalali, M.A., Hosseini, S.A., & Imanpour, M.R. (2008). Effect of vitamin  
526 E and highly unsaturated fatty acid enriched *Artemia urmiana* on growth  
527 performance, survival and stress resistance of Beluga (*Huso huso*) larvae.  
528 Aquaculture Research, 39, 1286-1291.

529 Kraul, S. (1993). Larviculture of the mahi-mahi *Coryphaena hippurus* in  
530 Hawaii, USA. Journal of the World Aquaculture Society, 24, 410–421.

531 Liu, J., Caballero, M.J., El-Sayed Ali, T., Izquierdo, M., Hernández-Cruz,  
532 C.M., Valencia, A., & Fernandez-Palacios, H. (2002). Effect of dietary  
533 lecithin and eicosapentaenoic acid on growth, survival, stress resistance  
534 and lipid transport in gilthead seabream (*Sparus aurata*). Fisheries  
535 Science, 6, 1165-1172.

536 Masuda, R., Takeuchi, T., Tsukamoto, K., Ishizaki, Y., Kanematsu, M., &  
537 Imaizumi, K. (1998). Critical involvement of dietary docosahexaenoic  
538 acid in the ontogeny of schooling behaviour in the yellowtail. Journal of  
539 Fish Biology, 53, 471–484.

540 Matsunari, H., Hashimoto, H., Oda, K., Masuda, Y., Imaizumi, H.,  
541 Teruya, K., Furuita, H., Yamamoto, T., Hamada, K., & Mushiake, K.  
542 (2012). Effects of docosahexaenoic acid on growth, survival and swim  
543 bladder inflation of larval amberjack (*Seriola dumerili*, Risso).  
544 Aquaculture Research, 44, 1696-1705.

545 Matsunari, H., Hashimoto, H., Iwasaki, T., Oda, K., Masuda, Y.,  
546 Imaizumi, H., Teruya, K., Furuita, H., Yamamoto, T., Hamada, K., &  
547 Mushiake, K. (2013). Effect of feeding rotifers enriched with taurine on  
548 the growth and survival of larval amberjack *Seriola dumerili*. Fisheries  
549 Science, 79, 815–821.

550 Mesa-Rodríguez A., Hernández-Cruz C.M., Socorro J.A., Fernández-  
551 Palacios H., Izquierdo M. & Roo J. (2014). Skeletal development and  
552 mineralization pattern of the vertebral column, dorsal, anal and caudal Fin  
553 complex in *Seriola Rivoliana* (Valenciennes, 1833) larvae. Journal of  
554 Aquaculture Research and Development, 5, 2155-9546.

555 Mesa-Rodríguez A., Hernández-Cruz C.M., Betancor. M.B, Fernández-  
556 Palacios H., Izquierdo M. & Roo J. (2016). Bone development of the  
557 skull, pectoral and pelvic fins in *Seriola rivoliana* (Valenciennes, 1833)  
558 larvae. Fish Physiology and Biochemistry, 42, 1777-1789.

559 Montero, D., Tort, L., Izquierdo, M., Robaina, L., & Vergara, J.M.  
560 (1998). Depletion of serum alternative complement pathway activity in  
561 gilthead seabream caused by  $\alpha$ -tocopherol and n-3 HUFA dietary  
562 deficiencies. *Fish Physiology and Biochemistry*, 18, 399-407.

563 Mourente, G., Tocher, D.R., Diaz-Salvago, E., Grau, A., & Pastor, E.  
564 (1999). Study of the high n-3 highly unsaturated fatty acids requirement  
565 and antioxidant status of *Dentex dentex* larvae at the Artemia feeding  
566 state. *Aquaculture*, 179, 291-307.

567 O'Neill, B., Le Roux, A., & Hoffman L.C. (2015). Comparative study of  
568 the nutritional composition of wild versus farmed yellowtail (*Seriola*  
569 *lalandi*). *Aquaculture*, 448, 169-175.

570 Papadakis, I.E., Chatzifotis, S., Divanach, P., & Kentouri, M. (2007).  
571 Weaning of greater amberjack (*Seriola dumerilii* Risso 1810) juveniles  
572 from moist to dry pellet. *Aquaculture International*, 16, 13–25.

573 Rezek, T.C., Watanabe, W.O., Harel, M., & Seaton, P.J. (2010). Effects of  
574 dietary docosahexaenoic acid (22:6n-3) and arachidonic acid (20:4n-6) on  
575 the growth, survival, stress resistance and fatty acid composition in black  
576 sea bass *Centropristis striata* (Linnaeus 1758) larvae. *Aquaculture*  
577 *Research*, 41, 1302–1314.

578 Rodríguez, C., Pérez, J.A., Díaz, M., Izquierdo, M., Fernández-Palacios,  
579 H., & Lorenzo, A. (1997). Influence of the EPA/DHA ratio in rotifers on  
580 gilthead seabream (*Sparus aurata*) larval development. Aquaculture, 150,  
581 77-89.

582 Rodríguez-Barreto, D., Jerez, S., Cejas, J.R., Martin, M.V., Acosta, N.G.,  
583 Bolaños, A., & Lorenzo, A. (2012). Comparative study of lipid and fatty  
584 acid composition in different tissues of wild and cultured female  
585 broodstock of greater amberjack (*Seriola dumerili*). Aquaculture, 360, 1-  
586 9.

587 Rodríguez-Barreto, D., Jerez, S., Cejas, J.R., Martin, M.V., Acosta, N.G.,  
588 Bolaños, A., & Lorenzo, A. (2014). Ovary and egg fatty acid composition  
589 of greater amberjack broodstock (*Seriola dumerili*) fed different dietary  
590 fatty acids profiles. European Journal of Lipid Science and Technology,  
591 116, 584-595.

592 Roo, J., Socorro, J.A., Izquierdo, M., Caballero, M.J., Hernández-Cruz,  
593 C.M., Fernández, A., & Fernández-Palacios (1999). Development of red  
594 porgy *Pagrus pagrus* visual system in relation with changes in the  
595 digestive tract and larval feeding habits. Aquaculture, 179, 499-512.

596 Roo, J., Hernandez-Cruz, C.M., Socorro, J.A., Fernandez-Palacios, H.,



597 Montero, D., & Izquierdo, M. (2009). Effect of DHA content in rotifers  
598 on the occurrence of skeletal deformities in red porgy *Pagrus pagrus*  
599 (Linnaeus, 1758). *Aquaculture*, 287, 84–93.

600 Roo, J., Hernandez-Cruz, C.M., Socorro, J., Fernandez-Palacios, H., &  
601 Izquierdo, M. (2010). Effect of rearing techniques on skeletal deformities  
602 and osteological development in red porgy *Pagrus pagrus* (Linnaeus,  
603 1758) larvae. *Journal of Fish Biology*, 77, 1309–1324.

604 Roo, J., Fernández-Palacios, H., Hernández-Cruz, C.M., Mesa-Rodriguez,  
605 A., Schuchardt, D., & Izquierdo, M. (2012). First results of spawning and  
606 larval rearing of longfin yellowtail *Seriola rivoliana* as a fast-growing  
607 candidate for European marine finfish aquaculture diversification.  
608 *Aquaculture Research*, 45, 689-700.

609 Saleh, R., Betancor, M.B., Roo, J., Benítez-Santana, T., Hernández-Cruz,  
610 C.M., Moyano, F.J., & Izquierdo, M. (2013). Optimum krill phospholipids  
611 content in microdiets for gilthead seabream (*Sparus aurata*) larvae.  
612 *Aquaculture Nutrition*, 19, 449-460.

613 Saleh, R., Betancor, M.B., Roo, J., Benítez-Dorta, V., Zamorano, M.J.,  
614 Bell, J.G., & Izquierdo, M. (2015). Effect of Krill phospholipids versus  
615 soybean lecithin in microdiets for gilthead seabream (*Sparus aurata*)

616 larvae on molecular markers of antioxidative metabolism and bone  
617 development. *Aquaculture Nutrition*, 21, 474 - 488.

618 Sargent, J.R., Bell, J.G., Bell, M.V., Henderson, R.J., & Tocher, D.R  
619 (1995). Requirement criteria for essential fatty acids. *Journal of Applied*  
620 *Ichthyology*, 11, 183- 198.

621 Seifert, M.E.B., Cerqueira, V.R., & Madureira, L.A.S. (2001). Effect of  
622 dietary (n-3) highly unsaturated fatty acids on growth and survival of fat  
623 snook (*Centropomus parallelus*, Pisces: Centropomidae) larvae during  
624 first feeding. *Brazilian Journal of Medical and Biological Research*, 34,  
625 645-651.

626 Shiozawa, S., Takeuchi, H., & Hirokawa, J. (2003). Improved seed  
627 production techniques for the amberjack, *Seriola dumerili*. *Saibai Giken*,  
628 31, 11-18.

629 Sokal, R.R., & Rolf, S.J. (1995). *Biometry: The principles and practice of*  
630 *statistics in Biological research*. (3rd ed.). NY: W.H. Freeman and  
631 Company,

632 Sims, N.A., & Key, G. (2011). Fish without footprints. *Oceans 2011*  
633 Conference, 19-22 September 2011, Waikoloa. Hawaii, United States of

634 America. pp: 1-6.

635 Takakuwa, F., Fukada, H., Hosokawa, H., & Masumoto, T. (2006).  
636 Optimum digestible protein and energy levels and ratio for greater  
637 amberjack *Seriola dumerili* (Risso) fingerling. *Aquaculture Research*, 37,  
638 1532–1539.

639 Takeuchi, T. (1997). Essential fatty acid requirements of aquatic animals  
640 with emphasis on fish larvae and fingerlings. *Reviews in Fisheries*  
641 *Science*, 5, 1-25.

642 Takeuchi, T. (2001). A review of feed development for early life stages of  
643 marine finfish in Japan. *Aquaculture*, 200, 203–222

644 Takeuchi, T. (2014). Progress on larval and juvenile nutrition to improve  
645 the quality and health of seawater fish: a review. *Fisheries Science*, 80,  
646 389–403.

647 Takeuchi, T., Ishizaki, Y., Watanabe, T., Imaizumi, K., & Shimizu, K.  
648 (1998). Effect of different DHA content in rotifers on the DHA  
649 requirement of larval yellowtail during *Artemia* feeding stage. *Nippon*  
650 *Suisan Gakkaishi*, 64, 280-275.

651 Tomas, A., de la Gandara, F., Garcia-Gomez, A., Perez, L., & Jover, M.

652 (2005). Utilization of soybean meal as an alternative protein source in the  
653 Mediterranean yellowtail, *Seriola dumerili*. Aquaculture Nutrition, 11,  
654 333–340.

655 Vandewalle, P., Gluckmann, I., & Wagemans, F. (1998). A critical  
656 assessment of the alcian blue/alizarin double staining in fish larvae and  
657 fry. Belgian Journal of Zoology, 128, 93 –95

658 Villeneuve, L., Gisbert, E., Le Delliou, H., Cahu, C.L., & Zambonino-  
659 Infante, J. (2005). Dietary levels of all-trans retinol affect retinoid nuclear  
660 receptor expression and skeletal development in European sea bass larvae.  
661 British Journal of Nutrition, 93, 1-12.

662 Villeneuve, L., Gisbert, E., Moriceau, J., Cahu, C.L., & Zambonino-  
663 Infante, J. (2006). Intake of high levels of vitamin A and polyunsaturated  
664 fatty acids during different developmental periods modifies the expression  
665 of morphogenesis genes in European sea bass (*Dicentrarchus labrax*).  
666 British Journal of Nutrition, 95, 677–687.

667 Watanabe, T., Izquierdo, M., Takeuchi, T., Satoh, S., & Kitajima, C.  
668 (1989). Comparison between eicosapentaenoic and docosahexaenoic acids  
669 in terms of essential fatty acids efficacy in larval red seabream. Nippon  
670 Suissan Gakkaishi, 5, 1635–1640.

671 Xu, H., Wang, J., Mai, K., Xu, W., Zhang, W., Zhang Y., & Ai, Q.  
672 (2014). Dietary docosahexaenoic acid to eicosapentaenoic acid  
673 (DHA/EPA) ratio influence growth performance, immune response, stress  
674 resistance and tissue fatty acid profile composition of juvenile Japanese  
675 seabass, *Lateolabrax japonicus* (Cuvier). Aquaculture Research, 47, 741-  
676 757.

677 Yamamoto, T., Teruya, K., Hara, T., Hokazono, H., Hashimoto, H.,  
678 Suzuki, N., Iwashita, Y., Matsunari, H., Furuita, H., & Mushiake, K.  
679 (2008). Nutritional evaluation of live food organisms and commercial dry  
680 feeds used for seed production of amberjack *Seriola dumerili*. Fisheries  
681 Science, 74, 1096–1108.

682

Table 1. Ingredients and proximate composition of the experimental microdiets containing increasing levels of DHA.

<i>Diet</i>	<i>DHA0</i>	<i>DHA1</i>	<i>DHA1.5</i>	<i>DHA2</i>	<i>DHA3</i>
<i>Ingredients (g kg<sup>-1</sup> diet)</i>					
<i>Defatted Squid meal</i> †	626.9	626.9	626.9	626.9	626.9
<i>DHA-50</i> ‡	0	20	50	70	90
<i>EPA 50</i> §	20.5	17.5	12.5	10.0	6.5
<i>ARA</i> ¶	12.5	12.5	10.0	10.0	8.0
<i>Oleic acid</i>	114.5	97.5	75.0	57.5	43.0
<i>Soy Lecithin</i>	30.0	30.0	30.0	30.0	30.0
<i>Vitamin mixture</i> ††	64	64	64	64	64
<i>Mineral mixture</i> ††	45.7	45.7	45.7	45.7	45.7
<i>Attractant</i> ††	55.9	55.9	55.9	55.9	55.9
<i>Gelatin</i>	30	30	30	30	30
<i>Proximate and FA analysis (g kg<sup>-1</sup> diet)</i>					
<i>Proteins (N×6.25)</i>	517.7	590.3	592.2	596.4	603.9
<i>Lipids</i>	205.4	194.6	204.9	191.1	185.2
<i>Moisture</i>	33.6	32.6	27.8	27.2	27.9
<i>Ash</i>	64.1	64.1	65.0	63.7	65.7
<i>Energy (MJ/kg)</i> †††	1,638.92	1,719.44	1,761.45	1,716.44	1,706.72
<i>DHA (%TFA/DW)</i>	2.76/ 0.25	8.90/ 0.75	18.35/ 1.64	25.83/ 1.99	35.26/ 3.17
<i>EPA</i>	6.42/ 0.58	6.58/ 0.56	5.91/ 0.53	5.64/ 0.44	4.88/ 0.44
<i>ARA</i>	3.36/ 0.3	3.73/ 0.32	3.76/ 0.94	4.14/ 0.32	4.11 / 0.37
<i>Saturated</i>	15.83/1.43	15.04/1.27	14.20/1.27	12.97/1.00	11.59/1.04
<i>Monosaturated</i>	56.74/5.12	50.87/4.3	42.07/3.75	36.00/2.78	28.40/2.55

685

686 † Squid meal (Agramar, Lorient, France),

687 ‡ DHA-50 Croda Chemicals Ltd. Goole, U.K.

688 § EPA-50 Croda Chemicals Ltd. Goole, U.K.

689 ¶ VEVODAR Oil.

690 †† Betancor et al., 2012

691 ††† Energy calculated as: fat×37.7 MJ/kg; protein×16.7 MJ/kg;

692

693 Table 2. *S. rivoliana* total length from 30 to 50 dah fed formulated diets with  
 694 increasing levels of dietary DHA (0.25, 0.75, 1.65, 2.0 and 3.17 g.kg<sup>-1</sup>dw DHA).  
 695 No significant differences were observed ( $P<0.05$ ).

696

697

698

699

700

	30 dah	42 dah	50 dah
<b>DHA0</b>	11.31 ± 1.79	15.91 ± 2.18	20.78 ± 3.54
<b>DHA1</b>	11.31 ± 1.79	14.87 ± 2.01	19.82 ± 3.49
<b>DHA1.5</b>	11.31 ± 1.79	15.02 ± 2.00	19.47 ± 2.86
<b>DHA2</b>	11.31 ± 1.79	14.66 ± 2.09	19.60 ± 3.35
<b>DHA3</b>	11.31 ± 1.79	14.94 ± 2.07	19.32 ± 2.59

701

702 DHA0, feed containing 0.25% DHA (dw); DHA1, feed containing 0.75% DHA  
 703 (dw); DHA1.5, feed containing 1.64% DHA (dw); DHA2, feed containing  
 704 1.99% DHA (dw); DHA3, feed containing 3.17% DHA (dw). Data expressed as  
 705 means ± SD (n = 3).

706

707 Table 3. Total fatty acid composition (%TFA) of 50dph larvae fed microdiets with  
708 increased levels of DHA (0.25, 0.75, 1.65, 2.0 and 3.17 g.kg<sup>-1</sup>dw DHA).

	DHA0	DHA1	DHA1.5	DHA2	DHA3
<i>Fatty acid content (%TFA)</i>					
<b>14:0</b>	0.26	0.27	0.29	0.30	0.39
<b>14:1n-5</b>	0.03	0.02	0.05	0.04	0.04
<b>14:1n-7</b>	0.01	0.01	0.01	0.01	0.01
<b>15:0</b>	0.12	0.13	0.16	0.16	0.21
<b>15:1n-5</b>	0.01	0.01	0.01	0.01	0.01
<b>16:0iso</b>	0.02	0.03	0.03	0.03	0.04
<b>16:0</b>	13.44	12.8	13.83	14.43	16.78
<b>16:1 n-7</b>	0.84	0.65	0.61	0.53	0.60
<b>16:1n-5</b>	0.07	0.07	0.10	0.13	0.16
<b>16:2n-6</b>	0.02	0.03	0.02	0.03	0.03
<b>16:2n-4</b>	0.17	0.21	0.24	0.31	0.38
<b>17:0</b>	0.03	0.03	0.03	0.04	0.04
<b>16:3n-4</b>	0.18	0.15	0.15	0.15	0.16
<b>16:3n-3</b>	0.04	0.04	0.05	0.06	0.07
<b>16:3n-1</b>	0.47	0.71	0.71	0.89	0.97
<b>16:4n-3</b>	0.45	0.64	0.58	0.68	0.65
<b>16:4 n-1</b>	0.05	0.10	0.11	0.13	0.14
<b>18:0</b>	5.8	6.48	6.90	8.01	9.19
<b>18:1 n-9</b>	41.11 <sup>d</sup>	31.02 <sup>c</sup>	23.79 <sup>b</sup>	20.58 <sup>ab</sup>	17.67 <sup>a</sup>
<b>18:1 n-7</b>	1.19	1.99	2.06	2.02	2.19
<b>18:1 n-5</b>	0.04	0.04	0.04	0.05	0.06
<b>18:2n-9</b>	0.09	0.09	0.08	0.09	0.11
<b>18:2 n-6</b>	12.18 <sup>b</sup>	10.73 <sup>b</sup>	10.72 <sup>a</sup>	8.66 <sup>a</sup>	8.40 <sup>a</sup>
<b>18:2n-4</b>	0.09	0.09	0.07	0.07	0.06
<b>18: 3n-6</b>	0.30	0.31	0.29	0.18	0.22
<b>18:3n-4</b>	0.06	0.063	0.05	0.03	0.04
<b>18:3 n-3</b>	1.30	1.16	1.18	0.94	0.94
<b>18:3n-1</b>	0.006	0.007	0.004	0.004	0.002
<b>18:4 n-3</b>	0.30	0.33	0.31	0.25	0.22
<b>18:4 n-1</b>	0.037	0.033	0.024	0.024	0.028
<b>20:0</b>	0.35	0.32	0.34	0.40	0.47



<b>20:1 n-9</b>	0.041	0.044	0.06	0.06	0.07
<b>20: 1n-7</b>	0.95	0.88	0.89	0.98	1.11
<b>20: 1n-5</b>	0.065	0.076	0.08	0.09	0.12
<b>20: 2n-9</b>	0.04	0.041	0.04	0.037	0.046
<b>20:2 n-6</b>	0.27	0.25	0.26	0.31	0.35
<b>20:3n-9+n-</b>	0.02	0.02	0.015	0.017	0.015
<b>20:3 n-6</b>	0.35	0.31	0.24	0.24	0.20
<b>20:4 n-6 (ARA)</b>	4.68 <sup>ab</sup>	5.25 <sup>b</sup>	4.74 <sup>ab</sup>	4.92 <sup>ab</sup>	4.51 <sup>a</sup>
<b>20: 3n-3</b>	0.17	0.18	0.20	0.22	0.24
<b>20:4 n-3</b>	0.31	0.26	0.21	0.18	0.17
<b>20:5 n-3 (EPA)</b>	5.34 <sup>c</sup>	5.23 <sup>c</sup>	4.19 <sup>ab</sup>	3.28 <sup>bc</sup>	2.55 <sup>a</sup>
<b>22:1 n-11</b>	0.05	0.07	0.10	0.08	0.12
<b>22:1 n-9</b>	0.23	0.25	0.24	0.25	0.30
<b>22:4 n-6</b>	0.28	0.32	0.34	0.40	0.42
<b>22:5 n-6</b>	0.20	0.69	1.13	1.37	1.49
<b>22:5 n-3</b>	1.28	1.30	1.12	1.06	0.98
<b>22:6 n-3 (DHA)</b>	6.68 <sup>a</sup>	16.26 <sup>b</sup>	23.26 <sup>c</sup>	27.23 <sup>c</sup>	26.97 <sup>c</sup>
<b>Saturated</b>	19.97 <sup>a</sup> ±2.66	20.04 <sup>a</sup> ±1.19	21.55 <sup>a</sup> ±1.49	23.35 <sup>ab</sup> ±0.97	27.09 <sup>b</sup> ±3.47
<b>Monoenoics</b>	44.63 <sup>d</sup> ±4.44	35.12 <sup>c</sup> ±0.98	28.05 <sup>b</sup> ±1.89	24.82 <sup>ab</sup> ±0.83	22.47 <sup>a</sup> ±2.49
<b>Total n-3</b>	15.87±3.36	25.40±2.28	31.10±2.35	33.91±1.93	32.80±5.22
<b>Total n-6</b>	18.27±0.92	17.90±1.10	17.76±0.86	16.11±0.78	15.62±1.37
<b>Total n-9</b>	41.51±3.26	31.44±0.71	24.20±1.55	21.01±0.62	18.19±1.78
<b>Total n-3PUFA</b>	13.78±3.01	23.22±2.18	28.98±2.16	31.98±1.80	30.91±5.09
<b>ARA</b>	4.67±0.44	5.25±0.19	4.74±0.08	4.92±0.04	4.51±0.25
<b>EPA</b>	5.34±0.93	5.23±0.26	4.18±0.39	3.28±0.20	2.55±0.37
<b>DHA</b>	6.68±1.68	16.26±1.78	23.26±1.70	27.23±1.57	26.97±4.53
<b>ARA/EPA</b>	0.88 <sup>a</sup> ±0.10	1.01 <sup>a</sup> ±0.72	1.13 <sup>a</sup> ±0.21	1.50 <sup>b</sup> ±0.19	1.76 <sup>b</sup> ±0.69
<b>DHA/EPA</b>	1.25 <sup>a</sup> ±0.11	3.11 <sup>b</sup> ±0.21	5.55 <sup>c</sup> ±0.62	8.30 <sup>d</sup> ±0.25	10.55 <sup>c</sup> ±0.63
<b>DHA/ARA</b>	1.43 <sup>a</sup> ±0.27	3.09 <sup>b</sup> ±0.22	4.90 <sup>c</sup> ±0.27	5.53 <sup>c</sup> ±0.28	5.97 <sup>c</sup> ±0.74
<b>oleic/DHA</b>	6.16±1.91	1.91±0.38	1.02±0.87	0.76±0.37	0.65±0.37
<b>oleic/n-3PUFA</b>	2.98±1.07	1.34±0.31	0.82±0.69	0.64±0.3	0.57±0.33
<b>n-3/n-6</b>	0.87 <sup>a</sup> ±0.16	1.42 <sup>b</sup> ±0.15	1.75 <sup>bc</sup> ±0.15	2.11 <sup>c</sup> ±0.17	2.10 <sup>c</sup> ±0.23

709

710

711

712


713

714

715

716

PUFA, polyunsaturated fatty acid; DHA, docosahexaenoic acid; EPA, eicosapentaenoic acid; ARA, arachidonic acid. DHA0, feed containing 0.25% DHA (dw); DHA1, feed containing 0.75% DHA (dw); DHA1.5, feed containing 1.64% DHA (dw); DHA2, feed containing 1.99% DHA (dw); DHA3, feed containing 3.17% DHA (dw). Data expressed as means ± SD (n = 3). Different superscript letters within a row denote significant differences among diets ( $P < 0.05$ ).




717

718 **Figure 1.** Survival rates (% of initial population) of *S. rivoliana* larvae fed  
 719 formulated diets with increasing levels of dietary DHA (0.25, 0.75, 1.65, 2.0 and  
 720 3.17 g.kg<sup>-1</sup>dw DHA) from 30 to 50 dah. Points show mean ± standard deviation  
 721 of three replicate tanks per diet, same letters denote that data are not significantly  
 722 different ( $P > 0.05$ ). The regression model represented by a line: survival =  
 723 1.137\*(DHA)<sup>2</sup> - 4.121\*DHA + 73.48, where DHA is g kg<sup>-1</sup> of dietary DHA  
 724 (polynomial regression, order 2).

725

726


727



728

729 **Figure 2.** Survival rates 24 h after activity test of *S. rivoliana* larvae fed  
 730 formulated diets with increasing levels of dietary DHA (0.25, 0.75, 1.65, 2.0 and  
 731 3.17 g.kg<sup>-1</sup>dw DHA) from 30 to 50 days after hatch. Activity test at 50 dah  
 732 consisted of 30 s air exposure. Points show mean  $\pm$  standard deviation of  
 733 different treatments, different letters denote that data were significantly different  
 734 ( $P < 0.05$ ). (Pearson correlation,  $r$ , is 0.99 with a significance of  $P=0.001$ ). The  
 735 regression model represented by a line: survival =  $0.859 \times (\text{DHA})^2 - 19.35 \times \text{DHA}$   
 736 + 7.033, where DHA is g. kg<sup>-1</sup> dw of dietary DHA (polynomial regression, order  
 737 2).

738



739

740 **Figure 3.** Incidence of skeletal deformities in *S. rivoliana* larvae fed formulated  
 741 diets with increasing levels of dietary DHA (0.25, 0.75, 1.65, 2.0 and 3.17 g kg<sup>-1</sup>  
 742 <sup>1</sup>dw DHA) at 50 dah. Sum. Anomalies (cranial + abnormal vertebra + fusion of  
 743 vertebra + Kyphosis + Lordosis); Sum. Abnormal vertebra (fusion of vertebra +  
 744 abnormal vertebra); Cranial (abnormal jaw).

745

746

747

748

749

750