

This is the peer reviewed version of the following article: Broome, A., Long, D., Ward, L. K. and Park, K. J. (2017), Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Appl Veg Sci*, 20: 397–409, which has been published in final form at <https://doi.org/10.1111/avsc.12303>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

**Promoting natural regeneration for the restoration of *Juniperus communis*:
a synthesis of knowledge and evidence for conservation practitioners.**

Alice Broome, Deborah Long, Lena K. Ward & Kirsty J. Park.

Broome, A. (Corresponding author, alice.broome@forestry.gsi.gov.uk)^{1,2}

Long, D. (Deborah.Long@plantlife.org.uk)³

Ward, L. (lkw@ceh.ac.uk)⁴

Park, K. (k.j.park@stir.ac.uk)²

¹Forest Research, Centre for Ecosystems, Society & Biosecurity, Northern Research Station, Roslin, Midlothian, EH25 9SY, UK

²University of Stirling, Biological & Environmental Sciences, Stirling, FK9 4LA, UK

³Plantlife Scotland, Balallan House, Allan Park, Stirling, 2QG, UK

⁴Centre for Ecology and Hydrology MacLean Bldg, Benson Ln, Crowmarsh Gifford, Wallingford, Oxfordshire OX10 8BB, UK

Key words: Juniper; Restoration; Conservation management; Cattle grazing; Scarification; Rotovating; Turf stripping; Soil stripping; Soil fertility; Moss cover; Microsites; Seed viability.

Nomenclature: Stace (2010) for plants; Rodwell (1991–2000) for plant communities; Hill et al. (2006) for mosses

Abstract

Questions: Natural regeneration is central to plant conservation strategies. Worldwide, many *Juniperus* species are threatened due to their failure to regenerate. We focus on *Juniperus communis* in areas of NW Europe where it is declining and ask: what advice is available to land managers on natural regeneration methods, and when applied, how effective has this been?

Methods: We synthesise knowledge on the efficacy of management interventions and conditions associated with *J. communis* regeneration. In field trials, we test interventions where knowledge is lacking. We assess regeneration of *J. communis*, creation of regeneration microsites and germination of sown seed in response to the interventions.

Results: Although *J. communis* occurs in different habitats, there is consistency in site conditions important for regeneration (unshaded/open, short ground vegetation, disturbed/bare ground, low herbivore pressure). In calcareous grasslands, areas with regeneration are stony/bare or vegetation is short or sparse; in upland acid grasslands and dry heathlands regeneration locations are disturbed areas sometimes with a moss cover. Several interventions (grazing, scarification, turf stripping) can create regeneration conditions. The synthesis identified cattle grazing and ground scarification for further testing on upland acid grasslands. In the resulting field trials, regeneration was rare and recorded on only one cattle grazed site. An exposed moss layer characterised regeneration microsites but there was insufficient evidence that either intervention increased regeneration microsite frequency. Few sown seeds germinated.

Conclusions: Different interventions or intensities of these appear to be required depending on habitat type. Broadly, on calcareous grassland intense scarification or soil stripping is needed, while on dry heathlands light scarification is suitable. On upland acid grassland, cattle grazing and ground scarification do not reliably result in regeneration. Creation of favourable mossy regeneration microsites is unlikely following intervention, unless soil fertility is low. Land use change, increased climate warming and pollution are pressures acting on *J. communis* and may cause habitat loss and altered site conditions (e.g. soil fertility), making it difficult to create regeneration microsites at all *J. communis* sites. Other constraints on regeneration may operate (e.g. seed predation and low seed viability) and managers should assess population and site potential before undertaking management.

Key words: Grasslands; Heathlands; Juniper; Management intervention; Microsites; Restoration; Seed viability.

Introduction

Self-sustainability in plant populations is a measure of ecological restoration success, with the occurrence of natural regeneration used as an indicator of a functioning ecosystem (Ruiz-Jaen & Aide 2005; Shackelford et al. 2013). Natural regeneration is central to plant conservation strategies and is considered the key process enabling species to adapt to climate change whilst maintaining local, site adapted, genetic resources and avoiding the risks associated with introducing plant material, such as novel pests and pathogens (Koskela et al. 2013; Lefèvre et al. 2013). For conifer species, natural regeneration has been widely and successfully achieved (Matthews 1989).

World wide, conifer species in the genus *Juniperus* show varying success of natural regeneration. For example, the North American species *Juniperus occidentalis* (western juniper) and *J. virginiana* (eastern redcedar) are currently undergoing population expansion whereas challenges to regeneration threatens *J. procera* (African pencilcedar) throughout its geographic range from the Arabian Peninsula to Zimbabwe (Miller et al. 2005; Meneguzzo & Liknes 2015; Negash & Kagne 2013). More typically, juniper species are a conservation concern in only part of their range, generally due to their failure to regenerate. This is the case for the montane species *J. thurifera* (Spanish juniper, incense juniper) of western Mediterranean regions and North Africa, and *J. communis* (common juniper), which occurs in western and eastern hemispheres, north of the equator (Farjon 2013a, www.iucnredlist.org/details/42255/0, accessed 28 November 2015; Farjon 2013b, www.iucnredlist.org/details/42229/0, accessed 28 November 2015). Although *J. communis* is not threatened with extinction globally in any of its forms (subspecies or varieties) (Farjon & Filer 2013), the species is struggling to survive in some areas, with changes in land use practices and site management identified as a factor driving reduction in plant survival and recruitment (Farjon 2013b).

Within Europe, *J. communis* (represented by the varieties *J. communis* L. var. *communis* and *J. communis* L. var. *saxatilis*) is an important component of designated habitats (calcareous heaths / grasslands, and coastal dunes) as given in Annex 1 of the EU Habitats Directive (92/43/EEC 1992). In boreal, alpine and eastern countries of Europe, these habitats are considered to be in ‘favourable’ condition (European Commission 2009)

and unwelcome invasions of *J. communis* into agricultural and other designated grassland habitats have even been reported in Scandinavia and Poland (Rosen 2005; Falinski 1998). However, within the Atlantic North and the Atlantic Central zones of Europe (EBONE 2009, www.ebone.wur.nl/UK, accessed 1 February 2012), both in the designated habitats and more widely, *J. communis* populations are declining and *J. communis* is of conservation concern. Factors impacting on natural regeneration of *J. communis* are noted as the main threats to the species (European Commission 2009; Joint Nature Conservation Committee 2010).

Success of natural regeneration is influenced by the availability of a seed source and microsites offering the correct conditions for germination and seedling survival (Eriksson & Ehrlén 1992). For a dioecious, sexually reproducing plant such as *J. communis*, all stages of the plant's life cycle have to be supported: pollination, viable seed production, seed dispersal and plant establishment, growth and development to reproductive maturity. As shown by studies of other long-lived (c. 200 years) conifer species, adult survival is likely to have a large influence on *J. communis* population dynamics with lesser importance placed on recruitment of individuals (indicated by successful germination or young seedling presence) for population survival (Thomas et al. 2007; Münzbergova et al. 2013; Kroiss & HilleRisLambers 2015). Nevertheless, recruitment appears to be a challenge for *J. communis* populations in the Atlantic North and the Atlantic Central zones of Europe.

Several studies have investigated reasons behind low seed production and viability. Many *J. communis* populations are aging and this is considered to reduce reproductive vigour (Ward 1982). Diffuse pollution has been shown to interrupt pollination, fertilisation and embryo development (Mugnaini et al. 2007), and nitrogen deposition, sulphur deposition, and increased temperatures can have a similar effect (García 2001; Verheyen et al. 2009; Ward 2010; Gruwez et al. 2014). A wide array of arthropods can act as pre-dispersal predators in *J. communis*, including mites (*Trisetacus quadrisetus*), and the chalcid wasp (*Megastigmus bipunctatus*) (Ward 1982; García 2001). Further, there may be decreased seed dispersal in areas where bird (*Turdus* spp.) numbers are lower (Eaton et al. 2009). Although it is thought that *J. communis* had historically high levels of pollen and seed-mediated gene flow, recent population fragmentation could be reducing effective gene flow with potential implications for the long-term fitness and survival of small populations even where viable seed production occurs (Van Der Merwe et al. 2000; Provan et al. 2008; Vanden-Broeck et al. 2011).

Timing the provision of suitable microsites is critical for successful regeneration of conifers as most have occasional mast years and the seed germinates when shed or following a short chilling period (e.g. Nixon & Worrell 1999). *Juniperus communis* has occasional years when seed production is abundant (Raatikainen & Tanska 1993; García et al. 1999; Bonner 2008; Ward 2010), but seed also displays a relatively deep dormancy which requires a longer period of exposure to natural winter conditions to break (Baskin & Baskin 2001; Bonner 2008). Seed is unlikely to germinate until the second spring following an autumn sowing and even then, germination can be sporadic (Broome 2003). Conditions suggested for successful germination and establishment of *J. communis* are associated with high light levels and unrestricted water availability is (Livingston 1972; Grubb et al. 1996; García et al. 1999). *Juniperus communis* is a community dominant in a range of open habitat types including upland acid grasslands, dry heathlands and lowland calcareous grasslands, and also occurs as a understorey species in pine woods and upland acid oak woodlands (Barkman 1985; Rodwell 1998a, b; Rodwell 1991; Thomas 2007). We therefore might expect the appearance of regeneration microsites and the processes by which they are created to vary with habitat type. Further, failure of *J. communis* to germinate and establish is thought to be due to a reduction in habitat suitability. Changes in site management leading to increased herbivore pressure are given as the primary causes for reduced suitability (Thomas et al. 2007). Therefore, there may be an opportunity to enhance natural regeneration of *J. communis* if management appropriate for the habitat conditions can be identified. *Juniperus communis* is declining within the Atlantic North and the Atlantic Central zones of Europe and here efforts to conserve the species and address the declines are required by European and country level legislation. In order to develop better guidance for conservation practitioners we conducted a literature review and field trials to:

- i. Synthesise information on conditions associated with *J. communis* regeneration and the efficacy of potential management interventions
- ii. Test the most suitable candidate interventions identified from the synthesis in field trials, particularly those which appear most practical to implement on the type of sites where managers are keen to restore *J. communis* populations. The specific objectives of the trials were (1) to evaluate natural regeneration of *J. communis* in response to two interventions (scarification and summer grazing by cattle), (2) to identify plant cover and composition of microsites where regeneration occurred and (3) assess whether the interventions created three measures of microsite condition identified in (2) and in the literature review. Given the uncertainties of seed viability and

dispersal for this species, a further objective (4) was to assess the germination of seed directly sown at the sites.

Methods

Methods are described in full in the online resource (Appendix S1); an outline of literature review and field trial methodologies is given here.

Literature review

We searched for information in two categories: i) surveys of *J. communis* (var. *communis* and var. *saxatilis*) sites where presence of regeneration was recorded, hereafter referred to as “regeneration surveys”; and ii) studies where management interventions had been applied in an attempt to enhance *J. communis* regeneration, referred to as “management studies”. Information was sought from countries within Atlantic North and the Atlantic Central environmental zones of Europe (EBONE 2009). The scientific literature was searched (up to November 2015) using *Juniperus communis* as the key word. Further information was sourced from book chapters and from published and unpublished reports produced by conservation agencies and organisations.

Field trials

We implemented a six-year trial on upland acid grassland habitats to test whether *J. communis* regeneration could be enhanced by ground scarification and /or by summer grazing by cattle. Four study sites in Scotland were used (Table S1 in Appendix S1): ‘Dorback’, ‘Pentland Hills’, ‘Fungarth’ and ‘Ballyoukan’. All sites had a population of *J. communis* var. *communis* containing reproductive female bushes growing within upland grassland and/or bracken (*Pteridium aquilinum*) communities. Scarification was applied at Dorback and Pentland Hills in the first year of the trial to produce a patchwork of bare areas with the matrix of unscarified vegetation acted as the control. The trial followed a blocked design (Figure S1 in Appendix S1). Grazing by cattle in the summer was applied annually at Fungarth and Ballyoukan to a target stocking intensity (Table S1 in Appendix S1). A control area was provided at each site using stock proof fencing (Figure S1 in Appendix S1). Seeds collected from local *J. communis* bushes were sown at a rate of 1600 seeds per m² in both treatment and control areas. Seeds were extracted from berries prior to sowing and tested for viability. This was estimated as: 41% (Dorback), 29% (Pentland Hills), 49% (Ballyoukan) and 6% (Fungarth). Germination was assessed annually in all the sown plots. In addition, natural regeneration was assessed by searching a 10 m wide buffer

around a sample of female bushes, annually. In the final year of monitoring, a systematic search of the site for *J. communis* seedlings was conducted at Ballyoukan and Fungarth only. Root collar diameter of seedlings was recorded, together with a description of the ground vegetation around each plant. Year of germination was estimated by dividing seedling root collar diameter by annual diameter stem increment figures. Vegetation monitoring was by annual measurements of vegetation height, vascular plant composition, and percentage cover of bare ground, litter and of all ground vegetation and moss species from permanent quadrats (0.25 m x 0.25 m). Depending on the complexity of the vegetation, between 80 and 180 quadrats were located on each site, evenly distributed between treatment and control areas and between vegetation types (where within-site differences occurred) (Table S1 in Appendix S1).

Data analysis

We implemented all tests in the package R (version 2.13.1; The R Foundation for Statistical Computing www.R-project.org/). For example: effect of cattle grazing on *J. communis* seedling occurrence at the Fungarth site, was tested using a Welch Two Sample t-test; associations between plant composition and occurrence of *J. communis* seedlings was investigated in a Principal Component Analysis (PCA); the relationship of seedling age with plant composition was tested using general linear models. The PCA was the first stage of the investigation of vegetation characteristics of microsites where regeneration occurred and contained plant composition and cover data (untransformed prior to analysis) from quadrats containing recently regenerated (1 to 2 year old) *J. communis* seedlings and a random sample of the permanent vegetation quadrats of an equivalent area containing no *J. communis* seedlings. In the second stage, we examined the effect of the variables with the strongest loadings on PCA axis 1 and 2 in general linear models with root collar diameter (a proxy for seedling age) as the response variable, for all the quadrats containing regeneration (seedlings ≤ 10 years). We investigated whether the two interventions applied in our trial produced the site condition measures (vegetation height, occurrence of exposed bare ground, and the occurrence of exposed moss cover) associated with regeneration microsites. We used linear mixed effects models to test the effect of scarification treatment on vegetation height and exposed bare ground although due to failure of model convergence, results for the latter have been presented descriptively. Due to failure of model convergence, result for the grazing treatments are presented descriptively.

Results

Site conditions associated with *Juniperus communis* regeneration from the review of regeneration surveys and studies

Results from seventeen regeneration surveys (Appendix S2) and seven management studies (Appendix S3) have been considered in this review and summarised (Table 1). These surveys and studies represent *J. communis* populations occurring on the full range of habitat types species occupies in Britain and other countries in the Atlantic North and the Atlantic Central European environmental zones (lowland dry heathlands, calcareous grasslands, upland pine-birch woodlands, upland acid grasslands and montane/coastal heath).

Results from the from the review of regeneration surveys

The survey methods followed in sixteen of the regeneration surveys was consistent: there was an element of identifying sample units of *J. communis* (usually populations), recording evidence of recent regeneration and providing information about site conditions and the land use/ site management. Regeneration was defined by the presence of juniper individuals estimated to be around five years old or younger although detection of very young (1 to 2 years old) seedlings is noted as difficult (A. Appleyard 2014, Botanical Surveyor, Salisbury, Wiltshire, personal communication). Two regeneration surveys were repeat surveys separated by several decades, a further five were designed to resample historical records and the remainder were generally searches of particular areas of interest e.g. nature reserves (Appendix S2).

Frequency of regeneration was generally low. Reports on eight of the regeneration surveys provide figures for the number of *J. communis* samples containing regeneration out of the total surveyed. These show an occurrence of regeneration in between 5% and 33% (median = 23%) of the sample units, per regeneration survey, respectively (Appendix S2). One further survey provided a cumulative count of 160 seedlings per hectare appearing at one site over the course of three years (Appendix S2). For the remainder of the regeneration surveys, results are descriptive with only the terms ‘very little’, ‘several’ and ‘a few’ used to describe the occurrence of regeneration, or counts of seedlings reported but no area of survey given (Appendix S2). Regeneration appears to relate to parent population size for most habitat types (Table 1) with, for example, minimum populations size of c.50 bushes required for regeneration to occur in upland grassland habitats (Appendix S2). However, where repeat surveys were conducted, a decline in the frequency of regeneration or poor inter-annual seedling survival (P. Woodruffe, A. Appleyard & S. Fitzpatrick 2016, Botanical Surveyors,

Salisbury, Wiltshire, personal communication) have been reported for *J. communis* in lowland calcareous grassland but not other habitat types (Appendix S2).

A short sward, and disturbed and bare ground/exposed mineral surface are associated with regeneration in the surveys of *J. communis* on upland grassland, lowland calcareous grassland and dry heathland sites (Table 1; Appendix S2). For *J. communis* populations on upland grassland in Scotland and heathland and grassland sites in Ireland, regeneration appeared to be associated with more nitrogen-limited sites (Appendix S2). On the very poor, dry heathland sites in the Netherlands, there are indications that regeneration is more prevalent on sites with higher base saturation of soil (e.g. 42% compared to 23%) or where there are pockets of higher pH (e.g. mean Hill-Ellenberg R value = 2.4) relative to the acidic surroundings (e.g. mean Hill-Ellenberg R value = 1.4) and also relatively more grass, fewer dwarf shrubs and more early successional mosses (Table 1; Appendix S2). Further, sites with little competition but shaded due to topographical position have been suggested as sites suitable for regeneration indicated by the presence of certain moss (*Hylocomium splendens*) and liverwort (e.g. *Lophozia ventricosa*) species characteristic of young *J. communis* stands (Appendix S3).

Reduced intensity of management (land use and site management) appeared to relate to presence of *J. communis* regeneration in several of the studies. For example reduced intensity of grazing by stock (usually sheep (*Ovis aries*) and cattle (*Bos taurus*) and other herbivores, e.g. rabbits (*Oryctolagus cuniculus*), is associated with regeneration in surveys of *J. communis* on upland grassland, lowland calcareous grassland and dry heathland sites (Table 1; Appendix S2; Appendix S3). Less intensive land management appeared to favour regeneration in open ground *J. communis* populations in Scotland, with regeneration being more frequent on land used for a combination of grazing and game interests rather than where management was for stock grazing only (Table 1; Appendix S2).

Management interventions used to encourage natural regeneration of *Juniperus communis* from the review of management studies

Reports on seven management studies were sourced; these investigated one or more of the following interventions: ground disturbance; reducing vegetation competition; reducing herbivore pressure; and changing soil pH (Table 1; Appendix S3). Three management studies were designed and monitored sufficiently to allow statistical analysis whilst the remaining management studies provided observational data only. Details on the

type, extent and duration of interventions aimed at encouraging the regeneration of *J. communis* are given in Appendix S3. Low impact ground disturbance e.g. by turf stripping or scarifying ground by dragging cut *J. communis* bushes increased regeneration for all habitat types in nearly all the management studies (Appendix S3) with failure in one study attributed to the limited area over which interventions were applied and/or poor seed viability (Appendix S3). It has been suggested that greater ground disturbance caused by cultivation is detrimental to survival of regenerating and young *J. communis* bushes (Appendix S3). Reducing vegetation competition through grazing was successful in two management studies on a calcareous grassland site, as *J. communis* seedlings regenerated either in the presence of sheep grazing (Appendix S3) during the summer months or during breaks in the grazing regime (Appendix S3) although seedling height was reduced by grazing. In one study, however, vegetation control by mowing or by herbicide treatment did not enhance recruitment of *J. communis* (Appendix S3). Successful germination, in a trial where seeds were sown on a dry heathland site, was attributed to increasing soil pH by liming yet this intervention did not encourage natural regeneration when applied within the adjacent *J. communis* stand (Appendix S3). In two separate management studies on upland acid grassland sites, reducing herbivore pressure (e.g. by excluding rabbits) was reported to benefit *J. communis* regeneration although the significance of the treatment effects could not be tested (Appendix S3). The interventions which appeared to promote regeneration most often (in 11 out of 12 surveys/studies) were those involving ground disturbance and reducing vegetation competition from the surrounding sward, particularly where they were applied in a less intensive way e.g. turf stripping rather than cultivation, sheep grazing rather than mowing. Lack of seed supply (or insufficient area over which treatment applied to successfully intercept available seed) was suspected as a cause of treatment failure in several of the management studies.

Germination of directly sown *Juniperus communis* seed in the field trials

Only two *J. communis* seedlings were recorded out of the 8000 (2500 of which were estimated as being viable) sown across all four sites, on both occasions in a scarified patch protected from stock grazing at the Pentland Hills site.

Response of *Juniperus communis* natural regeneration to intervention observed in the field trials

We wanted to evaluate the effect of two interventions (scarification and grazing) on *J. communis* regeneration. No regeneration was recorded from within the areas monitored around the *J. communis* bushes at any of the

sites. However, more widely within the site Fungarth, four seedlings (maximum height of 33cm) were found in 2008 and seven more seedlings were found in 2009. This indicated regeneration was occurring so a more widespread, systematic search of the cattle grazing sites was instigated. By 2011, a total of 33 seedlings (approximately 10 seedlings/ha) had been recorded and all at Fungarth; 23 in the grazed and 10 in ungrazed area. The grazing intervention had been applied in 2006. The ten ‘seedlings’ germinating prior to 2006 are equally distributed between the treatment and control areas. However a comparison of the number of seedling germinating in each of the six years when cattle grazing was applied shows that more *J. communis* seedlings germinating in the treated area compared to the control ($t = 2.60$, $df = 7$, $P = 0.032$, $n = 12$).

Vegetation cover and composition characterising regeneration microsites at one cattle grazed field trial site (Fungarth)

Axis 1 of the PCA bi-plot (Figure 1) describes a continuum from quadrats with a high % cover of moss and herbs and low % cover of grass or *Pteridium aquilinum* to quadrats with a low % cover of moss and herbs and high % cover of grass or *Pteridium aquilinum*. Axis 2 describes quadrats with a vegetation community dominated by *Pteridium aquilinum* to those of the grass dominated community. Quadrats containing *J. communis* seedlings (1 to 2 years old) are associated with higher % cover of moss and form a cluster relatively separate from the samples containing no regeneration. Together axis 1 and 2 explain 46% of the variation. Loading values for the species groups used in the PCA are given in Appendix S4. Regeneration microsites therefore appear to be characterised by an exposed moss cover i.e. a cover of moss that is not overlaid by other ground vegetation. When percentages of plant cover (moss, herbs, grass, *Pteridium aquilinum* (the variables with the strongest loadings on PCA axis 1 and 2) occurring with *J. communis* ‘seedlings’ up to 10 years old are analysed using general linear models, % moss cover appears as the best explanatory variable of root collar diameter (model with moss having lowest AIC value); ‘moss’ cover was higher ($F_{1,27} = 10.80$, $P = 0.003$; $R^2 = 0.29$, $n = 29$) where the seedlings were younger, as indicated by smaller root collar diameters (Figure 2). ‘Moss’ cover was negatively correlated with ‘grass’ cover (correlation coefficient = -0.64). These relationships suggest that there was more than 80% moss cover and very little grass cover present at the regeneration microsites at time of germination. The main moss species were typical of upland acid grassland sites in Britain (e.g. *Rhytidiadelphus squarrosus*, *Pseudoscleropodium purum*, *Pleurozium schreberi* and *Hylocomium splendens*)

(Rodwell 1998b).




Figure 1: Principal component analysis bi-plot showing the distribution of ground vegetation cover variables at the field trial site, Fungarth (both grazed and ungrazed area), in samples ($n = 45$) with (○) and without (Δ) *Juniperus communis* natural regeneration; regeneration indicated by the presence of 1 to 2 year old seedlings.




Figure 2: Vegetation composition of the *Juniperus communis* regeneration microsites and root collar diameter of seedlings at the field trial site, Fungarth (both grazed and ungrazed area) in 2011. Vegetation composition described as % cover of all

moss species; relationships were analysed using general linear models (solid line indicates the lines of best fit with 95% confidence intervals shown as dashed lines).

Creation of regeneration microsites by scarification and grazing treatments used in the field trial

Scarification produced exposed bare ground microsites which reverted to a grass sward after two and three growing seasons at Pentland Hills and Dorback, respectively (Appendix S5). An exposed moss cover was observed in very few of the quadrats (c.10%) during the re-vegetation process indicating that scarification does not reliably result in a layer of moss covering the ground prior establishment of other ground flora species.

Differences in vegetation height between control and treatment plots were still detectable at Dorback and Pentland Hills after five growing seasons following scarification (ANOVA: $F_{1,3} = 12.13$, $P = 0.040$, $n = 80$ for Dorback; $F_{1,3} = 10.67$, $P = 0.047$, $n = 157$ for Pentland Hills). Scarified areas had a mean vegetation height of 6.10 ± 1.14 cm compared to 13.00 ± 1.65 cm in the control at Dorback, and 9.80 ± 2.49 cm in the scarified areas compared to 13.30 ± 2.90 cm in the control at Pentland Hills.

At the grazed sites, exposed bare ground and moss microsites were infrequent (Appendix S6) and although there is some evidence that frequency of exposed bare ground increased in the final year of the trial at Ballyoukan in the grazed area, there appears to be no effect of grazing treatment on frequency of bare ground at Fungarth or exposed moss at either site. Grazing appears to reduce sward height at Ballyoukan, and at Fungarth there is some evidence of grazing causing a reduction in sward height especially in comparison with pre-grazed conditions (Appendix S7).

Discussion

Unlike other parts of Europe (e.g. Alps and Scandinavia) where *J. communis* is in favourable condition (European Commission 2009), *J. communis* sites within Atlantic North and the Atlantic Central European environmental zones require action to perpetuate *J. communis* populations in the face of multiple threats. It is not clear why these regional differences exist but in an attempt to control for variations in wide scale possible influences (e.g. climatic) on *J. communis* regeneration, we examined the options for promoting *J. communis* natural regeneration in the Atlantic North and Atlantic Central zones only. The literature available for synthesis

comprised seventeen regeneration surveys and seven management studies. Within these, data are often reported qualitatively or the quality of design and monitoring of the surveys and studies are insufficient to allow statistical analysis of data. As with any seedling survey where abundance is low, detection of infrequent and very young seedlings may be difficult and early stages of regeneration may be under estimated (McCarthy et al. 2013). However, an attempt to synthesise this range of information has not been undertaken before and this study provides insights in to the site conditions and management practices related to successful natural regeneration of *J. communis*. With an aim to strengthen the findings we tested the most suitable candidate interventions identified from the synthesis, in field trials.

Site and microsite conditions associated with *Juniperus communis* regeneration

The microsite conditions of the seed bed appears to be an important factor influencing regeneration of *J. communis* across the range of habitat types it occupies in the Atlantic North and the Atlantic Central zones of Europe. The regeneration surveys and management studies reviewed here indicate that reducing ground vegetation competition either by ground disturbance or lowering vegetation height, resulting in open/unshaded site conditions, are required for regeneration. However, regeneration microsites vary in the different habitat types. In calcareous grassland habitats, a bare surface appears to be the primary requirement for the regeneration of *J. communis* (Appendix S2; Wilkins & Duckworth 2011). Disturbed ground is also required within pine/birch woodland, acid grassland and dry heathland sites (Appendix S2). Microsites where we observed *J. communis* regeneration in the field trials conducted on acid grasslands were characterised by a cover of moss but an absence of taller vegetation. *J. communis* regeneration has been observed associated with a cover of unshaded moss on acid heathland sites and abandoned agricultural land (Falinski 1998; Appendix S2). Presence of a moss cover indicates that microsites must have high humidity at ground level - a requirement shown for *J. communis* regeneration in areas of Europe affected by summer drought (García et al. 1999). Mosses, along with lichens are often the first colonisers of nutrient poor-sites, where *J. communis* seedlings frequently occur (Wells et al. 1976). Moss cover has several positive effects on seed bed conditions, such as ameliorating temperature fluctuations, reducing frost heave as well as maintaining moist conditions (Parker et al. 1997; Groeneveld et al. 2007). The importance of preventing desiccation of seed of *Juniperus* species with extended stratification requirements such as *J. communis*, has long been recognised within the nursery trade (e.g. Heit, 1967). The surface of stones and rock fragments produced when calcareous soils are exposed may also maintain high humidity at the ground surface. Stones can act as mulch, reducing evapotranspiration of soil moisture (e.g. Perez

1998; Ma 2011) or provide a micro-watershed effect, creating suitable conditions for seedling establishment (Livingston 1972). In one survey, regeneration was associated with rock crevices which are assumed to have higher humidity (Appendix S2) and eroding chalk cliffs and limestone outcrops (often created by quarrying) have long been noted as suitable substrates for regeneration (Appendix S2; Grubb 1977; Ward 1981).

Soil fertility (usually reported in the reviewed literature as nitrogen availability and pH) also appears to be an important factor in determining appropriate site and microsite conditions. On acid habitat types, variations in soil fertility, even within a site, affected the occurrence of natural regeneration (Appendix S2). Vegetation studies on acid grassland sites reported that soil fertility affected vegetation succession on cleared ground, with an herbaceous sward as opposed to moss developing on sites with higher nutrient status (Miles 1973). On calcareous sites, the lack of soil in the regeneration microsite results in a relatively lower fertility of the surface material (Wells et al. 1976) and conservation practitioners have observed that on such sites, remaining topsoil acts as a growing medium and seed source for competitive native species, e.g. *Rubus fruticosus* (bramble), which rapidly colonise and shade areas prepared for *J. communis* regeneration (Wilkins & Duckworth 2011; F. Scully 2014, Community & Learning Ranger, National Trust, Guildford, Surrey, personal communication).

Management methods which create site and microsite conditions for *Juniperus communis* regeneration

The findings of this review suggest that management to create regeneration conditions for *J. communis* on all habitat types should aim to reduce competition from surrounding ground vegetation and provide protection from herbivores, primarily rabbits (Appendix S2). Reduced competition was most successfully achieved by manipulating herbivore management or mechanically removing ground vegetation. However, the outcome of applying similar management prescriptions is not always consistent between sites with differing soil fertility and highlights the difficulty of achieving both reduced vegetation competition and herbivore control (Appendix S3). Some evidence suggests that on acid dry heathland intensive disturbance (e.g. by cultivation) creates conditions less suitable for regeneration than the removal of surface litter and vegetation but there is limited evidence from the literature for the appropriate level of intervention on acid, upland grassland sites (Appendix S3). In our trials we tested scarification and cattle grazing in the summer but found they did not reliably enhance the natural regeneration of *J. communis*. Of the two interventions, summer cattle grazing appeared to have more potential for stimulating natural regeneration but confidence in predicting the results of this treatment at other sites is low. Scarification is clearly an inappropriate treatment on upland acid grassland sites. Evidence of creating

regeneration microsite conditions by the two management treatments used in the field trials is also lacking.

Unlike other studies (Ozols & Ozol 2007; Takala et al. 2012), we failed to show that a prolonged period of cattle grazing in the summer months increased the area of the site dominated by a moss cover. This was despite the partial removal of the bracken canopy by the cattle at our trial site (Fungarth), the effect of which has been linked with moss colonisation in other studies (Novak 2007). We found scarification was ineffective in creating an exposed moss cover that persisted for several years, instead a grass sward rapidly developed. Perhaps failure to develop a moss cover at our trial site was due to soil fertility being too high as a result of increasing nitrogen mineralisation from the soil disturbance (Russell 1961). On calcareous sites the success of management in promoting *J. communis* regeneration appears to be influenced by the depth of soil overlying the calcareous rock (and therefore the fertility of the site), and structure of the surface. Regeneration was reported to occur in a short sward resulting from stock grazing on thin soils in two of the management studies (Appendix S3). These types of sites also appear from the regeneration surveys to be the most suited to producing bare ground microsites by appropriate levels of stock grazing (Appendix S2). However, the experience of conservation practitioners suggests that scarification is a more reliable method of producing microsites which support *J. communis* regeneration particularly when the surface is composed of large chalk fragments (Wilkins & Duckworth 2011; F. Scully 2014, Community & Learning Ranger, National Trust, Guildford, Surrey, personal communication; J. Carey 2014, Countryside Officer, Bucks County Council, Aylesbury, Buckinghamshire, personal communication). These subtleties of interactions between disturbance type/ intensity, habitat type and site fertility, and development of the correct microsite may help explain the apparent contradictions over management regimes (type, duration and periodicity) which give rise to *J. communis* regeneration.

Wider constraints on *Juniperus communis* regeneration

Land use and site management changes, particularly changes in herbivore management and pressure, are viewed as strong drivers for changes in site suitability (Thomas et al. 2007; Fajhon 2013b). Over the last few centuries changing economic pressures has led to marginal land (often steeper slopes or nutrient poor grassland and heathland which is often associated with *J. communis* regeneration) has been over or under grazed by sheep/cattle, abandoned or ploughed (e.g. Wells et al. 1976; Appendix S3; Ward 1981). There is potential to reinstate grazing or increase protection of sites from herbivores and these types of manipulations have been identified as useful in our synthesis and further tested in the field trials. There have also been wide scale changes in soil fertility over last few decades. These changes have been associated with increased soil nitrogen levels

and acidification as a result of atmospheric deposition of ammonia and nitrogen oxides and by sulphur dioxide, respectively, and, although atmospheric deposition levels across Europe are lower than they were 20 years ago, there are still exceedances of critical loads for nitrogen (RoTAP 2012). Where site conditions have changed it may be possible to apply habitat manipulation to develop a seedbed but the causal factors for site change may have wider impacts. Failure to produce viable seed has been linked to high temperatures and nitrogen and sulphur deposition interrupting embryo development (Gruwez et al. 2014). It is possible that at many sites the unsuitability of microsites due to site fertility may indicate more fundamental failures in *J. communis* regeneration.

The importance of an adequate seed supply for successful natural regeneration is supported by the review; all the regeneration surveys which considered population size indicated a positive relationship between population size and regeneration (Appendix S2). All the *J. communis* populations studied in our trial produced berries and viability of sown seed (per population) ranged from 6% to 49%. This would seem adequate for natural regeneration, as Gruwez et al. (2013) reports recruitment at sites with 13% seed viability but no recruitment with 3% seed viability. However, the germination rate was low, with only two of the estimated 2500 viable seeds sown germinating. The absence of germination in our trial may be due to post-dispersal seed predation for example by mice (*Apodemus sylvaticus*) (García et al. 2001). Seedlings may have also been removed by small herbivores (rodents and slugs) before we recorded them; losses reported from other studies can be large, e.g. 6 seedlings out of 10,000 seeds survived the first year (García 2001). In hindsight, it may have been prudent to have provided protection to the patches of sown seed in our trial.

Conclusions

Natural regeneration is a fundamental process in the conservation of plant populations, and for *J. communis*, may be the only conservation option where risk of spread of pathogens e.g. *Phytophthora austrocedrae* (Green et al. 2014) from planting stock is high. The focus of this work is on relationships between site conditions and habitat management aimed at the restoration, through natural regeneration, of *J. communis* populations in the parts of NW Europe where the species is declining. By drawing together and adding to the existing body of information, we have further highlighted the difficulty in promoting the natural regeneration of *J. communis*. These findings, however, should be considered in the context of the wider constraints to natural regeneration recognised for *J. communis*: population fragmentation influencing gene flow, senescing/aging parent population

and pre-dispersal seed predation causing poor and infrequent seed production, reduced dispersal success and low seed viability.

Site conditions tolerated by the parent bushes of *J. communis* can differ from the microsites required for germination therefore habitat manipulation is required to develop a seedbed. Regeneration microsites need to be open (unshaded by ground vegetation) and provide moist conditions but may vary in appearance in different habitats. In calcareous grasslands, regeneration microsites are stony/bare or vegetation is short or sparse; in upland acid grasslands and dry heathlands microsites are disturbed areas sometimes with a moss cover. Grazing and ground disturbance are two commonly used techniques to create regeneration conditions. However, we found neither treatment to be a reliable intervention for enhancing natural regeneration of *J. communis* in upland acid grassland habitats. Similarly, many regeneration trials reviewed produced inconclusive findings and this synthesis showed mixed results for similar management interventions. No single type and intensity of management intervention appears best in all situations although grazing a site continuously appears inappropriate. Greater knowledge could be gained if more consistent and quantifiable methods are used in future management studies and regeneration surveys.

We suggest that where interventions are attempted, the soil fertility as well as the moisture availability and vegetation competition should be considered. For example, on acid grasslands focusing on sites where soil nutrient regime is poor, so that the intervention produces an extended successional stage of moss cover prior to development of a grass sward. Further, protection measures should be included as browsing by herbivores particularly rabbits is associated with failure of *J. communis* to regenerate, and post-dispersal seed predation may also reduce regeneration success. Managers should assess the potential of the site to support regeneration microsites, apply management measures for a minimum of five years, and be prepared to wait longer for results, as regeneration times are long.

Acknowledgments

We would like to thank Andrew McBride of Borders Forest Trust, Graham Sullivan of Scottish Natural Heritage and Pat Woodruffe, Anne Appleyard and Susan Fitzpatrick for making their data available, and Thomas Stolz, Patrick Hommel, Tim Wilkins and Tony Mundell for supplying reports of their research. The field trials were

made possible by SITA Tayside Biodiversity Action Fund, Scottish Natural Heritage and Forestry Commission Scotland, and support of the private owners and site managers. We would also like to thank Forest Research's Technical Support Unit staff, Jason Weber, Andrew Peace, Tom Connolly, Rachel Thompson, Mike Strachan, Helen Taylor and Tim Barratt for their assistance. Thanks also go to Deborah Long, Sallie Bailey, Joan Cottrell, Shelagh McCartan, Chris Quine and Helen McKay for comments on earlier drafts.

References

- Barkman, J.J. 1985. Geographical variation in associations of juniper scrub in the central European plain. *Vegetatio* 59: 67-71.
- Baskin, C. C. & Baskin, J.M. 2001. *Seeds: ecology, biogeography and evolution of dormancy and germination* Academic Press Inc, San Diego.
- Bonner, F. T. 2008. *Juniperus* L. Pages 607-614. In: Bonner FT, Karrfalt RP (eds) *Woody Plant Seed Manual* USDA FS Agriculture Handbook 727. US Agriculture Department, Forest Service, Washington, D.C.
- Broome, A. 2003. *Growing juniper: propagation and establishment practices*. Forestry Commission Information Note 50. Forestry Commission Edinburgh.
- Eaton, M.A., Brown, A.F., Noble, D.G., Musgrove, A.J., Hearn, R.D., Aebischer, N.J., Gibbons, D.W., Evans, A., Gregory, R.D. 2009. Birds of Conservation Concern 3: The population status of birds in the United Kingdom, Channel Islands and Isle of Man. *British Birds* 102: 296-341.
- Eriksson, O., Ehrlén, J. 1992. Seed and microsite limitation of recruitment in plant populations. *Oecologia* 91: 360-364.
- European Biodiversity Observation Network (EBONE) (2009) European Environmental Stratification.
- European Commission. 2009. Composite report on the conservation status of Habitat Types and Species as required under Article 17 of the Habitats Directive. Report from the Commission to the Council and the European Parliament [COM(2009), No. 358/F1] Brussels, Belgium.
- Falinski, J.B. 1998. Dioecious woody pioneer species (*Juniperus communis*, *Populus tremula*, *Salix* sp. div.) in the secondary succession and regeneration. *Phytocoensis*, Vol. 10 (N.S.) *Supplementum Cartographiae Geobotanicae* 8, Warszawa – Białowieża.
- Farjon, A. 2013a. *Juniperus thurifera*. The IUCN Red List of Threatened Species 2013.
- Farjon, A. 2013b. *Juniperus communis*. The IUCN Red List of Threatened Species 2013.

- Farjon, A., Filer, D. 2013. *An atlas of the world's conifers: an analysis of their distribution, biogeography, diversity and conservation status*. Brill, Leiden, Netherlands.
- García, D., Zamora, R., Hódar, J.A., Gómez, J.M. 1999. Age structure of *Juniperus communis* L. in the Iberian peninsula: Conservation of remnant populations in Mediterranean mountains. *Biological Conservation* 87: 215-220.
- García, D., Zamora, R., Gómez, J.M., Hódar, J.A. 2001. Frugivory at *Juniperus communis* depends more on population characteristics than on individual attributes. *Journal of Ecology* 89: 639 – 647.
- García, D. 2001. Effects of seed dispersal on *Juniperus communis* recruitment on a Mediterranean mountain. *Journal of Vegetation Science*, 12: 839-848.
- Green, S., Elliot, M., Armstrong, A., Hendry, S.J. 2014. *Phytophthora austrocedrae* emerges as a serious threat to juniper (*Juniperus communis*) in Britain. *Plant Pathology* 64: 456-466.
- Groeneveld, E.V.G., Massé, A., Rochefort, L. 2007. *Polytrichum strictum* as a nurse-plant in peatland restoration. *Restoration Ecology* 15: 709-719.
- Grubb, P.J. 1977. The maintenance of species-richness in plant communities: the importance of the regeneration niche. *Biological Reviews* 2: 107–145.
- Grubb, P.J., Lee, W.G., Kollmann, J., Wilson, J.B. 1996. Interaction of irradiance and soil nutrient supply on growth of seedlings of ten European tall-shrub species and *Fagus sylvatica*. *Journal of Ecology* 84: 827 – 840.
- Gruwez, R., Leroux, O., De Frenne, P., Tack, W., Viane, R., Verheyen, K. 2013. Critical phases in the seed development of common juniper (*Juniperus communis*) *Plant Biology* 15: 210–219.
- Gruwez, R., Frenne, P.D., Schrijver, A.D., Leroux, O., Vangansbeke, P., Verheyen, K. 2014. Negative effects of temperature and atmospheric depositions on the seed viability of common juniper (*Juniperus communis*). *Annals of Botany* 113: 489-500.
- Heit, C.E. 1967. Propagation from seed. Part 9: Fall sowing of conifer seeds. *American Nurseryman* 126: 60-69.
- Joint Nature Conservation Committee. 2010. UK Priority Species data collation *Juniperus communis* version 2 updated on 15/12/2010. Pages 1-18. In: Maddock A (ed.) *Species pages for 2007 UK BAP priority species*, Peterborough, UK.
- Koskela, J., Lefèvre, F., Schueler, S., Kraigher, H., Olrik, D.C., Hubert, J., Longauer, R., Bozzano, M., Yrjänä, L., Ditlevsen, B. 2013. Translating conservation genetics into management: Pan-European minimum

- requirements for dynamic conservation units of forest tree genetic diversity. *Biological Conservation* 157: 39-49.
- Kroiss, S.J., HilleRisLambers, J. 2015. Recruitment limitation of long-lived conifers: implications for climate change responses. *Ecology* 96:1286–1297.
- Lefèvre, F., Boivin, T., Bontemps, A., Courbet, F., Davi, H., Durand-Gillmann, M., Fady, B., Gauzere, J., Gidoïn, C., Pichot, S. 2013. Considering evolutionary processes in adaptive forestry. *Annals of Forest Science* DOI: 10.1007/s13595-013-0272-1.
- Livingston, R.B. 1972. Influence of Birds, Stones and Soil on the Establishment of Native Juniper (*Juniperus communis*) and Red Cedar (*J. virginiana*) in New England Pastures. *Ecology* 53: 1142 – 1147.
- Ma, Y.-J., Li, X.-Y. 2011. Water accumulation in soil by gravel and sand mulches: Influence of textural composition and thickness of mulch layers. *Journal of Arid Environments* 75: 432-437.
- Matthews, J.D. 1989. *Silvicultural Systems*. Clarendon Press, Oxford.
- McCarthy, M.A., Moore, J.L., Morris, W.K., Parris, K.M., Garrard, G.E., Vesk, P.A., Rumpff, L., Giljohann, K.M., Camac, J.S., Bau, S., Friend, T., Harrison, B., Yue, B. 2013. The influence of abundance on detectability. *Oikos* 122: 717-726.
- Meneguzzo, D.M., Liknes, G.C. 2015. Status and Trends of Eastern Redcedar (*Juniperus virginiana*) in the Central United States: Analyses and Observations Based on Forest Inventory and Analysis Data *Journal of Forestry* 113.3: 325-33.
- Miles, J. 1973. Natural recolonization of experimentally bared soil in callunetum in north-east Scotland. *Journal of Ecology* 61: 399-412.
- Miller, R.F., Bates, J.D., Svejcar, T.J., Pierson, F.B., Eddleman, L.E. 2005. *Biology, Ecology, and Management of Western Juniper* (*Juniperus occidentalis*) Technical Bulletin 152. Corvallis: Agricultural Experiment Station, Oregon State University.
- Mugnaini, S., Nepi, M., Guarnieri, M., Piotto, B., Pacini, E. 2007. Pollination drop in *Juniperus communis*: response to deposited material. *Annals of Botany* 100: 1475-1481.
- Münzbergová, Z., Hadincová, V., Wild, J., Kindlmannová, J. 2013. Population growth as a key factor in the invasion success of *Pinus strobus* *PLOS ONE* 8.2: e56953
- Negash, L., Kagnew, B. 2013. Mechanisms for the successful biological restoration of the threatened African pencilcedar (*Juniperus procera* Hochst. ex. Endl., Cupressaceae) in a degraded landscape. *Forest Ecology and Management* 310: 476-482.

- Nixon, C.J., Worrell, R. 1999. *The potential for the natural regeneration of conifers in Britain*. Forestry Commission Bulletin 120. Forestry Commission, Edinburgh.
- Novak, J. 2007. Regulation of *Pteridium aquilinum* (L.) Kuhn with triclopyr. *Ekologia* (Bratislava) 26: 211-221.
- Ozols, U., Ozols, R. 2007. Observations on vegetation development along pasture fence lines. *Tuexenia* 27: 307-326.
- Parker, W.C., Watson, S.R., Cairns, D.W. 1997. The role of hair-cap mosses (*Polytrichum* spp.) in natural regeneration of white spruce (*Picea glauca* (Moench) Voss). *Forest Ecology and Management* 92: 19-28.
- Pérez, F.L. 1998. Conservation of soil moisture by different stone covers on alpine talus slopes (Lassen, California). *Catena* 33: 155-177.
- Provan, J., Beatty, G.E., Hunter, A.M., McDonald, R.A., McLaughlin, E., Preston, S.J., Wilson, S. 2008. Restricted gene flow in fragmented populations of a wind-pollinated tree. *Conservation Genetics* 9:1521-1532.
- Pyatt DG, Ray D, Fletcher J (2001) An Ecological Site Classification for Forestry in Great Britain. Forestry Commission Bulletin 124, Forestry Commission, Edinburgh
- Raatikainen, M., Tanska, T. 1993. Cone and seed yields of the juniper (*Juniperus communis*) in southern and central Finland. *Acta Botanica Fennica* 149: 27-39.
- Rodwell, J.S. 1991. *British plant communities, I: Woodlands and Scrub*. Cambridge University Press, Cambridge, UK.
- Rodwell, J.S. 1998a. *British plant communities, vol. 2: Mires and heaths*. Cambridge University Press, Cambridge, UK.
- Rodwell, J.S. 1998b. *British plant communities, vol. 3: Grassland and montane communities*. Cambridge University Press, Cambridge, UK.
- Rosén, E. 2006. Alvar vegetation of Öland - changes, monitoring and restoration. *Biology and Environment: Proceedings of the Royal Irish Academy* 106, 387-399.
- RoTAP. 2012. *Review of Transboundary Air Pollution: Acidification, Eutrophication, Ground Level Ozone and Heavy Metals in the UK*. Contract Report to the Department for Environment, Food and Rural Affairs. Centre for Ecology and Hydrology, UK.

- Ruiz-Jaén, M.C. & Aide, T. M. 2005. Vegetation structure, species diversity, and ecosystem processes as measures of restoration success. *Forest Ecology and Management* 218: 159-173.
- Russell, E.W. 1961. Chapter 16: Nitrification in Forest Soils. Pages 310 - 317. In: *Soil Conditions and Plant Growth*. 9th edition. Longmans, London, UK.
- Shackelford, N., Hobbs, R.J., Burgar, J.M., Erickson, T.E., Fontaine, J.B., Lalibert, E., Ramalho, C.E., Perring, M.P., Standish, R.J. 2013.. Primed for Change: Developing Ecological Restoration for the 21st Century. *Restoration Ecology* 21: 297-304.
- Takala, T., Tahvanainen, T., Kouki, J. 2012. Can re-establishment of cattle grazing restore bryophyte diversity in abandoned mesic semi-natural grasslands? *Biodiversity and Conservation* 21: 981-992.
- Thomas, P., El-Barghathi, M., Polwart, A. 2007. Biological Flora of the British Isles: *Juniperus communis* L. *Journal of Ecology* 95: 1404-1440.
- Vanden-Broeck, A., Gruwez, R., Cox, K., Adriaenssens, S., Michalczyk, I.M., Verheyen, K. 2011. Genetic structure and seed-mediated dispersal rates of an endangered shrub in a fragmented landscape: a case study for *Juniperus communis* in northwestern Europe. *BMC GENETICS* 12: 1-16.
- Van der Merwe, M., Winfield, M.O., Arnold, G.M., Parker, J.S. 2000. Spatial and temporal aspects of the genetic structure of *Juniperus communis* populations. *Molecular Ecology* 9: 379 – 386.
- Verheyen, K., Adriaenssens, S., Gruwez, R., Michalczyk, I.M., Ward, L.K., Rosseel, Y., Van den Broeck, A., García, D. 2009. *Juniperus communis*: victim of the combined action of climate warming and nitrogen deposition? *Plant Biology* 11: 49-59 .
- Ward, L.K. 1981. The demography, fauna and conservation of *Juniperus communis* in Britain. In: *The Biological aspects of rare plant conservation*, ed. H Synge, 319-329. John Wiley & Sons. Chichester.
- Ward, L.K. 1982. The conservation of juniper: longevity and old age. *Journal of Applied Ecology* 19: 917-928.
- Ward, L.K. 2010. Variation in ripening years of seed cones of *Juniperus communis* L. *Watsonia* 28:11-19.
- Wells, T.C.E., Sheail, J., Ball, D.F., Ward, L.K. 1976. Ecological studies on the Porton ranges: Relationships between vegetation, soils and land-use history. *Journal of Ecology* 64: 589-626.
- Wilkins, T.C., Duckworth, J.C. 2011. *Breaking new ground for juniper- a management handbook for lowland England*. Plantlife, Salisbury, UK.

Table 1: Support for the effects of site variables on promoting (Δ) or restricting (\blacktriangledown) regeneration of *Juniperus communis* by habitat type. Number of regeneration surveys (S) or management studies (T) reporting effect are indicated beside triangle. With the exception of Lowland Heathland, soil pH values are drawn from Pyatt et al. 2001 and soil moisture regime (SMR) and soil nutrient regime (SNR) from Pyatt unpubl. 2000 (SNR classes: Very Poor (VP), Poor (P), Medium (M), Very Rich (VR); SMR classes: Moist (M), Fresh (F), slightly Dry (SD)).

Habitat		Bare ground/ disturbance	Reduced sward height	Herbivore pressure	Soil nutrient levels	Soil moisture	Parent population size
Upland/Lowland							
Lowland calcareous grassland/ scrub (NVC types CG3, CG4 & W21); soil pH = 4.5 -7.5, SMR = F, SNR = VR.	S	Δ^2	Δ^2	\blacktriangledown^2 -if no break in stock grazing	\blacktriangledown^1 - better if soil fertility is low	Δ^1 -negative effect of summer drought	Δ^2 - particularly young bushes a bushes bearing berries
				\blacktriangledown^3 -if browsing by rabbits	Δ^1 - better if soil pH is higher		
				Δ^3 -if grazing intermittent			
	T	Δ^1	Δ^1	Δ^1 -if summer sheep grazing used			
Lowland heathland (NL/Belgium/ Denmark); soil pH = 3.8-4.8 ^{7,10} .	S	Δ^2	Δ^3	Δ^2 -if grazing is managed	Δ^2 -with base enrichment of soil		
				\blacktriangledown^1 -if browsing by rabbits			
	T	\blacktriangledown^1			Δ^1 -with base enrichment of soil.		
Upland acid & mesotrophic grassland/scrub and heathland (UK-NVC type W19 & H15); soil pH = 3-5, SMR = M-F, SNR = VP-M.	S	Δ^2	Δ^1	Δ^2 -if fluctuating or in pulses	\blacktriangledown^2 -better on more acidic and nitrogen limited sites	Δ^2 -layering promoted (bushes propagate by stems touching ground)	Δ^5
				\blacktriangledown^4 -if exposed to grazing/more extensive landuse			
	T	Δ^1		\blacktriangledown^2 -if stock and rabbits are			

Natural regeneration of *Juniperus communis*

not excluded									
Pine/birch woodland (UK –NVC type W17/18); soil pH = 3-4, SMR = SD-F, SNR = VP-P.	S					▼ ¹	-better on more nitrogen limited sites	Δ ¹	Δ ¹
	T	Δ ¹	Δ ¹	▼ ¹	-small rodents and slugs				

Supporting information to the paper

Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S1. Methods

Literature search and general review methods

To assess site conditions associated with *J. communis* regeneration and management interventions likely to promote regeneration, we searched for information in two categories: i) surveys of *J. communis* sites and ii) studies where management interventions had been applied. To allow comparisons of *J. communis* response on a comparable range of habitat types and relatively similar range of climatic conditions encountered, within the natural range of *J. communis*, information was sought from countries within Atlantic North and the Atlantic Central environmental zones of Europe (EBONE 2009, <http://www.ebone.wur.nl/UK/>> project information and products /european environmental stratification page, accessed 1 February 2012).

The scientific literature was searched (up to November 2015) primarily using the Web of Knowledge within the subject areas of environmental sciences, ecology, forestry and biodiversity conservation using *Juniperus communis* as the key word. Further information was sourced from book chapters and from published and unpublished reports produced by conservation agencies and organisations. Information extracted from the literature has been summarised under a common set of headings in tables two tables (surveys, management studies). The surveys are described (location/habitat type, observations made) and any site characteristics positively or negatively associated with natural regeneration are listed. Similarly for the management studies, data on location/habitat type as well as interventions and outcomes have been listed. Records of soil pH and levels of nitrogen (N), phosphorus (P), potassium (K), aluminium (Al) and calcium (Ca) given in the surveys or studies have been replicated in the tables. Otherwise, we have used any ground vegetation data in conjunction with their indicator values (Hill et al. 1999) to derive scores of soil pH (R), soil nitrogen availability ('N') and soil moisture (F) using a mean abundance/frequency weighted approach (Pyatt et al. 2001). Where possible we have also reported the soil nutrient regime (R + 'N') and soil moisture regime (F) classes associated with the scores (Pyatt et al. 2001). In addition, original data sets have been made available to the authors from two Scottish *J. communis* surveys (Sullivan 2003 and Borders Forest Trust 1997) allowing a more detailed analysis to be conducted. Given the limited number of surveys and studies available, and the wide variation in methodologies followed and types of data collected by these, we followed a literature synthesis approach (e.g. Humprey et al. 2015) rather than a full systematic review or meta-analysis to assess the data (Koricheva & Gurevitch 2013).

Study areas

The study was conducted from January 2005 to March 2011. Four study sites, located in three administrative regions of Scotland were used (Table S1): Highland (with one site 'Dorback'); Midlothian (with one site 'Pentland Hills'); and Perthshire (with two sites 'Fungarth' and 'Ballyoukan'). Prior to the experimental management, sites were subject to various levels of grazing throughout the year by sheep (Dorback; Pentlands - a subsection only) and deer and rabbits (all), resulting in a tight sward and/or dense thatch of

litter. Very little natural regeneration of *J. communis* had been observed recently at any of the sites, which was felt to be due to inappropriate seed bed conditions resulting from site management prior to the start of the trial (R. Thompson 2003, conservation advisor, Scottish Natural Heritage, Battleby, Perthshire, personal communication; D. Granger 2004, local land manager, Dunkeld, Perthshire, personal communication).

Management treatments

Two sites were subject to scarification treatment (Dorback and Pentland Hills) . At Dorback scarification was performed by a tractor mounted rotary cultivator which produced a patchwork of bare areas of approximately 0.25m² with 1 metre spacing. At Pentland Hills the vegetation was cut and the ground surface was scarified in 1m² patches using a hand spade, again at 1m spacing. At Dorback and Pentland Hills, the trial had a block design and was blocked with respect to extensive stock grazing at Pentland Hills (see Figure S1 for schematic). At both sites deer had access and, with the exception of the extensively grazed area at Pentland Hills, rabbits were excluded.

At the other two sites, Fungarth and Ballyoukan, the treatment was grazing by cattle in the summer (Table S1) Stocking densities of cattle were within the range for upland birchwoods in Scotland, where woodland regeneration was occurring in the presence of grazing (Pollock et al. 2005; Table S1). The cattle-grazed sites each contained two different vegetation types (Table S1) which were represented in both the treatment and control areas. A set of permanent 25 cm x 25 cm quadrats were located in all treatment and control areas (Figure S1, Table S1). Grazing treatment commenced in 2006 at Fungarth and 2007 at Ballyoukan. Deer and rabbits were excluded from Fungarth but were present in low numbers at Ballyoukan.

Table S1: Summary descriptions of sites used, management interventions applied and monitoring in the *Juniperus communis* regeneration trial

Site	Lat & Long (NGR)	Elevation & Aspect	Solid geology	Size ³ (ha)	Intervention - type	- year applied	Habitat/Vegetation type (NVC community code ⁴),Site Fertility (SNR) & Wetness (SMR) ⁵	Monitoring - number of quadrats	- year (months) ⁶
Dorback	57°7'N, 3°5'W (NJ056192)	380m SW	Granite, syenite, granophyre and allied types ¹	5.60	Scarification;	2005	Improved upland acid grassland (U4).	20 (Treatment)	2005-2010
					Release from sheep grazing		SNR M; SMR F-M	20 (Control)	(July - October)
				0.75			Upland acid grassland (U4).	20 (Treatment)	
							SNR P- M; SMR F -M	20 (Control)	
Pentland Hills	55°9'N, 3°2'W (NT229649)	335m SE	Lower Old Red Sandstone - rhyolite and felsite ²	0.25	Scarification	2005	Wavy hair-grass grassland (U2)	40 (Treatment)	2005-2010
							SNR VP-P; SMR F-M.	40 (Control)	(September)
				0.13			Bracken community (U20)	40 (Treatment)	
							SNR P; SMR M	40 (Control)	
Fungarth	56°6'N, 3°6'W (NO045425)	130-320m NW	Devonian and Old Red Sandstone- andesitic and basaltic lavas and tuffs ¹	25	Summer cattle grazing (0.44 LSU /ha /year ⁷ . Breed= Limousin)	2006	Mosaic of upland acid grassland (U4) & bracken community (U20).		2005-2010
							U4: SNR P – M; SMR F – M	40 (Treatment)	(September - November)
								40 (Control)	
							U20: SNR P; SMR F – M.	50 (Treatment)	
								50 (Control)	

Ballyoukan	56°7'N, 3°7'W (NN968570)	180-285m SW	Upper Dalradian- quartz-mica-schist, grit, slate and phyllite ¹	12	Summer cattle grazing (0.41 LSU /ha /year ⁷ . Breed=Highland	2007	Mosaic of upland acid grassland (U4) and purple moor-grass mire (M25).	2005-2010 (August- October)
							U4: SNR P- M; SMR F-M	50 (Treatment)
								50 (Control)
							M25: SNR VP-P; SMR VM -W.	30 (Treatment)
								30 (Control)

¹ British Geological Survey (1979).

² British Geological Survey (1928).

³ includes both control and treated areas

⁴ National Vegetation Classification (Rodwell 1991; Rodwell 1998a; Rodwell 1998b)

U4: *Festuca ovina*-*Agrostis capillaris*- *Galium saxatile* grassland

U20 *Pteridium aquilinum*-*Galium saxatile* community

M25 *Molinia caerulea*-*Potentilla erecta* mire

U2 *Deschampsia flexuosa* grassland

⁵ Soil Nutrient Regime (SNR) and Soil Moisture Regime (SMR) derived from vascular plant composition (Pyatt et al. 2001) prior to intervention, indicates site fertility and site wetness at start of the trial; increasing soil fertility with SNR classes Very Poor (VP) < Poor (P) <Medium (M); increasing soil wetness with SMR classes Fresh (F) < Moist (M) < Very Moist (VM) < Wet (W). Pyatt, et al. 2001).

⁶ Assessment carried out once in this period of the year.

⁷ Live Stock Units (LSU) – livestock unit value for suckler cow (including calf at foot) Highland breed = 0.7, Limousin breed = 1.1 (Chesterton 2006); LSU/ha/year = (monthly livestock number in the treatment area averaged over whole year x the livestock unit value for the breed)/ size of treatment area (ha).

Direct sowing

At all four sites at the start of the trial, berries were collected from local *J. communis* bushes (Fungarth, using a local, 4 km distant, population). Seeds were extracted (McCartan & Gosling 2013) and sown in both treatment and control areas. Sowing took place at each site in 20 of the 25 cm x 25 cm quadrats, evenly distributed between treatment and control areas (Table 1; Figure S1). As 100 seeds were sown per quadrat, each site received two thousand seeds. Viability of seeds sown was estimated based on a sample of seeds which was tested using the tetrazolium test (Gosling 2003).

Germination and regeneration assessments

J. communis seed germination was monitored annually in all the sown plots. Natural regeneration of *J. communis* was also monitored annually but due to time limitations, this was conducted by searching a 10 m wide buffer around a sample of female bushes (20 bushes per site - 5 bushes per block at Dorback and Pentland Hills, 10 bushes in the treatment area, 10 in the control area at Fungarth and Ballyoukan). a systematic search of the sites for *J. communis* seedlings and young plants was conducted in the final year of monitoring, at the two sites with cattle grazing only. Plant height and root collar diameter were recorded, together with a description of the ground vegetation in a 1 m x 1 m square area around each plant. Species cover was assessed in different vegetation layers to give a total cover value for a quadrat; this value could therefore exceed 100%.

Vegetation monitoring

An even number of permanent 25 cm x 25 cm quadrats were randomly located in the treatment and control areas at each site. At Fungarth and Ballyoukan, sampling was further stratified by vegetation type (Table S1). In the annual assessments, species cover (as well as cover of bare ground and litter) was assessed in different vegetation layers to give a total cover value for a quadrat; this value could therefore exceed 100%.

Data analysis

In order to relate regeneration to management treatment the year when *J. communis* seedlings germinated needs to be known.

This could be recorded for those seedlings appearing during the trial period in the plots and the monitored buffer around the bushes. However, for the seedlings found outside these areas (during the systematic search), year of germination had to be estimated by dividing seedling root collar diameter by annual diameter stem increment figures. These figures were from a *J. communis* dendrochronology study of the early growth of bushes at a comparable site to our study sites (Glen Artney, Perthshire, Scotland; A. Tene 2006, Forest Scientist, Forest Research, Rosin, Midlothian, personal communication).

In order to identify vegetation characteristics of microsites suitable for regeneration, comparisons between vegetation composition and cover for quadrats containing *J. communis* seedlings and quadrats (a random sample of the 2010 permanent quadrats) where no seedlings were found were made using Principal Component Analysis. As vegetation cover and composition is expected to

change rapidly following disturbance, quadrats with the youngest (1 to 2 year old) *J. communis* seedlings only were included. These seedlings germinated in 2010 or 2011. For the general linear models and after inspection of data, root collar diameter (the response variable and proxy for seedling age) was log transformed to stabilise data dispersion, and percentages of plant cover (moss, herbs, grass, *Pteridium aquilinum*), were checked for co-linearity. Plant cover terms that were correlated were tested in separate models as potential explanatory variables. A gaussian error structure was followed. Automated model simplification using Akaike's information criterion (AIC) was applied to find the minimal adequate model with the greatest fit, and residuals were examined for normality (Crawley, 2005).

For the investigation of intervention and site condition measures associated with regeneration microsites, we used ground vegetation data from annual monitoring to describe three site condition measures: vegetation height, occurrence of exposed bare ground (this we defined as >80% bare ground and < 20% for the sum of grass, herbs and moss) and the occurrence of exposed moss cover (>80% moss cover and < 20% for the sum of grass, herbs and bare ground), i.e. 'exposed' describes conditions where bare ground or moss cover is not shaded or over stood by other ground vegetation. For the linear mixed effects model incorporating random block and quadrat effects, and fixed treatment effect, vegetation height data from Dorback and Pentland Hills was square root transformed to stabilise data dispersion; examination of residuals suggested no further data transformation was required. Due to the failure of model convergence for the response of bare ground to treatment, data have been presented descriptively. There were too few data for occurrence of exposed moss in response to treatment to be tested. Due to lack of within-site replication of grazing treatments applied at Ballyoukan and Fungarth, results can only be descriptive, and means and 95% confidence intervals of the three site condition measures have been presented per treatment. All errors reported are standard errors.




Figure S1: Generic design for trial layout at the two Scottish trial sites receiving scarification treatment (Dorback and Pentland Hills) and two Scottish trial sites receiving cattle grazing treatment (Fungarth and Ballyoukan), indicating blocking (I- IV) or site stratification by vegetation type (V1; V2), and replication of treatments (scarification or cattle grazing (T); control (C)).

References for Appendix S1

Borders Forest Trust (1997) Common juniper (*Juniperus communis* L.): a review of biology and status in the Scottish Borders.

Borders Forest Trust Occasional Paper [No. 1], Ancrum, Scottish Borders, UK.

British Geological Survey (1979) Solid geology map of Scotland, northern England, Isle of Man and Northern Ireland — 1:625 000 third edition. .Keyworth, Nottingham, British Geological Survey.

British Geological Survey (1928) Sheet 32- Edinburgh. Geological survey of Scotland,.1:63,360 geological map series. Keyworth, Nottingham, British Geological Survey.

Chesterton C (2006) Revised Calculation of Livestock Units for Higher Level Stewardship Agreements. Rural Development Service Technical Advice Note 33 (second edition), Rural Development Service, UK.

Crawley MJ. 2005. Statistics: An introduction to using R. John Wiley & Sons Ltd., Chichester.

Gosling PG (2003) Chapter 24: Seed viability testing. Pages 445-481 In: Smith RD, Dickie JB, Linington SH, Pritchard, HW, Probert RJ (eds) Seed conservation: turning science into practice. The Royal Botanic Gardens, Kew, London, UK

GraphPad (2009), GraphPad Software version 3.10, 32 bit for Windows SanDiego, California, USA.

Hill MO, Mountford JO, Roy DB, Bunce RGH (1999) Ellenberg's indicator values for British plants. ECOFACT Volume 2 Technical Annex, Institute of Terrestrial Ecology, Huntingdon

- Humphrey JW, Watts K, Fuentes-Montemayor E, Macgregor NA, Peace AJ, Park KJ (2015) What can studies of woodland fragmentation and creation tell us about ecological networks? A literature review and synthesis. *Landscape Ecology* 30:21–50
- Koricheva J, Gurevitch J (2013) Place of meta-analysis among other methods of research synthesis. Pages 3-13. In: Koricheva J, Gurevitch J, Mengersen K (eds) *Handbook of Meta-Analysis in Ecology and Evolution*. Princeton University Press, Princeton and Oxford.
- McCartan SA, Gosling PG (2013) Guidelines for Seed Collection and Stratification of Common Juniper (*Juniperus communis* L.) Tree Planters Notes 56: 24-29
- Pollock ML, Milner JM, Waterhouse A, Holland JP, Legg CJ (2005) Impacts of livestock in regenerating upland birch woodlands in Scotland. *Biological Conservation* 123: 443-452
- Pyatt DG, Ray D, Fletcher J (2001) *An Ecological Site Classification for Forestry in Great Britain*. Forestry Commission Bulletin 124, Forestry Commission, Edinburgh
- Rodwell JS (1991) *British plant communities, I: Woodlands and Scrub*. Cambridge University Press, Cambridge, UK
- Rodwell JS (1998a) *British plant communities, vol. 2: Mires and heaths*. Cambridge University Press, Cambridge, UK
- Rodwell JS (1998b) *British plant communities, vol. 3: Grassland and montane communities*. Cambridge University Press, Cambridge, UK
- Sullivan G (2003) *Extent and condition of juniper scrub in Scotland*. Scottish Natural Heritage [Contract No. BAT/AC205/01/02/96], Edinburgh, UK.
- R Development Core Team (2011) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S2. *Juniperus communis* regeneration surveys conducted within the Atlantic North and the Atlantic Central environmental zones of Europe.

Citation/ Source	Location and context	Habitat type	Duration	Density (D)/ Frequency (F) of regeneration	Quality of description for:		Conditions associated with regeneration	Conditions not associated with regeneration
					juniper populations ¹	site conditions ²		
Ward 1973	UK- southern England Survey sites selected on historical records of <i>J. communis</i> presence.	Lowland calcareous grassland/ scrub (NVC ³ type CG3, CG4, W21).	Four years (1968 to 1971).	F= 28% (in 86 of 309 1km squares)	Medium	High (+)	- bare ground (reduced competition from other vegetation; no intensive shading) - release from rabbit pressure - shallow soils (on steep chalk slopes, quarries and old track-ways).	
Ward & King 2006.	UK- southern England (county of Sussex). Re- survey (after 30 years) of <i>J.</i> <i>communis</i> sites.	Lowland calcareous grassland/ scrub (NVC ³ type CG3, CG4 and W21).	Three years (2001-2003).	F =14% (in 3 of 22 populations)	Medium	Low (+)	- sparse open grassland - release from sheep and cattle grazing.	- summer drought - rabbit browsing.
Woodruffe, et al. 2016	UK-southern England (counties of Hampshire & Wiltshire)	Lowland calcareous grassland/ scrub (NVC ³ type CG3, CG4 and W21).	Three years (2014-2016).	D =160 ha ⁻¹ (cumulative seedling count)	Medium	Low (-)	- fruiting mature junipers, - short grass, moss and bare areas - rabbit grazing	- severe rabbit grazing, - long grass. - survival less near parent plant
Clifton et al. 1995.	UK- northern England (county of Northumbria) Surveys in two different years- in 1973, 130 sites; 1994 a subset of 83 sites (those extant in 1973 plus new records).	Upland acid grassland/scrub (NVC ³ type W19)	Two, 1-year surveys (1973 and 1994).	'Very little'	High	High (+)	- fluctuations in grazing - freedom from grazing - wetter conditions (leading to layering of bushes)	
Gilbert 1980.	UK- northern England (Upper	Upland acid grassland/scrub	Observations made over 15	D =1 m ² - 5 m ²	Low	High (+)	- bare ground and thin turf adjacent to mature bushes	- closed vegetation of common bent and heath

	Teesdale) 10km ² where a concentration of <i>J. communis</i> sites	(NVC ³ type– W19)	years (1960's & 70's).				- disturbance of ground vegetation following clearing birch scrub - pulses of heavy stock grazing.	bedstraw protection from sheep but not rabbits.
Douglas 2015	UK- northern England (county of Cumbria) Census of upland <i>J. communis</i> scrub	Upland acid grassland/scrub /heathland (NVC ³ – W19, H15)	Four years (2011-2014)	F =33% (in 82 of the 252 sites)	Low	Low (+)		
Long & Williams 2007	UK- upland areas of Britain. Survey questionnaires completed by members of the public	Grassland, moorland, broadleaved woodland/scrub, montane, conifer woodland.	On year (2004-2005)	F = 13% (in 43 of 342 sites)	High	Low (+)	- large population sizes (seedlings recorded on 11 of the 203 sites with <50 bushes and 17 of the 42 sites with 50+ bushes)	- no significant associations noted for seedling presence/absence and: habitat type; grazing animals; rabbits)
Sullivan 2003	UK- Scotland (all). Sample survey stratified geographically.	Upland acid grassland/ scrub, montane heathland, Scots pine woodland and upland oak/birch woodland (NVC ³ types– W19, H15, W18 &W11)	Two years (2001 to 2002).	F =28% (in 21 of 76 sites)	High	High (+)	- large population sizes (usually 50+ bushes) - extensive land use - low nitrogen availability and soil pH ⁴ - soil moisture availability ⁵	- stock grazing when on productive/ lowland sites, - higher levels of base and nitrogen enrichment ⁶ . - extremes of soil moisture availability ⁷ .
Mearns 2001.	UK-southern Scotland (region of Dumfries and Galloway) Survey covered c. 3700 km ² .	Upland acid grassland/scrub and montane heathland (NVC ³ types– W19 &H15).	Two years (1998 to 2000).	F = 5% (in 9 of the 189 populations)	Medium	Medium(-)	larger population sizes.	
Borders Forest Trust 1997.	Survey of <i>J. communis</i> sites in Scotland with historical records or where local knowledge indicated <i>J. communis</i>	Upland acid grassland/scrub (NVC ³ type– W19).	One year (1997)	F = 29% (in 19 of the 65 populations)	Medium	High (+)	- larger colony size - management regime of grazing and shooting - old, abandoned sheep tracks - fenced areas with disturbed sheep hefts - tall, ungrazed and unburnt heather	- population age structure

	presence.						<ul style="list-style-type: none"> - light summer grazing with stock - exclusion of rabbits 	
Cooper et al. 2012.	Ireland - 11 counties on Atlantic coast	Five habitat types ⁸ containing <i>J. communis</i> identified: 1- Wet grass/heath/ bog 2- Exposed calcareous rock 3- Dry calcareous heath & grassland 4- Dry siliceous heath 5-Dry calcareous /neutral grassland.	Three years (2008 – 2010).	F = 18% (in 22 of 125 sites) Average % seedlings by habitat type: - 0.5% (habitat type 1); 0.5% (2); 1.2% (3); 3.5% (4); 5.9% (5) (calculated from sites where ≥ 50 bushes, n=45) Significantly higher % seedlings for habitat types 4 and 5.	High	Medium(+)	<ul style="list-style-type: none"> - large parent populations, - high bush density, and presence of berries - relatively lower nitrogen levels⁹ or more calcareous sites (pH = 7.4 versus pH = 6.8) - rocky crevices 	<ul style="list-style-type: none"> - intensive grazing pressure - relatively nitrogen rich sites¹⁰
Stolz 2010.	Netherlands- province of Drenthe (Dwingelderveld National Park) (3700 ha)	wet and dry heathlands	One year (2010).	D = 106 seedlings within 'focussed survey area'.	Low	Low (+)	<ul style="list-style-type: none"> - shorter ground vegetation - vegetation dominated with grasses¹¹ - periods of reduced rabbit densities. 	- vegetation dominated with ericaceous shrubs ¹²
Ginkel & Bulten 2007.	Netherlands- province of Drenthe Survey of 25%.of <i>J. communis</i> stands on state, nature conservation organisation and private land.	Heathlands	One year (2005).	D =100 seedlings - at 3 locations of unspecified size	Low	Low (-)	<ul style="list-style-type: none"> - recently grazing - recent removal of shrubs. 	
Vedel 1961	Field survey of <i>J. communis</i> throughout Denmark; information on Swedish	Calcareous and heathland sites	Not stated (assume several years).	'several'	Low	Medium(+) Additional attribute: proximity to Juniper seed source	<ul style="list-style-type: none"> - sparse vegetation cover - bare ground (on heaths, sites with poor sandy soil or bare rock) - deer and sheep grazing creating short vegetation 	

	populations (from literature).						and bare ground - fluctuation in grazing.	
Hommel et al. 2009.	Netherlands-seven provinces (20 nature reserves).	Heathlands	One year (2007).	'A few'	Low	Medium(-)	- shallow litter layer (average depth 1cm) - short, open vegetation (average % cover: dwarf shrubs-7.3, early successional mosses- 46.5) - base enrichment of soil (pH in A and E horizons at 4.8).	- relatively deep litter layer (average depth 1.9cm) - relatively tall, dense vegetation (average % cover: dwarf shrubs-24.4, early successional mosses- 18.5) - relatively acidic soil conditions (pH of 4.3 and 4.4 in A and E horizons, respectively).
Lucassen et al. 2011.	Netherlands-Maasduinen area (11 sites), Guelderland, Overijssel and Drenthe provinces (5 sites) plus 3 reference sites in Germany and Norway.	Heathlands	Four Months (winter 2010/11)	Regeneration occurrence (no, some and strong).	Low, Additional attributes: seed viability ¹³ - berry infestation with mites - foliar (and ripe berry) chemistry	Low (-)	- base saturation of soil mean = 42% - low infestation mites/scale insects (Carulaspis J. communisi) (classed as 0.5) - viability of seeds 35%	- low base saturation of soil mean 23%) - high extractable Al and Al/Ca ratio in soil and plants(at least 2 times higher than at sites where regeneration = 'many') - low P and K concentration and high N/P ratios - high infestation with mites (classed as >2) - low viability of seeds (15%)
Miles & Kinnaird 1979.	Observation of field seed sowing experiment and establishment and regeneration in Scottish Highlands.	pine/birch communities (NVC ³ type-W17/18)	At least five years (1970's).	Not stated.	Low Additional attributes: - survival of germinating seed -establishment of bushes.	Low (+)	- bare ground and short turf adjacent to <i>J. communis</i> stands - protection from rodents (mice eat seed; seedling mortality from small rodents and slugs).	

¹ Quality of description for *J. communis* populations: of the 5 attributes recorded to describe *J. communis* populations (number of bushes, bush condition/size, age structure, regeneration presence, berry presence- 'High' where 4 or 5 attributes recorded, 'Medium' where 3 attributes recorded, 'Low' where 1 or 2 attributes recorded.

² Quality of description of site conditions: of 10 attributes describing site conditions, 'High' - where 5 or 6 recorded, 'Medium' - 3 or 4 recorded, 'Low' - less than 3 recorded; 10 attributes describing site conditions recorded are geology, altitude, slope, aspect, associated habitat/NVC type, associated vascular plant species, vegetation height, soil type, soil pH/soil chemistry, site features e.g. rock outcrops. +/- = including/excluding information on land use/management.

³ National Vegetation Classification (Rodwell 1991, 1998a, 1998b)

⁴ mean site Hill-Ellenberg R values of 2-3, 'N' values of 1.5-2.5 (Hill et al. 1999); soil nutrient regime = 'Very Poor' (Pyatt et al. 2001)).

⁵ mean site Hill-Ellenberg F values of 3-5 (Hill et al. 1999); soil moisture regime = 'Dry' to 'Fresh' (Pyatt et al. 2001)).

⁶ mean site Hill-Ellenberg R values of 2-6, 'N' values of 2-5; soil nutrient regime = 'Very Poor' to 'Rich'.

⁷ mean site Hill-Ellenberg F values of 1-7; soil nutrient regime = 'Very Dry' to 'Very Moist'.

⁸ Fossitt, JA2000.

⁹ mean site Hill-Ellenberg 'N' value of ≤ 2.8 (Hill et al. 1999).

¹⁰ Mean site Hill-Ellenberg 'N' value ≥ 3.2 .

¹¹ mean site Hill-Ellenberg R value of 2.4.

¹² mean site Hill-Ellenberg R value of 1.4.

¹³ Gosling 2003.

References for Appendix S2

Borders Forest Trust (1997) Common juniper (*Juniperus communis* L.): a review of biology and status in the Scottish Borders.

Borders Forest Trust Occasional Paper [No. 1], Ancrum, Scottish Borders, UK.

Clifton SJ, Ranner DS, Ward L (1995) The conservation of juniper in Northumbria. English Nature Research Report [152], Peterborough, UK.

Cooper F, Stone RE, McEvoy P, Wilkins T, Reid N (2012) The conservation status of juniper formations in Ireland. National Parks and Wildlife Service, Department of Environment, Heritage and Local Government [Rep. No. 63], Dublin, Ireland

Douglas M (2015) Report on the 'Uplands for Juniper Project' Cumbria Wildlife Trust, Kendal, Cumbria, UK. URL:

<http://www.cbdc.org.uk/data/projects/Juniper-Project> (accessed 22 March 2016)

Fossitt JA (2000) A guide to the habitats in Ireland. Republic of Ireland. Kilkenny: Heritage Council

Gilbert OL (1980) Juniper in Upper Teesdale. *Journal of Ecology* 68: 1013 - 1024

Ginkel J, Bulten M (2007) Management helps regeneration of juniper. *Vakblad Natuur Bos Landschap* 4: 18-19

Gosling PG (2003) Chapter 24: Seed viability testing. Pages 445-481 In: Smith RD, Dickie JB, Linington SH, Pritchard, HW, Probert RJ (eds) Seed conservation: turning science into practice. The Royal Botanic Gardens, Kew, London, UK

Hill MO, Mountford JO, Roy DB, Bunce RGH (1999) Ellenberg's indicator values for British plants. ECOFACT Volume 2 Technical Annex, Institute of Terrestrial Ecology, Huntingdon

Hommel P, Griek M, Haveman R, den Ouden J, deWaal RW (2009) Herstel van jeneverbestruwelen. *De Levende Natuur* 110: 135-137

Long D, Williams J, Juniper in the British uplands : the Plantlife Juniper Survey results survey results. Plantlife Scotland, Stirling, UK.

Lucassen E, Loeffen L, Popma J, Verbaarschot E, Remke E, de Kort S, Roelofs J (2011) Bodemverzuring lijkt een sleutelrol te spelen in het verstoorde verjongingsproces van Jeneverbes. *De Levende Natuur* 112: 235- 239

Mearns R (2001) Juniper, *Juniperus communis*, in Dumfries and Galloway. The Transactions of the Dumfriesshire and Galloway Natural History and Antiquarian Society 75:1-28

Miles J, Kinnaid JW (1979) Establishment and Regeneration of Birch, Juniper and Scots Pine in the Scottish Highlands. *Scottish Forestry* 33: 102 -119

Pyatt DG, Ray D, Fletcher J (2001) An Ecological Site Classification for Forestry in Great Britain. Forestry Commission Bulletin 124, Forestry Commission, Edinburgh

Stolz T (2010) Impact of Rabbits on Juniper Recruitment in Western Europe. M.Sc. thesis, University of Utrecht, Netherlands.

Sullivan G (2003) Extent and condition of juniper scrub in Scotland. Scottish Natural Heritage [Contract No. BAT/AC205/01/02/96], Edinburgh, UK.

Rodwell JS (1991) British plant communities, I: Woodlands and Scrub. Cambridge University Press, Cambridge, UK

- Rodwell JS (1998a) British plant communities, vol. 2: Mires and heaths. Cambridge University Press, Cambridge, UK
- Rodwell JS (1998b) British plant communities, vol. 3: Grassland and montane communities. Cambridge University Press, Cambridge, UK
- Vedel H (1961) Natural regeneration in juniper. *Proceedings of the Botanical Society of the British Isles* 4:146-148
- Ward LK (1973) The conservation of juniper. I. Present status of juniper in Southern England. *Journal of Applied Ecology* 10: 165-188
- Ward LK, King M (2006) Decline of juniper in Sussex. *Quarterly Journal of Forestry* 100: 263-272
- Woodruffe, P., Appleyard, A. & Fitzpatrick, S. (2016) Botanical Surveyors, Salisbury, Wiltshire, personal communication of work in progress.

Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S3. *Juniperus communis* management studies conducted within the Atlantic North and the Atlantic Central environmental zones of Europe.

Citation/ source	Location/context	Interventions	Duration	Observations made	Response/outcomes/findings	Limitations
Fitter & Jennings 1975.	UK-southern England (Aston Rowant). Chalk grassland with <i>J. communis</i> (NVC1 type W21) Previously burnt & grazed.	Sheep grazing: - rate of 1.2 sheep per ha - timing: autumn, winter, summer - control with no grazing. Sites rabbit fenced prior to treatment.	Seven years	Seedling: - survival - height - crown diameter - stem diameter.	Seedlings regenerate in presence of summer grazed sheep. Growth not hindered by grazing in summer as it is by grazing at other times of the year.	Summer grazing regime assessed for only 2 years as rabbit fencing around treatment failed. Unreplicated within site, only conducted at one site.
Morris et al. 1993.	UK – southern England (Old Winchester Hill). Chalk grassland with <i>J. communis</i> (NVC1 type W21)	Sheep grazing: -rate to remove 75–100% of the herbage per grazing period - timing: spring, summer, autumn -rotated to provide periods of grazing (5years) and no grazing (4 years) different paddocks treated in different years	Twelve years	Seedling: - year of germination - height - survival Established bush height, survival Vegetation height	Regeneration occurs under rotational grazing management (e.g. maximum 60 seedlings/ha/year). More seedlings where: - close to female bushes - short sward height - in period when grazing ceases. Growth and survival of seedlings and young bushes (<10years old) reduced by period of grazing (e.g. 85% lost after grazing period).	Errors likely in seedling number by year estimates, as difficult to: -detect one year old seedlings -determine seedling age. Results difficult to analyse: too few seedlings recording for some treatments individual protection not consistently provided for all seedlings in all years.
Wilkins 2011.	UK- southern England. <i>J. communis</i> scrub sites (10) on chalk soils (NVC ¹ type W21)	Turf stripping to produce bare ground (1m x 1m) scrapes next to <i>J. communis</i> bushes: - protection (vole, rabbit and larger herbivores) with wire mesh cages (T) - unprotected control (C) Scrapes next to male <i>J. communis</i> bushes sown with cleaned seed, female-bush scrapes unsown.	Three years	Seedling: - number recruited - location	Regeneration at four sites: - two control sites (1 plus 2 seedlings in scrapes) and two Treatment sites (1 plus 3 in caged scrapes). - occurred in third year (trial started in Autumn 2008, seedlings recorded summer 2011).	- Not a balanced design: local control only installed at some sites; some sites with treatment or control plots, only.
Kerr 1968. Sykes 1976 (seedling data)	UK-southern Scotland (Tynron Juniper wood) <i>J. communis</i> scrub (5	Interventions for regeneration: -stock fencing -rabbit control	Twelve years (1955 to	Seedling (from surveys of reserve): - number	Fire can successfully prepare ground for <i>J. communis</i> establishment. More recruitment of seedlings when	Not a replicated design for treatments Short duration of monitoring

Citation/ source	Location/context	Interventions	Duration	Observations made	Response/outcomes/findings	Limitations
analysis).	ha) in upland acid grassland/scrub vegetation community. (NVC1 type W19). Re-analysis of seedling data in 1976.	-bracken cutting -providing bird perches - burning (1 ha, unplanned) Establishment treatments: - weeding, plastic mulching, caging, planting and sowing.	1967). Fourteen years (1960 to 1974) for seedling survey.	- height - survival Seedling (when caged /uncaged): - survival - growth Plant (under different establishment treatments): -survival	rabbit numbers are lowered.	for most treatments Some interventions were small scale e.g. 2, 1m ² areas sown with seed, once. For seedling data analysis (1976): Seedling locations not mapped so can't relate to treatments, seedbed conditions or conditions favouring survival/growth-
Clifton et al. 1995. Sutherland 1993.	UK- northern England (Upper Teesdale) Upland acid grassland/scrub habitat (NVC1 type W19).	Three J. communis bush treatments: - coppiced to ground level - coppiced to 1m - removed by dragging/winning. Sites stock and rabbit fenced prior to treatment.	Ongoing – report end third year (1990)	Seedling: - number - location Local site conditions.	Regeneration occurred: when herbivores excluded + dragging bushes (caused ground disturbance and shedding of berries). in 3 years following treatment most in areas shaded by bracken. Survival only where protection in winter from sheep and rabbits.	Presence/absence of control not confirmed. Details of monitoring not given. Monitoring of short duration compared to the known germination profile of sown berries.
Verheyen et al. 2005.	Belgium- province of Limburg (Heiderbos nature reserve). J. communis scrub (10 ha) on dry, heathland site. Regeneration of extant population thought due to abandonment of traditional heathland management 50 years ago. Site ungrazed for last 50 years.	Four treatments - sod cutting + selective herbicide treatment of grasses - cultivating + Calluna vulgaris sown following year - sod cutting All plots mown in at least one year; woody plants cut regularly.	Fourteen years	Demographic change in J. communis population since treatment (23 years later) based on bush: - location - height - stem girth Growth response models used to determine if emergence of new J. communis bushes correlated with management treatments.	Management did not produce younger cohort of J. communis: recruitment (estimated at 5 individuals per ha per year):not enhanced: established bush mortality promoted. conditions created by cultivation less suitable for survival of regenerating/ young bushes than other treatment e.g. sod cutting. Limited availability of bare ground for germination and low viability of seeds, may explain lack of success.	Regeneration interpreted from long term (20years) survival of plants (not annual monitoring). Different treatments applied in different time periods; could be an undetermined treatment *year interaction. Seeds thought to have very low viability. Factors (other than management), negative for regeneration acting during trial period but not during period when extant population established (e.g. lowering water table and nitrogen deposition).
Hommel et al. 2012 (in Dutch).	Netherlands (2 sites)- province of Drenthe (Balingertzand) &	Treatments(Trts): - control /no management (Trt 1)	Four years	Germination trial: Seedling emergence (monitored 2-3	Germination trial: germination capacity low (0.03% average; 0.28% maximum).	Very low seedling numbers makes detection of positive influences on germination

Citation/ source	Location/context	Interventions	Duration	Observations made	Response/outcomes/findings	Limitations
	Overijssel (De Borkeld) J. communis scrub on dry heathland. Germination trial (seed sown in enclosed plots) and natural regeneration trial (within the J. communis stand) repeated at both sites.	<ul style="list-style-type: none"> - shallow sod cutting (litter and vegetation removed) (Trt 2) - deep sod cutting (8cm depth , organic topsoil removed) (Trt 3) - deep sod cutting + liming (rate of 200g/m²) (Trt 4) spading (to mix soil) (Trt 5) adding of J. communis litter (Trt 6) <p>Germination trial: 1 block of 25, 1m x 1m plots with 0.5m buffers. Four reps of each Trt. Each plot split for sowings in March 2008 (1000 berries, 3 origins, per plot) and 2009 (800, 1). Over 300,000 seeds sown; four origins used.</p> <p>Natural regeneration trial: Trts 2, 3 and 4 (5 reps of each), applied along a transect of 15, 1m x1m Trt plots; 5 Trt 1 plots located adjacent to Trt 2 plots.</p>		<p>times annually)</p> <p>Natural regeneration trial:</p> <ul style="list-style-type: none"> - seedling emergence - vegetation development & inventory of species and plant communities associated with J. communis. <p>Soil composition and chemistry for each trial site(0 -10cm, 10-20 cm):</p> <ul style="list-style-type: none"> -clay content - organic matter content - pH - nitrogen, - phosphorous - base ion saturation. 	<p>treatment effect for all but Trt 6;Trt 3, 4 and 5 had most germination (but did not differ significantly); no germination with Trt 1 (control). seed origin effect not significant</p> <p>location effect (germination capacity higher at Markelo). Soil influence: more germination where higher: clay content, organic matter content, phosphorus and base ion availability, but lower where higher calcium utilization.</p> <p>Regeneration trial: none recorded.</p> <p>Vegetation survey suggests indicator species (mosses and liverworts) of past regeneration microsites.</p> <p>Indicator species infrequent; not increased by Trts; plots colonised by other but frequent mosses.</p>	hard to detect.

¹National Vegetation Classification (Rodwell, 1991).

References for Appendix S3

- Clifton SJ, Ranner DS, Ward L (1995) The conservation of juniper in Northumbria. English Nature Research Report [152], Peterborough, UK
- Fitter AH, Jennings RD (1975) The effects of sheep grazing on the growth and survival of seedling junipers (*Juniperus communis*). *Journal of Applied Ecology* 12: 637-642
- Hommel P, Haveman R, Huiskes HPJ, deWaal RW (2012) Herstel van jeneverbesstruwelen Resultaten OBN-onderzoek 2007 - 2011. Alterra internal report, Wageningen, Netherlands
- Kerr AJ (1968) Tyron Juniper Wood National Nature Reserve. The Nature Conservancy Council, Edinburgh, UK
- Morris MG, Clarke RT, Rispin WE, Ward LK, Thomas JA (1993) Invertebrate studies at Old Winchester Hill NNR. Institute of Terrestrial Ecology (Natural Environment Research Council) [EN/NERC contract No F74-05-18], Wareham, UK.
- Sykes JM (1976) Regeneration of juniper. A summary of data collected at Tynron Juniper Wood NNR 1960 -1974. Institute of Terrestrial Ecology Project [387], Grange-over-Sands, UK
- Sutherland WJ (1993) Juniper regeneration -Habitat Management News. *British Wildlife* 4: 194
- Rodwell JS (1991) British plant communities, I: Woodlands and Scrub. Cambridge University Press, Cambridge, UK
- Verheyen K, Schreurs K, Vanholen B, Hermy M (2005) Intensive management fails to promote recruitment in the last large population of *Juniperus communis* (L.) in Flanders (Belgium). *Biological Conservation* 124: 113-121
- Wilkins, T. 2011, Woodland Advisor, Plantlife, Salisbury, Wiltshire, personal communication


Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S4. Loading values (superior to 0.1) for species groups used in the Principal Component Analysis (Figure 1) . The Principal component Analysis investigated the distribution of ground vegetation cover variables at the field trial site, Fungarth (both grazed and ungrazed area), in samples with and without *Juniperus communis* natural regeneration; regeneration indicated by the presence of 1 to 2 year old seedlings.

Species groups	PCA1	PCA2
bracken (<i>Pteridium aquilinum</i>)	0.150	0.734
herbs	-0.337	-0.218
grass	0.497	-0.526
shrubs	-0.321	-0.152
moss	-0.671	-0.112
bare ground	0.165	
litter	0.190	-0.309


Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S5. Change in exposed bare ground over six years at the two trial sites (Dorback (D) and Pentland Hills (P)) subject to initial scarification treatment, in control (C) and scarified (T) areas; 95% confidence intervals shown. Data points for the same year have been offset for ease of viewing.



Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S6. Change in occurrence of exposed bare ground and exposed moss cover over six years at the cattle grazed trial sites (Ballyoukan (B) and Fungarth (F)) in control (C) and grazed (T) areas; 95% confidence intervals shown. Data points for the same year have been offset for ease of viewing.



Broome, A. et al. Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. *Applied Vegetation Science*.

Appendix S7. Change in vegetation height at the two annually cattle grazed sites (Ballyoukan (B) and Fungarth (F)) over six years in control (C) and grazed (T) areas (no grazing in year one); 95% confidence intervals shown. Data points for the same year have been offset for ease of viewing.

