

Accepted refereed manuscript of:

Nightingale TE, Metcalfe R, Vollaard N & Bilzon JLJ (2017) Exercise guidelines to promote cardiometabolic health in spinal cord injured humans: time to raise the intensity?, *Archives of Physical Medicine and Rehabilitation*, 98 (9), pp. 1693-1704.

DOI: [10.1016/j.apmr.2016.12.008](https://doi.org/10.1016/j.apmr.2016.12.008)

© 2017, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Accepted Manuscript

Exercise guidelines to promote cardiometabolic health in spinal cord injured humans: time to raise the intensity?

Tom E. Nightingale, PhD, Richard S. Metcalfe, PhD, Niels BJ. Vollaard, PhD, James L.J. Bilzon, PhD



PII: S0003-9993(17)30004-7

DOI: [10.1016/j.apmr.2016.12.008](https://doi.org/10.1016/j.apmr.2016.12.008)

Reference: YAPMR 56765

To appear in: *ARCHIVES OF PHYSICAL MEDICINE AND REHABILITATION*

Received Date: 4 November 2016

Accepted Date: 15 December 2016

Please cite this article as: Nightingale TE, Metcalfe RS, Vollaard NB, Bilzon JLJ, Exercise guidelines to promote cardiometabolic health in spinal cord injured humans: time to raise the intensity?, *ARCHIVES OF PHYSICAL MEDICINE AND REHABILITATION* (2017), doi: 10.1016/j.apmr.2016.12.008.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Running head:** Exercise intensity in SCI

2

3 **Title:** Exercise guidelines to promote cardiometabolic health in spinal cord injured
4 humans: time to raise the intensity?

5

6 **Authors:** Tom E Nightingale, PhD¹, Richard S Metcalfe, PhD², Niels BJ Vollaard,
7 PhD^{1,3} & James L J Bilzon, PhD¹

8

9 **Affiliations:**

10 ¹ Department for Health, University of Bath, Bath, BA2 7AY, UK

11 ² Sport and Exercise Sciences Research Institute, Ulster University, Northern Ireland

12 ³ Faculty of Health Sciences and Sport, University of Stirling, Scotland

13

14 **Acknowledgement of prior presentation:** None

15

16 **Acknowledgement of financial support:** None

17

18 **Explanation of conflicts of interest:** The authors declare no conflicts of interest

19

20 **Corresponding Author:** Professor James Bilzon, Department for Health, University of
21 Bath, Bath, BA2 7AY, UK

22 Email: J.Bilzon@Bath.ac.uk

23 Tel: +44 (0)1225 384809

24

25 **Clinical trial registration number:** Not applicable

1 **Running head:** Exercise intensity in SCI

2

3 **Title:** Exercise guidelines to promote cardiometabolic health in spinal cord injured

4 humans: time to raise the intensity?

5

6

7

8

9

10

11

12

13

14 **Acknowledgement of prior presentation:** None

15

16 **Acknowledgement of financial support:** None

17

18 **Explanation of conflicts of interest:** The authors declare no conflicts of interest

19

20

21

22

23

24

25 **Clinical trial registration number:** Not applicable

Abstract

Spinal cord injury (SCI) is a life changing event that, as a result of paralysis, negatively influences habitual levels of physical activity and hence cardiometabolic health. Performing regular structured exercise therefore appears extremely important in persons with SCI. However, exercise options are mainly limited to the upper-body, which involves a smaller activated muscle mass compared to the mainly leg-based activities commonly performed by non-disabled individuals. Current exercise guidelines for SCI focus predominantly on relative short durations of moderate-intensity aerobic arm cranking exercise, yet contemporary evidence suggests this is not sufficient to induce meaningful improvements in risk factors for the prevention of cardiometabolic disease in this population. As such, these guidelines and their physiological basis, require reappraisal. In this special communication, we propose that high-intensity interval training (HIIT) may be a viable alternative exercise strategy, to promote vigorous-intensity exercise and prevent cardiometabolic disease in persons with SCI. Supplementing the limited data from SCI cohorts with consistent findings from studies in non-disabled populations, we present strong evidence to suggest that HIIT is superior to moderate-intensity aerobic exercise for improving cardiorespiratory fitness, insulin sensitivity and vascular function. The potential application and safety of HIIT in this population is also discussed. We conclude that increasing exercise intensity could offer a simple, readily available, time-efficient solution to improve cardiometabolic health in persons with SCI. We call for high-quality randomised controlled trials to examine the efficacy and safety of HIIT in this population.

Key words: Spinal cord injury, Cardiometabolic health, High-intensity interval training, Vigorous-intensity exercise, Cardiorespiratory fitness

Abbreviations:

- CVD- cardiovascular disease,
FMD- flow-mediated dilation,
HbA1c- glycated haemoglobin,
HDL-C- high-density lipoprotein cholesterol
HIIT- high-intensity interval training,
HRmax- maximum heart rate,
LDL-C- low-density lipoprotein cholesterol
MICT- moderate-intensity continuous training,
OGTT- oral glucose tolerance test,
PAG-SCI- physical activity guidelines for people with a spinal cord injury,
RPE- rating of perceived exertion,
SCI- spinal cord injury,
SIT- sprint interval training,
T2DM- type-2 diabetes mellitus,
 $\dot{V}O_{2peak}$ - maximal oxygen uptake.

76 1 Introduction

77

78 Spinal cord injury (SCI) creates a complex pathophysiology, characterised by paralysis,
 79 which has wide-ranging implications for multiple body systems. For persons with SCI,
 80 chronic cardiometabolic diseases occur at a heightened frequency and earlier in the
 81 lifespan compared to non-disabled individuals ¹⁻³. Given that more than 2 million people
 82 currently live with SCI worldwide and the incidence of SCI is highest among young
 83 adults ⁴, it is clear that there is an increased and prolonged demand on medical and
 84 support resources for persons aging with paralysis. Despite the known, undisputed
 85 health benefits of physical activity in non-disabled individuals ⁵⁻⁷, research suggests
 86 patients with SCI perform little to no physical activity ⁸⁻¹¹, and this is likely a key driver
 87 of the greater prevalence of cardiometabolic disease in this population ^{12, 13}. Therefore,
 88 it is a priority to develop evidence-based, effective physical activity recommendations
 89 for the prevention of chronic disease in persons with SCI.

90

91 The recently re-published Physical Activity Guidelines for Spinal Cord Injury (PAG-
 92 SCI) recommends at least 20 minutes of moderate to vigorous-intensity aerobic exercise
 93 twice a week (40 min/wk) ¹⁴, while a recent position statement from Exercise and Sports
 94 Science Australia recommends ≥ 150 min/wk of moderate-intensity or ≥ 60 min/wk of
 95 vigorous-intensity exercise ¹⁵. Both of these guidelines also include strength training ≥ 2
 96 day/wk ^{14, 15}. Regardless of the large discrepancy between these guidelines in terms of
 97 the recommended volume of moderate-intensity exercise, they remain indifferent from
 98 the *minimum* amount of exercise which is promoted by reputable, international health
 99 authorities [Centers for Disease Control (CDC) and World Health Organisation (WHO)]
 100 in order to reduce the risk of developing cardiometabolic disease in the general

101 population. However, it is noteworthy that the exercise guidelines for non-disabled
102 populations are based on lower-body or whole-body activity (e.g. walking, running,
103 cycling), whereas exercise for persons with SCI is primarily restricted to the smaller
104 upper-body skeletal muscles [e.g. arm-crank exercise or wheelchair propulsion]. As a
105 result of the smaller active muscle mass and blunted haemodynamic responses with
106 SCI, the absolute capacity for physical exercise is reduced¹⁶⁻¹⁸. Therefore, at the same
107 relative intensity, the absolute energy expenditure, cardiovascular strain, and whole-
108 body metabolic demand, will always be lower during moderate-intensity arm-crank
109 exercise or wheelchair propulsion compared with moderate-intensity walking or
110 cycling. The ability for skeletal muscle to adapt to the same stimulus will not be
111 reduced; however, the smaller active muscle mass means that modest training-induced
112 adaptations in the arm are less likely to impact biomarkers of cardiometabolic health. As
113 such, to promote a *lower* volume of exercise in this population would seem
114 physiologically counterintuitive, whilst promoting a *similar* volume of exercise would
115 likely be less effective. In accordance with this, a recent randomised controlled trial
116 demonstrated that performing PAG-SCI for 16 weeks was insufficient to promote
117 clinically meaningful changes in both novel and traditional biomarkers of
118 cardiovascular disease (CVD)¹⁹. Moreover, a systematic review requested by the
119 Consortium for Spinal Cord Medicine²⁰ concluded that the current evidence is
120 insufficient to determine whether these volumes of exercise are associated with positive
121 changes in carbohydrate and lipid metabolism (and associated disorders) amongst adults
122 with SCI. Therefore, we contend that these guidelines, and their physiological
123 justification, require reappraisal, and that there is need to develop more effective,
124 alternative approaches.

125

126 There are numerous psychosocial and environmental barriers to engage in physical
127 activity for individuals that use wheelchairs²¹⁻²⁴. Moreover, compromised venous
128 return in persons with SCI blunts cardiac output²⁵, which can lead to an early onset of
129 muscle fatigue²⁶, thus reducing ones capacity for prolonged exercise. Therefore,
130 promoting a *larger* volume of moderate-intensity exercise might not be feasible in this
131 population. Functional electronic stimulation²⁷⁻³⁰ and body weight supported treadmill
132 training³¹, have received considerable research attention, but have numerous practical
133 limitations (i.e. significant cost and specialist resources required), and may have limited
134 application outside the laboratory. One potential alternative approach, which has
135 received less attention, would be to recommend high-intensity interval training (HIIT)
136 as a practical means of increasing vigorous-intensity exercise. The benefit of vigorous-
137 intensity physical activity is supported by a number of epidemiological studies, albeit in
138 non-disabled individuals, demonstrating superior reductions in the risk of
139 cardiovascular^{32, 33} and all-cause mortality³⁴⁻³⁶, in comparison to light-to-moderate
140 intensity physical activity. Moreover, accumulating evidence, from studies applying
141 HIIT in non-disabled populations, demonstrates that HIIT promotes superior peripheral
142³⁷ and whole-body adaptations³⁸⁻⁴⁰, compared with moderate-intensity continuous
143 training (MICT). HIIT may therefore offer a simple, more effective alternative to
144 current approaches for improving cardiometabolic health in persons with SCI. In the
145 following sections we put forward the case for recommending HIIT in SCI, and
146 subsequently consider its potential practical application and safety in this population.

147

148

149

2 High-intensity Interval Training to Facilitate Vigorous-intensity Exercise in Spinal Cord Injury

HIIT encompasses exercise performed above the intensity which elicits the maximal lactate steady state. Any exercise above this threshold results in the progressive accumulation of intramuscular and systemic metabolites that are implicated in fatigue. As such, exercise intensities above this threshold ($\sim 80\text{--}85\%$ $\dot{V}O_{2\text{peak}}$) cannot be maintained for a prolonged period of time. The exercise must therefore be performed in intervals interspersed with periods of low-intensity or resting recovery. The main justification for HIIT is that it allows a greater volume of vigorous-intensity exercise to be accrued in a single exercise session, and accumulating evidence suggests that this can be of great physiological and clinical benefit³⁸⁻⁴⁰.

A wide range of HIIT protocols have been utilised in the literature but with limited standardisation of the terminology used to classify different protocols. Furthermore, studies have prescribed exercise intensities as a percentage of different maximal physiological responses [e.g. maximum heart rate (HR_{max})⁴¹, heart rate reserve⁴², age-predicted max heart rate⁴³ and peak oxygen uptake ($\dot{V}O_{2\text{peak}}$)⁴⁴] and, for these reasons, may not be directly comparable, particularly in individuals with low baseline fitness⁴⁵. Nevertheless, for the purposes of this review, we adopt the terminology proposed by Weston *et al.*³⁸, whereby HIIT describes protocols using intensities between 80-100% of HR_{max} , whereas protocols using ‘all-out’ efforts, or efforts $\geq 100\%$ $\dot{V}O_{2\text{peak}}$, are referred to as sprint interval training (SIT) (Figure 1). There is good evidence that both HIIT and SIT provide equal or even superior physiological adaptations compared with MICT⁴⁶⁻⁵⁰. However, as SIT protocols may be more difficult to adapt in order to

175 provide a practical intervention for persons with SCI, in this review we draw mainly on
 176 HIIT studies to support the argument for vigorous-intensity exercise. Example HIIT
 177 protocols tested in both the SCI and non-disabled literature are described in Table 1.

178

179 *[INSERT FIGURE 1 ABOUT HERE]*

180

181 *[INSERT TABLE 1 ABOUT HERE]*

182

183

184 **3 Moderate vs Vigorous-intensity Exercise for Cardiometabolic Health**

185

186 **3.1 Cardiorespiratory Fitness and Skeletal Muscle Oxidative Capacity**

187

188 Poor cardiorespiratory fitness has been widely reported in individuals with SCI^{52, 53}.

189 Although just ~90 min/wk^{44, 54} of MICT is sufficient to promote modest improvements

190 (~10%) in $\dot{V}O_{2peak}$, a substantially larger volume (180 min/wk) is necessary for greater

191 improvements (~19%)⁵⁵. Vigorous-intensity exercise offers superior benefits and is

192 more time efficient. Of the two studies which have used time-matched training

193 protocols in SCI (Table 2) there are negligible (12% vs. 10%)⁴² and considerable (50%

194 vs. 17%)⁵⁶ improvements in $\dot{V}O_{2peak}$ with vigorous-intensity compared to moderate-

195 intensity exercise, respectively. The larger improvement in the De Groot *et al*,⁵⁶ study

196 could be due to participants having acute (< 225 days) injuries or the greater volume of

197 accumulated vigorous-intensity activity (additional 48 min/wk). More recently,

198 unpublished data from Sæter⁵⁷, which adopted a more robust isocaloric study design,

199 demonstrated a superior stimulus for $\dot{V}O_{2peak}$ and PPO with vigorous-intensity exercise

200 compared to MICT. Furthermore, a case-study demonstrated a 52% increase in $\dot{V}O_{2peak}$
 201 in a 42 year old man with SCI following just 6 weeks of HIIT⁵⁸.
 202
 203 Several studies have directly compared the effects of energy-matched HIIT and MICT
 204 on $\dot{V}O_{2peak}$ in deconditioned (non-disabled) individuals with pre-existing
 205 cardiometabolic disease and these have clearly demonstrated that HIIT results in
 206 superior improvements. These studies were summarised in a recent meta-analysis
 207 which, using data from 10 studies and 273 participants, showed that the increase in
 208 $\dot{V}O_{2peak}$ following HIIT was approximately twice (~ 3 ml/kg/min) that observed
 209 following MICT³⁸. This finding has been reproduced in various non-disabled
 210 populations including healthy young and middle-aged sedentary men^{59,60}, overweight
 211 and obese men and women⁶¹, and in individuals with type-2 diabetes mellitus (T2DM)
 212⁶². A 3 ml/kg/min improvement in cardiorespiratory fitness is associated with a 15% and
 213 19% reduction in all-cause and CVD mortality, respectively, and is on par with a 7 cm
 214 reduction in waist circumference, a 5 mmHg reduction in systolic blood pressure, or a 1
 215 mmol/L drop in fasting plasma glucose^{63,64}. Given that cardiorespiratory fitness
 216 consistently manifests as the strongest predictor of cardiometabolic disease risk and
 217 longevity in epidemiological studies⁶⁵⁻⁶⁸, these findings are an important point of
 218 reference in the argument for applying HIIT, as a model to increase vigorous-intensity
 219 physical activity, in individuals with SCI.
 220
 221 Although still a subject of debate⁶⁹⁻⁷¹, recent evidence supports, at least partially, the
 222 role of peripheral muscle characteristics, in particular absolute mitochondrial capacity
 223 (i.e. maximal mitochondrial oxygen utilization), in limiting $\dot{V}O_{2peak}$, and hence
 224 underpinning changes in $\dot{V}O_{2peak}$ with exercise training^{72,73}. It is noteworthy then that a

recent study convincingly demonstrated that cycling based HIIT induced superior mitochondrial adaptations compared with MICT, in muscle taken from the lower limb³⁷. Arm exercise training may not be sufficient to induce central hemodynamic adaptations⁷⁴, but can be expected to induce peripheral mitochondrial adaptations. Thus, if the superior effects observed with HIIT compared with moderate-intensity cycling and walking/running in non-disabled individuals are translatable to arm exercise training in persons with SCI, then HIIT may provide a more effective intervention for improving $\dot{V}O_{2peak}$ in persons with SCI. Moreover, the superior changes in mitochondrial oxidative capacity with HIIT may have implications for other cardiometabolic risk factors such as insulin sensitivity and glycaemic control⁷⁵.

[INSERT TABLE 2 ABOUT HERE]

3.2 Insulin Action and Glycaemic Control

Insulin resistance is a pre-requisite to T2DM. It is characterised by the failure of insulin to exert the normal cellular effects on various tissues, leading to the impairment of insulin mediated glucose disposal. Fasting hyperglycaemia can persist due to the insensitivity of the liver to the suppressive effects of insulin on gluconeogenesis and reduced glycogenolysis⁷⁶. Consequently fasting plasma glucose concentrations have been shown to correlate with basal rates of hepatic glucose output⁷⁷. Therefore, as fasting plasma glucose concentrations tend to be only mildly elevated in individuals with SCI⁷⁸, it is most likely that peripheral insulin resistance is the major driver behind impaired glycaemic control in this population. The lack of stimulation and disuse because of paralysis can have a profound impact on skeletal muscle below the level of

injury, including i) atrophy of lean mass⁷⁹⁻⁸², which diminishes the tissue available for glucose disposal (Figure 2a)^{83, 84}, and ii) accumulation of intramuscular fat^{85, 86}.

[INSERT FIGURE 2 ABOUT HERE]

Recent publications have demonstrated that moderate-intensity arm-crank ergometry improves insulin resistance, as determined by HOMA-IR^{87, 88}. Although this is promising, HOMA-IR reflects hepatic insulin sensitivity, whereas indices derived during postprandial oral glucose tolerance tests (OGTT), such as the $ISI_{matsuda}$, represent predominantly peripheral insulin sensitivity^{89, 90}. Data from the HOMEX-SCI trial, including both fasting and provocative dynamic testing, would suggest arm-crank MICT (60 – 65% $\dot{V}O_{2peak}$, 180 min/wk) in persons with chronic paraplegia improves hepatic but not whole-body insulin sensitivity⁵⁵. Therefore, moderate-intensity arm-crank exercise might not be sufficient to overcome insulin resistance in peripheral tissues. There is a paucity of research comparing both fasting and dynamic glucose and insulin responses to HIIT or MICT in the context of arm-crank exercise in the SCI population. Insulin sensitivity data from De Groot *et al*⁵⁶ is counter-intuitive, in that it demonstrates non-significant improvements in the moderate-intensity group and reduced insulin sensitivity in the high-intensity group. This may be explained by a natural regression to the mean effect (i.e. greater proportion of insulin resistant individuals in the low-intensity group at baseline). These results should be viewed with caution due to the, (i) small sample size (n=3 per group) and, (ii) the marked age and sex differences between the two groups, which could impact exercise responses.

275 The impact of HIIT on insulin action and glycaemic control in non-disabled populations
 276 has recently been summarised by Jelleyman *et al*,⁴⁰ in a meta-analysis of 50 training
 277 studies. Their analyses demonstrated that HIIT was associated with improved insulin
 278 sensitivity (estimated via fasting or OGTT-derived indices) and reduced fasting glucose
 279 when compared to both baseline and/or changes in a no-exercise control group⁴⁰. The
 280 magnitude of change appeared to be greater in populations with insulin resistance (e.g.
 281 T2DM or metabolic syndrome) with reductions in glycated haemoglobin (HbA1c) also
 282 observed in this group⁴⁰. When compared with MICT there appeared to be greater
 283 improvements in markers of insulin sensitivity with HIIT (both fasting and dynamic
 284 combined), but no difference in the change in fasting glucose, insulin or HbA1c in
 285 isolation⁴⁰. These differences were apparent despite the fact that the methods varied
 286 considerably between studies. This included variations in the HIIT protocols utilised
 287 (e.g. SIT vs HIIT, cycling vs running), the techniques used to assess insulin sensitivity
 288 (e.g. fasting vs OGTT vs clamp) and the duration after the final training session in
 289 which the insulin sensitivity data was captured. Moreover, studies had been performed
 290 in a wide variety of populations. As such, there is sufficient evidence that in non-
 291 disabled populations with insulin resistance HIIT is associated with superior changes in
 292 markers of insulin sensitivity compared to MICT^{62, 91-95}.

293
 294 It is also important to consider the acute effects of MICT and HIIT on glycaemic
 295 control, although this has received less research attention, especially in SCI individuals.
 296 Two studies have examined the acute effects of HIIT vs MICT on glycaemic control,
 297 using continuous glucose monitors to capture 24-hour glucose profiles, and have shown
 298 superior effects with HIIT in both obese men⁹⁶ and individuals with T2DM⁹⁷. These
 299 effects are underpinned by a plausible mechanism given that high-intensity exercise is

300 associated with greater muscle glycogen utilisation⁹⁸ and muscle glycogen
 301 concentrations are an important driver of acute changes in insulin sensitivity with
 302 exercise⁹⁹⁻¹⁰¹. Clearly, the acute effects of exercise, as well as comparisons of HIIT and
 303 MICT, on glycaemic control in SCI individuals, is an important area of future research.

304

305

306 **3.3 Vascular Function and Blood Pressure**

307

308 Arterial stiffness¹⁰² and endothelial function^{103, 104} are important predictors of future
 309 cardiovascular health. Individuals with SCI are characterised by severe deterioration of
 310 structure and function of vessels below the level of injury¹⁰⁵, but evidence also suggests
 311 increased stiffness and impaired endothelial function within central and regional upper
 312 body arteries in SCI relative to non-disabled controls¹⁰⁶. Recent evidence suggests that
 313 achieving the PAG-SCI for 16-weeks is insufficient to improve the health of both lower
 314 and upper-limb, as well as central blood vessels¹⁹.

315

316 A recent meta-analysis, including 182 participants from 7 studies, demonstrated that
 317 HIIT was superior to MICT for improving markers of endothelial function³⁹. Within
 318 the meta-analysis, studies that had utilised a work-matched HIIT protocol, consisting of
 319 4 x 4 min at 85-90% HR_{max}, appeared to show the most consistent benefit of HIIT over
 320 and above improvements observed with MICT^{61, 91, 107, 108}. A 1% increase in FMD
 321 (flow-mediated dilation) is associated with a 13% reduction in the risk of cardiovascular
 322 events¹⁰³. Therefore the 2.6% magnitude of difference in the change in FMD observed
 323 between HIIT and MICT in this meta-analysis would be expected to result in clinically
 324 meaningful risk reduction³⁹.

325 Individuals with lower-level spinal cord lesions experience similar hypertension issues
 326 as the general population ¹⁰⁹, whereas individuals with higher-level lesions (\geq T6) often
 327 suffer from hypotension ¹¹⁰. A direct comparison of moderate and high-intensity
 328 exercise training on blood pressure is not available in SCI. However, in non-disabled
 329 individuals, evidence suggests that several months of HIIT or MICT are able to induce
 330 comparable changes in both systolic and diastolic blood pressure in a variety of
 331 populations ^{60, 61, 91, 111}.

334 3.4 Body Composition

336 Individuals with SCI demonstrate a greater propensity to accumulate excess body fat
 337 compared to non-disabled populations ^{112, 113}. Furthermore, due to the accelerated loss
 338 of lean mass, the distribution of adipose tissue in SCI also appears to be altered ¹¹⁴,
 339 which would be expected to exert detrimental metabolic effects ¹¹⁵⁻¹¹⁸. It is therefore
 340 important to consider the role physical activity plays in maintaining body composition
 341 and the potential contribution towards a sustained energy deficit to reduce adiposity.
 342 Yet, large additions to weekly total energy expenditure (TEE) through structured
 343 exercise (i.e. on top of baseline physical activity) are required to induce meaningful
 344 reductions in body fat ¹¹⁹. For example, Donnelly *et al.*, ¹²⁰ suggested that a meaningful
 345 body mass reduction requires an exercise energy expenditure in excess of 2000 kcal/wk.
 346 If we extrapolate from exercise data for inactive SCI participants in the HOMEX-SCI
 347 trial ⁵⁵, achieving this would require approximately 448 min/wk of moderate-intensity
 348 arm-crank exercise. Therefore, it is perhaps not surprising that following PAG-SCI for

16 weeks does not induce significant reductions in total and visceral fat mass, although it may be sufficient to reduce the rate of increase in adiposity¹⁹.

351

There is good evidence from non-disabled studies that HIIT can be an effective intervention for promoting positive changes in body composition, including reductions in total body mass^{59, 91, 121-123}, total fat percentage¹²²⁻¹²⁵, total abdominal fat mass^{91, 122-124} and waist circumference^{91, 122, 126}. However, perhaps as expected, studies that have compared energy-matched HIIT and MICT interventions (i.e. both interventions would increase TEE to a similar extent) over several months have demonstrated comparable changes in body composition^{61, 91, 121}. Interestingly, it also appears that HIIT protocols requiring lower exercise volumes (e.g. low-volume HIT or SIT) are associated with similar increases in total 24-hour energy expenditure to 30-50 min of MICT^{127, 128} and can also induce meaningful reductions in total and abdominal fat^{124, 129}, which are comparable to 30-45 min of MICT in overweight/obese individuals¹²³. Increases in leg lean mass have also been observed with cycling based HIIT^{122, 124}, and this has the potential to also translate to the upper-body musculature in patients with SCI. While HIIT does not appear to induce a greater reduction in adiposity than MICT, the reviewed evidence would suggest it is equally as effective, but with a reduction in exercise time commitment.

368

369

3.5 Fasting and Postprandial Dyslipidaemia

371

A recent meta-analysis¹³⁰ highlighted that persons with SCI have a unique lipid profile, primarily characterised by depressed high-density lipoprotein cholesterol (HDL-C). Hooker & Wells⁴² showed a trend for increased (21%) HDL-C and reduced (-15%)

low-density lipoprotein cholesterol (LDL-C) with vigorous but not moderate-intensity exercise over 8 weeks. Other trials specifically in persons with SCI demonstrated no impact of exercise-intensity on lipid profiles. Greater or similar improvements in HDL-C with HIIT compared to MICT have been shown in populations with cardiometabolic disease³⁸ and obese young men¹³¹, respectively. Currently the non-disabled literature is unclear as to whether HIIT offers superior adaptations than MICT for lipid profiles³⁹¹³². However, over 24 weeks O'Donovan *et al*,¹³³ demonstrated high-intensity exercise was more effective in improving lipid profiles than MICT of equal energy cost. It is possible interventions of longer durations are required to determine the true-impact of exercise intensity on lipid profiles.

The two studies which have used time-matched training protocols in SCI demonstrated a decrease in fasting triglyceride concentrations (-19%⁴² and -31%⁵⁶) pre-post with vigorous-intensity exercise, but no change with moderate-intensity exercise training. Elevated fasting triglyceride concentrations have long been associated with CVD^{134, 135}. Despite observing unremarkable concentrations of fasting triglycerides, participants with chronic paraplegia have shown exaggerated postprandial lipaemia^{136, 137}. This exaggerated postprandial lipaemia is an important stimulus for the development of atherosclerosis¹³⁸, and non-fasting triglyceride concentrations has revealed a stronger association with CVD than fasting¹³⁹. As a result of a more sedentary lifestyle, reduced lipoprotein lipase slows postprandial triglyceride extraction from the systemic circulation and the atrophy of leg lean mass limits the ability to metabolise postprandial triglycerides as a fuel source¹⁴⁰. To our knowledge, no studies have been conducted looking at the impact of upper-body exercise on postprandial lipaemia in persons with SCI. However, several studies have examined the effect of an acute bout of HIIT on the

400 postprandial triglyceride response to a high-fat mixed meal in able-bodied individuals.
 401 These were summarised in a systematic review which concluded that an acute bout of
 402 HIIT is similarly effective to MICT for reducing postprandial lipaemia ¹⁴¹.

403

404

405 **4 Cardiovascular Safety of HIIT**

406

407 Concerns have been raised over the safety of HIIT in populations at risk of
 408 cardiometabolic disease and this should be specifically considered with reference to
 409 SCI. Evidence from one recent non-disabled study, which included 5000 patients
 410 undergoing supervised cardiovascular rehabilitation over a 7-year period, suggested that
 411 the rate of adverse cardiovascular events was low with both HIIT and MICT, although
 412 the event rate was higher with HIIT ¹²⁶. Specifically, the study reported an adverse
 413 cardiovascular event rate of 1 per ~23,000 exercise hours during HIIT (2 non-fatal
 414 cardiac arrests) compared with 1 per 129,000 exercise hours during MICT (1 fatal
 415 cardiac arrest) ¹²⁶. However, various HIIT protocols have been used safely in patients
 416 with post infarction heart failure ^{142, 143}, diastolic dysfunction ¹⁴⁴, coronary artery disease
 417 ¹⁴⁵ and atrial fibrillation ¹⁴⁶, while also improving clinical symptoms. A systematic
 418 review of laboratory/hospital based exercise training studies in persons with SCI found
 419 that adverse events were not common and those that occurred were not serious ¹⁴⁷. It
 420 should be noted that the individuals in this review and within the studies mentioned
 421 above were subject to extensive screening, and the cardiovascular safety of HIIT in this
 422 population therefore requires further scientific appraisal. However, when appropriate
 423 pre-participation screening is adopted the risks of adverse events are relatively low and
 424 as previously suggested are '*likely comparable with the variant risks observed in the*
 425 *general population*' ¹⁴⁸. SCI-specific special considerations for exercise, including the

management of autonomic dysreflexia, have been thoroughly addressed elsewhere¹⁵,
¹⁴⁹. It is noteworthy that patients with SCI are usually well-educated regarding the
 symptoms and management of autonomic dysreflexia and there is no reason to speculate
 that the occurrence of this will be increased with HIIT. As with any exercise
 prescription, it would be recommended that individuals consult their clinician prior to
 engaging in such exercise training programmes.

432

433

434 **5 Considerations for the application of HIIT to SCI populations**

435

Individuals with SCI \geq T6 exhibit a blunted cardiovascular response due to an absence
 of cardiac sympathetic innervation¹⁵⁰ and a reduced catecholamine response during
 exercise¹⁵¹. As a result of autonomic dysregulation, HR_{peak} can be as low as 120 b/min.
 Consequently in these individuals it would be difficult to prescribe an appropriate
 exercise intensity using heart rate data. Evidence suggests that ratings of perceived
 exertion (RPE)¹⁵² and a talk test¹⁵³ can be effectively used to control exercise intensity
 in persons with paraplegia. Consequently we advise an RPE \geq 16 and ‘speaking is not
 comfortable’ as appropriate markers of ‘vigorous-intensity’ when performing upper-
 body exercise.

445

The advantage of HIIT is that it enables deconditioned individuals to do a substantial
 amount of work at a relatively high-intensity by incorporating rest periods, which
 reduce local muscular fatigue. Fatigue following an acute 20 minute bout of HIIT in
 patients with chronic fatigue syndrome was not clinically different to moderate-intensity
 continuous exercise of a comparable workload¹⁵⁴. Sensory impairment below the level

451 of injury can increase the risk of pressure sores when performing new activities for
 452 prolonged periods in the same position. Consequently as HIIT can be more time
 453 efficient and incorporates rest periods (ideal for performing regular pressure release)
 454 this could mitigate this risk and prevent skin breakdown.

455

456 Due to a reduced sweating capacity and inability to dilate superficial vasculature ¹⁵⁵,
 457 persons with higher-level injuries have an impaired heat loss during exercise ¹⁵⁶. While
 458 workload is increased with HIIT, possibly resulting in greater heat production, the total
 459 exercise time is less than MICT with recovery periods interspersed throughout.

460 Therefore we have no reason to believe that HIIT would impact core body temperature
 461 more than MICT. Still precautions should be taken when persons with SCI exercise in
 462 hot environments, as they have impaired thermoregulatory function ¹⁵⁷. Furthermore, to
 463 overcome blood pooling in lower extremities, associated with impaired venous return,
 464 an adequate cool down should be performed to prevent post-exercise hypotension.

465 Shoulder overuse injuries and musculoskeletal pain are also common in persons with
 466 SCI ^{158, 159}. While the higher workloads necessary to achieve vigorous-intensity might
 467 further contribute to these conditions, exercise has been proposed as a feasible,
 468 conservative, therapeutic treatment for shoulder pain in persons with SCI ¹⁶⁰.

469

470 Discussions regarding behaviour change and/or maintenance are outside the scope of
 471 this review. However, preliminary evidence would suggest that individuals with pre-
 472 diabetic conditions can adhere to HIIT over the short-term (4 weeks) and do so at a
 473 greater level than MICT ^{123, 161}. Questions have been raised regarding the adherence to
 474 HIIT over the long-term ^{162, 163} but we encourage researchers and practitioners to
 475 develop and evaluate strategies to incorporate HIIT into the everyday lives of persons

476 with SCI. We believe this is possible considering the evidence that non-disabled
477 participants enjoyed HIIT more and were equally as confident to engage in HIIT as they
478 were MICT¹⁶⁴. Reassuringly, unpublished data has also demonstrated persons with SCI
479 experienced greater enjoyment with HIIT and SIT protocols compared to MICT¹⁶⁵.
480 However, medical over protection may limit the prescription of vigorous-intensity
481 exercise rehabilitation in this population. To help overcome this, the safety and efficacy
482 of HIIT, particularly for persons with acute (<1 year) and higher level (\geq T6) SCI would
483 need to be demonstrated by well-controlled longitudinal training studies. This is
484 imperative when vigorous-intensity exercise has the potential to offer significantly
485 greater improvements in certain cardiometabolic outcomes than MICT in a population
486 at increased risk of chronic disease.

487

488

489 **6 Conclusions**

490

491 This special communication presents a case for the utility of HIIT as a strategy to
492 promote vigorous-intensity physical activity and reduce cardiometabolic disease in
493 persons with SCI. Data from SCI cohort studies, albeit collected using suboptimal
494 research designs, seem to agree with consistent findings from studies in the general
495 population that vigorous-intensity is superior to moderate-intensity exercise in
496 improving a variety of cardiometabolic health outcomes. Importantly, these findings can
497 be explained and supported by plausible physiological mechanisms. High-intensity
498 virtual reality arm-exercise is already being investigated in persons with SCI¹⁶⁶ and the
499 National Centre on Health, Physical Activity & Disability (NCHPAD) promote a
500 selection of adapted vigorous-intensity exercise options (e.g. wheelchair burpees).

501 Increasing exercise-intensity could offer a simple, readily available, time-efficient
502 solution to improve cardiometabolic health in persons with SCI. However, until stronger
503 evidence has been collated concerning the safety and efficacy of HIIT in this population
504 this is merely a call to action for researchers in the field and not necessarily an exercise
505 guideline to be prescribed by clinicians.

7 References

1. Devivo MJ, Shewchuk RM, Stover SL, Black KJ, Go BK. A cross-sectional study of the relationship between age and current health-status for persons with spinal-cord injuries. *Paraplegia* 1992;30(12):820-7.
2. Bauman WA, Spungen AM. Disorders of carbohydrate and lipid-metabolism in veterans with paraplegia or quadriplegia - a model of premature aging. *Metab-Clin Exp* 1994;43(6):749-56.
3. Garshick E, Kelley A, Cohen SA, Garrison A, Tun CG, Gagnon D et al. A prospective assessment of mortality in chronic spinal cord injury. *Spinal Cord* 2005;43(7):408-16.
4. Lee BB, Cripps RA, Fitzharris M, Wing PC. The global map for traumatic spinal cord injury epidemiology: update 2011, global incidence rate. *Spinal Cord* 2014;52(2):110-6.
5. Booth FW, Gordon SE, Carlson CJ, Hamilton MT. Waging war on modern chronic diseases: primary prevention through exercise biology. *J Appl Physiol* 2000;88(2):774-87.
6. Kesaniemi YA, Danforth E, Jensen MD, Kopelman PG, Lefebvre P, Reeder BA. Dose-response issues concerning physical activity and health: an evidence-based symposium. *Med Sci Sports Exerc* 2001;33(6):S351-S8.
7. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39(8):1423-34.
8. Washburn RA, Zhu WM, McAuley E, Frogley M, Figoni SF. The physical activity scale for individuals with physical disabilities: Development and evaluation. *Arch Phys Med Rehabil* 2002;83(2):193-200.
9. Ginis KAM, Arbour-Nicitopoulos KP, Latimer AE, Buchholz AC, Bray SR, Craven BC et al. Leisure Time Physical Activity in a Population-Based Sample of People With Spinal Cord Injury Part II: Activity Types, Intensities, and Durations. *Arch Phys Med Rehabil* 2010;91(5):729-33.
10. Tanhoffer RA, Tanhoffer AIP, Raymond J, Hills AP, Davis GM. Exercise, Energy Expenditure, and Body Composition in People With Spinal Cord Injury. *J Phys Act Health* 2014;11(7):1393-400.
11. Tanhoffer RA, Tanhoffer AIP, Raymond J, Johnson NA, Hills AP, Davis GM. Energy Expenditure in Individuals With Spinal Cord Injury Quantified by Doubly Labeled Water and a Multi-Sensor Armband. *J Phys Act Health* 2015;12(2):163-70.
12. Blair SN. Physical inactivity: the biggest public health problem of the 21st century. *Br J Sports Med* 2009;43(1):1-2.
13. Thyfault JP, Krogh-Madsen R. Metabolic disruptions induced by reduced ambulatory activity in free-living humans. *J Appl Physiol* 2011;111(4):1218-24.
14. Evans N, Wingo B, Sasso E, Hicks A, Gorgey AS, Harness E. Exercise Recommendations and Considerations for Persons With Spinal Cord Injury. *Arch Phys Med Rehabil* 2015;96(9):1749-50.
15. Tweedy SM, Beckman EM, Geraghty TJ, Theisen D, Perret C, Harvey LA et al. Exercise and sports science Australia (ESSA) position statement on exercise and spinal cord injury. *Journal of science and medicine in sport / Sports Medicine Australia* 2016.
16. Theisen D. Cardiovascular determinants of exercise capacity in the Paralympic athlete with spinal cord injury. *Exp Physiol* 2012;97(3):319-24.

17. Jehl JL, Gandmontagne M, Pastene G, Eyssette M, Flandrois R, Coudert J. Cardiac output during exercise in paraplegic subjects. *Eur J Appl Physiol Occup Physiol* 1991;62(4):256-60.
18. Jacobs KA, Burns P, Kressler J, Nash MS. Heavy reliance on carbohydrate across a wide range of exercise intensities during voluntary arm ergometry in persons with paraplegia. *J Spinal Cord Med* 2013;36(5):427-35.
19. de Zepetnek JOT, Pelletier CA, Hicks AL, MacDonald MJ. Following the Physical Activity Guidelines for Adults With Spinal Cord Injury for 16 Weeks Does Not Improve Vascular Health: A Randomized Controlled Trial. *Arch Phys Med Rehabil* 2015;96(9):1566-75.
20. Carlson KF, Wilt TJ, Taylor BC, Goldish GD, Niewoehner CB, Shamliyan TA et al. Effect of Exercise on Disorders of Carbohydrate and Lipid Metabolism in Adults With Traumatic Spinal Cord Injury: Systematic Review of the Evidence. *J Spinal Cord Med* 2009;32(4):361-78.
21. Rimmer JH, Riley B, Wang E, Rauworth A, Jurkowski J. Physical activity participation among persons with disabilities - Barriers and facilitators. *Am J Prev Med* 2004;26(5):419-25.
22. Kehn M, Kroll T. Staying physically active after spinal cord injury: a qualitative exploration of barriers and facilitators to exercise participation. *BMC Public Health* 2009;9.
23. Craig A, Tran Y, Wijesuriya N, Middleton J. Fatigue and tiredness in people with spinal cord injury. *J Psychosomat Res* 2012;73(3):205-10.
24. Gorgey AS. Exercise awareness and barriers after spinal cord injury. *World J Orthop* 2014;5(3):158-62.
25. Myers J, Lee M, Kiratli J. Cardiovascular Disease in Spinal Cord Injury. *Am J Phys Med Rehabil* 2007;86(2):142-52.
26. Nash MS. Exercise as a health-promoting activity following spinal cord injury. *J Neurol Phys Ther* 2005;29(2):87-106.
27. Hjeltne N, Galuska D, Bjornholm M, Aksnes AK, Lannem A, Zierath JR et al. Exercise-induced overexpression of key regulatory proteins involved in glucose uptake and metabolism in tetraplegic persons: molecular mechanism for improved glucose homeostasis. *Faseb J* 1998;12(15):1701-12.
28. Mohr T, Dela F, Handberg A, Biering-Sorensen F, Galbo H, Kjaer M. Insulin action and long-term electrically induced training in individuals with spinal cord injuries. *Med Sci Sports Exerc* 2001;33(8):1247-52.
29. Jeon JY, Weiss CB, Steadward RD, Ryan E, Burnham RS, Bell G et al. Improved glucose tolerance and insulin sensitivity after electrical stimulation-assisted cycling in people with spinal cord injury. *Spinal Cord* 2002;40(3):110-7.
30. Griffin L, Decker MJ, Hwang JY, Wang B, Kitchen K, Ding Z et al. Functional electrical stimulation cycling improves body composition, metabolic and neural factors in persons with spinal cord injury. *J Electromyogr Kinesiol* 2009;19(4):614-22.
31. Phillips SM, Stewart BG, Mahoney DJ, Hicks AL, McCartney N, Tang JE et al. Body-weight-support treadmill training improves blood glucose regulation in persons with incomplete spinal cord injury. *J Appl Physiol* 2004;97(2):716-24.
32. Moholdt T, Wisløff U, Nilsen TI, Slørdahl SA. Physical activity and mortality in men and women with coronary heart disease: a prospective population-based cohort study in Norway (the HUNT study). *Eur J Cardiovasc Prev Rehabil* 2008;15(6):639-45.

33. Wisløff U, Nilsen TI, Drøyvold WB, Mørkved S, Slørdahl SA, Vatten LJ. A single weekly bout of exercise may reduce cardiovascular mortality: how little pain for cardiac gain? 'The HUNT study, Norway'. *Eur J Cardiovasc Prev Rehabil* 2006;13(5):798-804.
34. Wen CP, Wai JP, Tsai MK, Yang YC, Cheng TY, Lee MC et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011;378(9798):1244-53.
35. Gebel K, Ding D, Chey T, Stamatakis E, Brown WJ, Bauman AE. Effect of Moderate to Vigorous Physical Activity on All-Cause Mortality in Middle-aged and Older Australians. *JAMA Intern Med* 2015;175(6):970-7.
36. Samitz G, Egger M, Zwahlen M. Domains of physical activity and all-cause mortality: systematic review and dose-response meta-analysis of cohort studies. *Int J Epidemiol* 2011;40(5):1382-400.
37. MacInnis MJ, Zacharewicz E, Martin BJ, Haikalis ME, Skelly LE, Tarnopolsky MA et al. Superior mitochondrial adaptations in human skeletal muscle after interval compared to continuous single-leg cycling matched for total work. *J Physiol* 2016.
38. Weston KS, Wisløff U, Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med* 2014;48(16):1227-34.
39. Ramos JS, Dalleck LC, Tjonna AE, Beetham KS, Coombes JS. The impact of high-intensity interval training versus moderate-intensity continuous training on vascular function: a systematic review and meta-analysis. *Sports Med* 2015;45(5):679-92.
40. Jolleyman C, Yates T, O'Donovan G, Gray LJ, King JA, Khunti K et al. The effects of high-intensity interval training on glucose regulation and insulin resistance: a meta-analysis. *Obes Rev* 2015;16(11):942-61.
41. Duran FS, Lugo L, Ramirez L, Eusse E. Effects of an exercise program on the rehabilitation of patients with spinal cord injury. *Arch Phys Med Rehabil* 2001;82(10):1349-54.
42. Hooker SP, Wells CL. Effects of low-intensity and moderate-intensity training in spinal cord-injured persons. *Med Sci Sports Exerc* 1989;21(1):18-22.
43. Midha M, Schmitt JK, Sclater M. Exercise effect with the wheelchair aerobic fitness trainer on conditioning and metabolic function in disabled persons: A pilot study. *Arch Phys Med Rehabil* 1999;80(3):258-61.
44. El-Sayed MS, Younesian A. Lipid profiles are influenced by arm cranking exercise and training in individuals with spinal cord injury. *Spinal Cord* 2005;43(5):299-305.
45. Swain DP, Leutholtz BC. Heart rate reserve is equivalent to %VO₂ reserve, not to %VO₂max. *Med Sci Sports Exerc* 1997;29(3):410-4.
46. Gibala MJ, Little JP, van Essen M, Wilkin GP, Burgomaster KA, Safdar A et al. Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *J Physiol* 2006;575(Pt 3):901-11.
47. Cocks M, Shaw CS, Shepherd SO, Fisher JP, Ranasinghe AM, Barker TA et al. Sprint interval and endurance training are equally effective in increasing muscle microvascular density and eNOS content in sedentary males. *J Physiol* 2013;591(Pt 3):641-56.
48. Shepherd SO, Cocks M, Tipton KD, Ranasinghe AM, Barker TA, Burniston JG et al. Sprint interval and traditional endurance training increase net intramuscular

- triglyceride breakdown and expression of perilipin 2 and 5. *J Physiol* 2013;591(Pt 3):657-75.
49. Gillen JB, Martin BJ, MacInnis MJ, Skelly LE, Tarnopolsky MA, Gibala MJ. Twelve Weeks of Sprint Interval Training Improves Indices of Cardiometabolic Health Similar to Traditional Endurance Training despite a Five-Fold Lower Exercise Volume and Time Commitment. *PLoS One* 2016;11(4):e0154075.
 50. Rakobowchuk M, Tanguay S, Burgomaster KA, Howarth KR, Gibala MJ, MacDonald MJ. Sprint interval and traditional endurance training induce similar improvements in peripheral arterial stiffness and flow-mediated dilation in healthy humans. *Am J Physiol Regul Integr Comp Physiol* 2008;295(1):R236-42.
 51. Gibala MJ, Gillen JB, Percival ME. Physiological and health-related adaptations to low-volume interval training: influences of nutrition and sex. *Sports Med* 2014;44 Suppl 2:S127-37.
 52. Janssen TWJ, Dallmeijer AJ, Veeger D, van der Woude LHV. Normative values and determinants of physical capacity in individuals with spinal cord injury. *J Rehabil Res Dev* 2002;39(1):29-39.
 53. Haisma JA, van der Woude LHV, Stam HJ, Bergen MP, Sluis TAR, Bussmann JBJ. Physical capacity in wheelchair-dependent persons with a spinal cord injury: a critical review of the literature. *Spinal Cord* 2006;44(11):642-52.
 54. Rosety-Rodriguez M, Camacho A, Rosety I, Fornieles G, Rosety MA, Diaz AJ et al. Low-Grade Systemic Inflammation and Leptin Levels Were Improved by Arm Cranking Exercise in Adults With Chronic Spinal Cord Injury. *Arch Phys Med Rehabil* 2014;95(2):297-302.
 55. Nightingale TE, Walhin, J. P., Thompson, D., Bilzon, J.L.J. Impact of Moderate-intensity Exercise on Metabolic Health and Aerobic Capacity in Persons with Chronic Paraplegia. *Med Sci Sports Exerc* 2016;48(5).
 56. de Groot PCE, Hjeltne N, Heijboer AC, Stal W, Birkeland K. Effect of training intensity on physical capacity, lipid profile and insulin sensitivity in early rehabilitation of spinal cord injured individuals. *Spinal Cord* 2003;41(12):673-9.
 57. Sæter TF. High versus moderate intensity arm-crank exercise for improving oxygen uptake and cardiovascular risk factors in spinal cord injured Trondheim Norwegian University of Science and Technology, NTNU - Trondheim 2014.
 58. Harnish CR, Daniels JA, Caruso D. Training response to high-intensity interval training in a 42-year-old man with chronic spinal cord injury. *J Spinal Cord Med* 2016;1-4.
 59. Helgerud J, Høydal K, Wang E, Karlsen T, Berg P, Bjerkaas M et al. Aerobic high-intensity intervals improve VO₂max more than moderate training. *Med Sci Sports Exerc* 2007;39(4):665-71.
 60. Nybo L, Sundstrup E, Jakobsen MD, Mohr M, Hornstrup T, Simonsen L et al. High-intensity training versus traditional exercise interventions for promoting health. *Med Sci Sports Exerc* 2010;42(10):1951-8.
 61. Schjerve IE, Tyldum GA, Tjønnå AE, Stølen T, Loennechen JP, Hansen HE et al. Both aerobic endurance and strength training programmes improve cardiovascular health in obese adults. *Clin Sci (Lond)* 2008;115(9):283-93.
 62. Karstoft K, Winding K, Knudsen SH, Nielsen JS, Thomsen C, Pedersen BK et al. The effects of free-living interval-walking training on glycemic control, body composition, and physical fitness in type 2 diabetic patients: a randomized, controlled trial. *Diabetes Care* 2013;36(2):228-36.
 63. Lee DC, Sui XM, Artero EG, Lee IM, Church TS, McAuley PA et al. Long-Term Effects of Changes in Cardiorespiratory Fitness and Body Mass Index on All-Cause

- and Cardiovascular Disease Mortality in Men The Aerobics Center Longitudinal Study. *Circulation* 2011;124(23):2483-U348.
64. Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA* 2009;301(19):2024-35.
 65. Blair SN, Kampert JB, Kohl HW, 3rd, Barlow CE, Macera CA, Paffenbarger RS, Jr. et al. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. *JAMA* 1996;276(3):205-10.
 66. Blair SN, Kohl HW, 3rd, Paffenbarger RS, Jr., Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 1989;262(17):2395-401.
 67. Blair SN, Kohl HW, 3rd, Barlow CE, Paffenbarger RS, Jr., Gibbons LW, Macera CA. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA* 1995;273(14):1093-8.
 68. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;346(11):793-801.
 69. Bassett DR, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc* 2000;32(1):70-84.
 70. Lundby C, Montero D. CrossTalk opposing view: Diffusion limitation of O₂ from microvessels into muscle does not contribute to the limitation of $\dot{V}O_2$ max. *J Physiol* 2015;593(17):3759-61.
 71. Wagner PD. CrossTalk proposal: Diffusion limitation of O₂ from microvessels into muscle does contribute to the limitation of $\dot{V}O_2$ max. *J Physiol* 2015;593(17):3757-8.
 72. van der Zwaard S, de Ruiter JC, Noordhof DA, Sterrenburg R, Bloemers FW, de Koning JJ et al. Maximal oxygen uptake is proportional to muscle fiber oxidative capacity - from chronic heart failure patients to professional cyclists. *J Appl Physiol* (1985) 2016:jap.00355.2016.
 73. Gifford JR, Garten RS, Nelson AD, Trinity JD, Layec G, Witman MA et al. Symmorphosis and skeletal muscle $\dot{V}O_2$ max : in vivo and in vitro measures reveal differing constraints in the exercise-trained and untrained human. *J Physiol* 2016;594(6):1741-51.
 74. Devillard X, Rimaud D, Roche F, Calmels P. Effects of training programs for spinal cord injury. *Annales de readaptation et de medecine physique* 2007;50(6):490-8, 80-9.
 75. Hoeks J, Schrauwen P. Muscle mitochondria and insulin resistance: a human perspective. *Trends Endocrinol Metab* 2012;23(9):444-50.
 76. Lee DE, Kehlenbrink S, Lee HN, Hawkins M, Yudkin JS. Getting the message across: mechanisms of physiological cross talk by adipose tissue. *Am J Physiol-Endocrinol Metab* 2009;296(6):E1210-E29.
 77. Campbell PJ, Mandarino LJ, Gerich JE. Quantification of the relative impairment in actions of insulin on hepatic glucose-production and peripheral glucose-uptake in non-insulin-dependent diabetes-mellitus. *Metab-Clin Exp* 1988;37(1):15-21.
 78. Bauman WA, Kahn NN, Grimm DR, Spungen AM. Risk factors for atherogenesis and cardiovascular autonomic function in persons with spinal cord injury. *Spinal Cord* 1999;37(9):601-16.

79. Nuhlicek DNR, Spurr GB, Barboriak JJ, Rooney CB, Elghatit AZ, Bongard RD. Body-composition of patients with spinal-cord injury. *Eur J Clin Nutr* 1988;42(9):765-73.
80. Kocina P. Body composition of spinal cord injured adults. *Sports Med* 1997;23(1):48-60.
81. Dionyssiotis Y, Petropoulou K, Rapidi CA, Papagelopoulos P, Papaioannou N, Galanos A et al. Body composition in paraplegic men. *J Clin Densitom* 2008;11(3):437-43.
82. Biering-Sorensen B, Kristensen IB, Kjaer M, Biering-Sorensen F. Muscle after spinal cord injury. *Muscle Nerve* 2009;40(4):499-519.
83. Capaldo B, Gastaldelli A, Antonello S, Auletta M, Pardo F, Ciociaro D et al. Splanchnic and leg substrate exchange after ingestion of a natural mixed meal in humans. *Diabetes* 1999;48(5):958-66.
84. DeFronzo RA. The Triumvirate: β -Cell, Muscle, Liver: A Collusion Responsible for NIDDM. *Diabetes* 1988;37(6):667-87.
85. Gorgey AS, Dudley GA. Skeletal muscle atrophy and increased intramuscular fat after incomplete spinal cord injury. *Spinal Cord* 2007;45(4):304-9.
86. Shah PK, Gregory CM, Stevens JE, Pathare NC, Jayaraman A, Behrman AL et al. Non-invasive assessment of lower extremity muscle composition after incomplete spinal cord injury. *Spinal Cord* 2008;46(8):565-70.
87. Bakkum AJT, Paulson TAW, Bishop NC, Goosey-Tolfrey VL, Stolwijk-Swuste JM, van Kuppevelt DJ et al. Effects of hybrid cycle and handcycle exercise on cardiovascular disease risk factors in people with spinal cord injury: a randomized controlled trial. *J Rehabil Med* 2015;47(6):523-30.
88. Kim DI, Lee H, Lee BS, Kim J, Jeon JY. Effects of a 6-Week Indoor Hand-Bike Exercise Program on Health and Fitness Levels in People With Spinal Cord Injury: A Randomized Controlled Trial Study. *Arch Phys Med Rehabil* 2015;96(11):2033-U325.
89. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing - Comparison with the euglycemic insulin clamp. *Diabetes Care* 1999;22(9):1462-70.
90. Radziuk J. Homeostatic Model Assessment and Insulin Sensitivity/Resistance. *Diabetes* 2014;63(6):1850-4.
91. Tjønnå AE, Lee SJ, Rognmo Ø, Stølen TO, Bye A, Haram PM et al. Aerobic interval training versus continuous moderate exercise as a treatment for the metabolic syndrome: a pilot study. *Circulation* 2008;118(4):346-54.
92. Mitranun W, Deerochanawong C, Tanaka H, Suksom D. Continuous vs interval training on glycemic control and macro- and microvascular reactivity in type 2 diabetic patients. *Scand J Med Sci Sports* 2014;24(2):e69-76.
93. Iellamo F, Manzi V, Caminiti G, Vitale C, Castagna C, Massaro M et al. Matched dose interval and continuous exercise training induce similar cardiorespiratory and metabolic adaptations in patients with heart failure. *Int J Cardiol* 2013;167(6):2561-5.
94. Iellamo F, Caminiti G, Sposato B, Vitale C, Massaro M, Rosano G et al. Effect of High-Intensity interval training versus moderate continuous training on 24-h blood pressure profile and insulin resistance in patients with chronic heart failure. *Intern Emerg Med* 2014;9(5):547-52.
95. Earnest CP, Lupo M, Thibodaux J, Hollier C, Butitta B, Lejeune E et al. Interval training in men at risk for insulin resistance. *Int J Sports Med* 2013;34(4):355-63.

96. Little JP, Jung ME, Wright AE, Wright W, Manders RJ. Effects of high-intensity interval exercise versus continuous moderate-intensity exercise on postprandial glycemic control assessed by continuous glucose monitoring in obese adults. *Appl Physiol Nutr Metab* 2014;39(7):835-41.
97. Karstoft K, Christensen CS, Pedersen BK, Solomon TP. The acute effects of interval- Vs continuous-walking exercise on glycemic control in subjects with type 2 diabetes: a crossover, controlled study. *J Clin Endocrinol Metab* 2014;99(9):3334-42.
98. van Loon LJ, Greenhaff PL, Constantin-Teodosiu D, Saris WH, Wagenmakers AJ. The effects of increasing exercise intensity on muscle fuel utilisation in humans. *J Physiol* 2001;536(Pt 1):295-304.
99. Bogardus C, Thuillez P, Ravussin E, Vasquez B, Narimiga M, Azhar S. Effect of muscle glycogen depletion on in vivo insulin action in man. *J Clin Invest* 1983;72(5):1605-10.
100. Newsom SA, Schenk S, Thomas KM, Harber MP, Knuth ND, Goldenberg N et al. Energy deficit after exercise augments lipid mobilization but does not contribute to the exercise-induced increase in insulin sensitivity. *J Appl Physiol* 2010;108(3):554-60.
101. Holtz KA, Stephens BR, Sharoff CG, Chipkin SR, Braun B. The effect of carbohydrate availability following exercise on whole-body insulin action. *Appl Physiol Nutr Metab* 2008;33(5):946-56.
102. Shokawa T, Imazu M, Yamamoto H, Toyofuku M, Tasaki N, Okimoto T et al. Pulse wave velocity predicts cardiovascular mortality: findings from the Hawaii-Los Angeles-Hiroshima study. *Circ J* 2005;69(3):259-64.
103. Inaba Y, Chen JA, Bergmann SR. Prediction of future cardiovascular outcomes by flow-mediated vasodilatation of brachial artery: a meta-analysis. *Int J Cardiovasc Imaging* 2010;26(6):631-40.
104. Celermajer DS, Sorensen KE, Bull C, Robinson J, Deanfield JE. Endothelium-dependent dilation in the systemic arteries of asymptomatic subjects relates to coronary risk factors and their interaction. *J Am Coll Cardiol* 1994;24(6):1468-74.
105. West CR, AlYahya A, Laher I, Krassioukov A. Peripheral vascular function in spinal cord injury: a systematic review. *Spinal Cord* 2013;51(1):10-9.
106. Totosy de Zepetnek JO, Ditor DS, Au JS, MacDonald MJ. Impact of shear rate pattern on upper and lower limb conduit artery endothelial function in both spinal cord-injured and able-bodied men. *Exp Physiol* 2015;100(10):1107-17.
107. Molmen-Hansen HE, Stolen T, Tjonna AE, Aamot IL, Ekeberg IS, Tyldum GA et al. Aerobic interval training reduces blood pressure and improves myocardial function in hypertensive patients. *Eur J Prev Cardiol* 2012;19(2):151-60.
108. Wisløff U, Støylen A, Loennechen JP, Bruvold M, Rognmo Ø, Haram PM et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation* 2007;115(24):3086-94.
109. Bristow S, Dalal, K., Santos, J.O., Martinez-Arizala, A., Banavac, K. Prevalence of hypertension, dyslipidemia, and diabetes mellitus after spinal cord injury. *Federal Practitioner* 2013:15- 8.
110. Wecht JM, Weir JP, Galea M, Martinez S, Bauman WA. Prevalence of Abnormal Systemic Hemodynamics in Veterans With and Without Spinal Cord Injury. *Arch Phys Med Rehabil* 2015;96(6):1071-9.
111. Ciolac EG, Bocchi EA, Bortolotto LA, Carvalho VO, Greve JM, Guimarães GV. Effects of high-intensity aerobic interval training vs. moderate exercise on

- hemodynamic, metabolic and neuro-humoral abnormalities of young normotensive women at high familial risk for hypertension. *Hypertens Res* 2010;33(8):836-43.
112. Gater DR. Obesity After Spinal Cord Injury. *Phys Med Rehabil Clin N Am* 2007;18(2):333-51.
 113. Gorgey A, Gater D. Prevalence of Obesity After Spinal Cord Injury. *Top Spinal Cord Inj Rehabil* 2007;12(4):1-7.
 114. Edwards LA, Bugaresti JM, Buchholz AC. Visceral adipose tissue and the ratio of visceral to subcutaneous adipose tissue are greater in adults with than in those without spinal cord injury, despite matching waist circumferences. *Am J Clin Nutr* 2008;87(3):600-7.
 115. Nakamura T, Tokunaga K, Shimomura I, Nishida M, Yoshida S, Kotani K et al. Contribution of visceral fat accumulation to the development of coronary-artery disease in nonobese men. *Atherosclerosis* 1994;107(2):239-46.
 116. Boyko EJ, Fujimoto WY, Leonetti DL, Newell-Morris L. Visceral adiposity and risk of type 2 diabetes - A prospective study among Japanese Americans. *Diabetes Care* 2000;23(4):465-71.
 117. Shah RV, Murthy VL, Abbasi SA, Blankstein R, Kwong RY, Goldfine AB et al. Visceral Adiposity and the Risk of Metabolic Syndrome Across Body Mass Index The MESA Study. *JACC-Cardiovasc Imag* 2014;7(12):1222-35.
 118. Gorgey AS, Mather KJ, Gater DR. Central adiposity associations to carbohydrate and lipid metabolism in individuals with complete motor spinal cord injury. *Metabolism* 2011;60(6):843-51.
 119. Thompson D, Karpe F, Lafontan M, Frayn K. Physical activity and exercise in the regulation of human adipose tissue physiology. *Physiol Rev* 2012;92(1):157-91.
 120. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. Appropriate Physical Activity Intervention Strategies for Weight Loss and Prevention of Weight Regain for Adults. *Med Sci Sports Exerc* 2009;41(2):459-71.
 121. Warburton DE, McKenzie DC, Haykowsky MJ, Taylor A, Shoemaker P, Ignaszewski AP et al. Effectiveness of high-intensity interval training for the rehabilitation of patients with coronary artery disease. *Am J Cardiol* 2005;95(9):1080-4.
 122. Heydari M, Freund J, Boutcher SH. The effect of high-intensity intermittent exercise on body composition of overweight young males. *J Obes* 2012;2012:480467.
 123. Shepherd SO, Wilson OJ, Taylor AS, Thøgersen-Ntoumani C, Adlan AM, Wagenmakers AJ et al. Low-Volume High-Intensity Interval Training in a Gym Setting Improves Cardio-Metabolic and Psychological Health. *PLoS One* 2015;10(9):e0139056.
 124. Gillen JB, Percival ME, Ludzki A, Tarnopolsky MA, Gibala MJ. Interval training in the fed or fasted state improves body composition and muscle oxidative capacity in overweight women. *Obesity* 2013;21(11):2249-55.
 125. Almenning I, Rieber-Mohn A, Lundgren KM, Shetelig Løvvik T, Garnæs KK, Moholdt T. Effects of High Intensity Interval Training and Strength Training on Metabolic, Cardiovascular and Hormonal Outcomes in Women with Polycystic Ovary Syndrome: A Pilot Study. *PLoS One* 2015;10(9):e0138793.
 126. Rognmo Ø, Moholdt T, Bakken H, Hole T, Mølsted P, Myhr NE et al. Cardiovascular risk of high- versus moderate-intensity aerobic exercise in coronary heart disease patients. *Circulation* 2012;126(12):1436-40.
 127. Skelly LE, Andrews PC, Gillen JB, Martin BJ, Percival ME, Gibala MJ. High-intensity interval exercise induces 24-h energy expenditure similar to traditional

- endurance exercise despite reduced time commitment. *Appl Physiol Nutr Metab* 2014;39(7):845-8.
128. Hazell TJ, Olver TD, Hamilton CD, Lemon PW. Two minutes of sprint-interval exercise elicits 24-hr oxygen consumption similar to that of 30 min of continuous endurance exercise. *Int J Sport Nutr Exerc Metab* 2012;22(4):276-83.
 129. Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism* 2010;59(10):1421-8.
 130. Gilbert O, Croffoot JR, Taylor AJ, Nash M, Schomer K, Groah S. Serum lipid concentrations among persons with spinal cord injury - A systematic review and meta-analysis of the literature. *Atherosclerosis* 2014;232(2):305-12.
 131. Fisher G, Brown AW, Bohan Brown MM, Alcorn A, Noles C, Winwood L et al. High Intensity Interval- vs Moderate Intensity- Training for Improving Cardiometabolic Health in Overweight or Obese Males: A Randomized Controlled Trial. *PLoS One* 2015;10(10):e0138853.
 132. Kessler HS, Sisson SB, Short KR. The potential for high-intensity interval training to reduce cardiometabolic disease risk. *Sports Med* 2012;42(6):489-509.
 133. O'Donovan G, Owen A, Bird SR, Kearney EM, Nevill AM, Jones DW et al. Changes in cardiorespiratory fitness and coronary heart disease risk factors following 24 wk of moderate- or high-intensity exercise of equal energy cost. *J Appl Physiol* 2005;98(5):1619-25.
 134. Austin MA, Hokanson JE, Edwards KL. Hypertriglyceridemia as a cardiovascular risk factor. *Am J Cardiol* 1998;81(4a):7b-12b.
 135. Sarwar N, Danesh J, Eiriksdottir G, Sigurdsson G, Wareham N, Bingham S et al. Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation* 2007;115(4):450-8.
 136. Nash MS, deGroot J, Martinez-Arizala A, Mendez AJ. Evidence for an exaggerated postprandial lipemia in chronic paraplegia. *J Spinal Cord Med* 2005;28(4):320-5.
 137. Emmons RR, Ciriigliaro, C.M., Moyer, J.M., Garber, C.E., Kirshblum, S.C., Spungen, A.M., Bauman, W.A. Exaggerated postprandial triglyceride response identified in individuals with spinal cord injury with cardiac risk factors. *Med Sci Sports Exerc* 2009;41:404 - 5.
 138. Zilversmit DB. Atherogenesis - postprandial phenomenon. *Circulation* 1979;60(3):473-85.
 139. Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM. Fasting compared with nonfasting triglycerides and risk of cardiovascular events in women. *JAMA* 2007;298(3):309-16.
 140. Cowan RE, Nash MS. Cardiovascular disease, SCI and exercise: unique risks and focused countermeasures. *Disability & Rehabilitation* 2010;32(26):2228-36.
 141. Burns SF, Miyashita M, Stensel DJ. High-Intensity Interval Exercise and Postprandial Triacylglycerol. *Sports Med* 2015;45(7):957-68.
 142. Wisloff U, Stoylen A, Loennechen JP, Bruvold M, Rognmo O, Haram PM et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation* 2007;115(24):3086-94.
 143. Haykowsky MJ, Timmons MP, Kruger C, McNeely M, Taylor DA, Clark AM. Meta-analysis of aerobic interval training on exercise capacity and systolic function in patients with heart failure and reduced ejection fractions. *Am J Cardiol* 2013;111(10):1466-9.

144. Hollekim-Strand SM, Bjørgaas MR, Albrektsen G, Tjønnå AE, Wisløff U, Ingul CB. High-Intensity Interval Exercise Effectively Improves Cardiac Function in Patients With Type 2 Diabetes Mellitus and Diastolic Dysfunction A Randomized Controlled Trial. *J Am Coll Cardiol* 2014;64(16):1758-60.
145. Madssen E, Moholdt T, Videm V, Wisløff U, Hegbom K, Wiseth R. Coronary Atheroma Regression and Plaque Characteristics Assessed by Grayscale and Radiofrequency Intravascular Ultrasound After Aerobic Exercise. *Am J Cardiol* 2014;114(10):1504-11.
146. Malmo V, Nes BM, Amundsen BH, Tjønnå AE, Stoylen A, Rossvoll O et al. Aerobic Interval Training Reduces the Burden of Atrial Fibrillation in the Short Term: A Randomized Trial. *Circulation* 2016;133(5):466-73.
147. Warme CA, Backus D, Rajan S, Bombardier CH, Schomer KG, Burns SP. Adverse events in cardiovascular-related training programs in people with spinal cord injury: a systematic review. *J Spinal Cord Med* 2014;37(6):672-92.
148. Ginis KA, Hicks AL, Latimer AE, Warburton DE, Bourne C, Ditor DS et al. The development of evidence-informed physical activity guidelines for adults with spinal cord injury. *Spinal Cord* 2011;49(11):1088-96.
149. Jacobs PL, Nash MS. Exercise recommendations for individuals with spinal cord injury. *Sports Med* 2004;34(11):727-51.
150. Furlan JC, Fehlings MG. Cardiovascular complications after acute spinal cord injury: pathophysiology, diagnosis, and management. *Neurosurg Focus* 2008;25(5):E13.
151. Steinberg LL, Lauro FAA, Sposito MMM, Tufik S, Mello MT, Naffah-Mazzacoratti MG et al. Catecholamine response to exercise in individuals with different levels of paraplegia. *Brazilian J Med Biol Res* 2000;33(8):913-8.
152. Goosey-Tolfrey V, Lenton J, Goddard J, Oldfield V, Tolfrey K, Eston R. Regulating intensity using perceived exertion in spinal cord-injured participants. *Med Sci Sports Exerc* 2010;42(3):608-13.
153. Cowan RE, Ginnity KL, Kressler J, Nash MS, Nash MS. Assessment of the Talk Test and Rating of Perceived Exertion for Exercise Intensity Prescription in Persons With Paraplegia. *Top Spinal Cord Inj Rehabil* 2012;18(3):212-9.
154. Sandler CX, Lloyd AR, Barry BK. Fatigue Exacerbation by Interval or Continuous Exercise in Chronic Fatigue Syndrome. *Med Sci Sports Exerc* 2016.
155. Petrofsky JS. Thermoregulatory stress during rest and exercise in heat in patients with a spinal-cord injury. *Eur J Appl Physiol Occup Physiol* 1992;64(6):503-7.
156. Price MJ, Campbell IG. Effects of Spinal Cord Lesion Level upon Thermoregulation during Exercise in the Heat. *Med Sci Sports Exerc* 2003;35(7):1100-7.
157. Griggs KE, Price MJ, Goosey-Tolfrey VL. Cooling athletes with a spinal cord injury. *Sports Med* 2015;45(1):9-21.
158. Dyson-Hudson TA, Kirshblum SC. Shoulder pain in chronic spinal cord injury, part I: Epidemiology, etiology, and pathomechanics. *J Spinal Cord Med* 2004;27(1):4-17.
159. Samuelsson KAM, Tropp H, Gerdle B. Shoulder pain and its consequences in paraplegic spinal cord-injured, wheelchair users. *Spinal Cord* 2004;42(1):41-6.
160. Cratsenberg KA, Deitrick CE, Harrington TK, Kopecky NR, Matthews BD, Ott LM et al. Effectiveness of Exercise Programs for Management of Shoulder Pain in Manual Wheelchair Users With Spinal Cord Injury. *J Neurol Phys Ther* 2015;39(4):197-203.

161. Jung ME, Bourne JE, Beauchamp MR, Robinson E, Little JP. High-Intensity Interval Training as an Efficacious Alternative to Moderate-Intensity Continuous Training for Adults with Prediabetes. *J Diabetes Res* 2015;2015:9.
162. Lunt H, Draper N, Marshall HC, Logan FJ, Hamlin MJ, Shearman JP et al. High intensity interval training in a real world setting: a randomized controlled feasibility study in overweight inactive adults, measuring change in maximal oxygen uptake. *PLoS One* 2014;9(1):e83256.
163. Saanijoki T, Nummenmaa L, Eskelinen JJ, Savolainen AM, Vahlberg T, Kalliokoski KK et al. Affective Responses to Repeated Sessions of High-Intensity Interval Training. *Med Sci Sports Exerc* 2015;47(12):2604-11.
164. Jung ME, Bourne JE, Little JP. Where does HIT fit? An examination of the affective response to high-intensity intervals in comparison to continuous moderate- and continuous vigorous-intensity exercise in the exercise intensity-affect continuum. *PLoS One* 2014;9(12):e114541.
165. Thum JS. Greater enjoyment in two modes of High Intensity Interval Training (HIIT) compared to continuous exercise training (CEX) in persons with spinal cord injury (SCI). California State University San Marcos; 2016.
166. Hasnan N, Engkasan JP, Husain R, Davis GM. High-Intensity Virtual-reality Arm plus FES-leg Interval Training in Individuals with Spinal Cord Injury. *Biomedizinische Technik Biomedical engineering* 2013.
167. Little JP, Gillen JB, Percival ME, Safdar A, Tarnopolsky MA, Punthakee Z et al. Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. *J Appl Physiol* 2011;111(6):1554-60.
168. Ross A, Leveritt M. Long-term metabolic and skeletal muscle adaptations to short-sprint training: implications for sprint training and tapering. *Sports Med* 2001;31(15):1063-82.
169. Gibala MJ, Little JP, Macdonald MJ, Hawley JA. Physiological adaptations to low-volume, high-intensity interval training in health and disease. *J Physiol* 2012;590(Pt 5):1077-84.

Table 1 High-intensity interval training (HIIT) protocols used in non-disabled and SCI research studies.

Table 2: Description of exercise training studies that have compared the impact of exercise intensity on cardiometabolic health markers in persons with SCI.

Figure 1: Schematic of sprint-interval training (SIT), high-intensity interval training (HIIT) and moderate-intensity continuous training (MICT) protocols (Adapted from Gibala *et al*,⁵¹ with permission).

Figure 2: Whole body Dual-energy X-ray absorptiometry (DEXA) scan of a female with neurological complete T7 injury sustained 6 years previously (a) and non-disabled female for comparative purposes (b). This figure visually highlights the drastic atrophy of lean mass and accumulation of intramuscular fat in the lower extremities of individuals with SCI.

Table 1

Authors	Exercise Intervals			Recovery Intervals		Total Session Time
	Number	Intensity	Duration	Intensity	Duration	
Little <i>et al.</i> , ¹⁶⁷ N-D Harnish <i>et al.</i> , ⁵⁸ SCI	10	90-110% Wmax ≥85% HRmax RPE ≥19	1-min	20-25% Wmax	1-2 min	~25 min
Tjønnna <i>et al.</i> , ⁹¹ N-D Sæter ⁵⁷ † SCI	4	~85% Wmax 85-95% HRmax RPE ≥17	2.5-4 min	20-25% Wmax	3-5 min	~30 min
MacInnis <i>et al.</i> , ³⁷ N-D Harnish <i>et al.</i> , ⁵⁸ SCI	3	~70% Wmax 80-85% HRmax RPE ≥ 16	4-5 min	20-25% Wmax	3-5 min	~30 min

Table 1 Legend: *HRmax* maximum heart rate, *N-D* non-disabled, *RPE* ratings of perceived exertion, *SCI* spinal cord injury, *Wmax peak power output (Watts)*, *achieved during an incremental test to fatigue*

Suggested frequency for training interventions is 3 sessions/week. Low-intensity warm-up and extended cool-down are not included in the table, but should be incorporated into any applied protocol to optimise circulation and prevent post-exercise hypotension (Evans *et al.*,¹⁴). We have suggested appropriate RPE values so that these protocols can be followed in patients with blunted cardiovascular responses to exercise (spinal cord injury lesions ≥T6). There is scope for variation in the above HIIT protocols, as the frequency, intensity and the duration of the high-intensity intervals, as well as the characteristics and duration of the recovery periods, may all be manipulated to change the nature of the exercise stimulus and thus potentially the physiological adaptations associated with training^{168, 169}

† Unpublished data


Table 2


Authors	Study Design	Participant Characteristics	Intervention	Outcome Measures		
				Change	No Change	
Hooker & Wells ^{42 *}	Pre-post parallel group WERG INT	6 (3F), 5 PARA, 1 TETRA, TSI; 4 mo - 19 yr Age; 26 - 36 yr	Frequency: 3 x wk Time: 20 min continuous Duration: 8 wks	Moderate-intensity (50 - 60% HRR)	↑ $\dot{V}O_2$ peak (10%), ↑ PPO (24%)	TC, TAG, LDL-C, HDL-C
		5 (2F), 3 PARA, 2 TETRA, TSI; 2 - 19 yr Age; 23 - 36 yr		High-intensity (70 - 80% HRR)	↑ $\dot{V}O_2$ peak (12%), ↑ PPO (13%) ↓ TAG (96 ± 28 to 78 ± 18 mg/dL; P ≤ 0.10), ↑ HDL-C (39 ± 11 to 47 ± 8 mg/dL; P ≤ 0.10), ↓ LDL-C (137 ± 26 to 116 ± 5 mg/dL; P ≤ 0.10)	TC
De groot <i>et al.</i> , ^{56 *}	Pre-post parallel group ACE INT	3 (2F), All PARA TSI; 61 - 225 days Age; 50 - 54 yr	Frequency: 3 x wk, Time: 60 min (3 & 2 minute work and rest intervals, respectively. Accumulated activity = 36 minutes) Duration: 8 wks	Moderate-intensity (40 - 50% HRR)	↑ $\dot{V}O_2$ peak (17%), ↑ PPO (24%)	TC, HDL-C, LDL-C, TAG. Non-significant improvement in IS (56%, measured via HOMA-CIGMA)
		3, 2 PARA, 1 TETRA TSI; 43 - 175 days Age; 20 - 38 yrs		High-intensity (70 - 80 % HRR)	↑ $\dot{V}O_2$ peak (50%), ↑ PPO (59%) ↓ TAG (-31%), ↓ IS (-33%, measured via HOMA-CIGMA)	TC, HDL-C, LDL-C
Sæter ^{57 †}	Pre-post parallel group ACE INT	5, All PARA TSI; 15 ± 11 yrs Age; 43 ± 14 yrs	Frequency: 3 x wk Time: ~ 49 min (373 kcal) Duration: 8 wks Moderate-intensity: 70% peak HR			$\dot{V}O_2$ peak, PPO, TC, HDL-C, LDL-C, TAG, glucose
		5, All PARA (1F) TSI; 15 ± 14 yrs Age; 46 ± 6 yrs	Frequency: 3 x wk Time: 28 min (including 12 min active recovery) Duration: 8 wks High-intensity: 85 - 95% peak HR (4 x 4 min intervals)		↑ $\dot{V}O_2$ peak (9%, trend for an interaction effect between groups, P = 0.051), ↑ PPO (28%)	TC, HDL-C, LDL-C, TAG, glucose

Table 1 Legend: *ACE* arm crank exercise, *HDL-C* high density lipoprotein cholesterol, *HR* heart rate, *HRR* heart rate reserve, *INT* intervention, *IS* insulin sensitivity, *LDL-C* low-density lipoprotein cholesterol, *PARA* paraplegic, *PPO* peak power output, *TAG* triglyceride, *TC* total cholesterol, *TETRA* tetraplegic, *TSI* time since injury, $\dot{V}O_2 peak$ peak oxygen uptake, *WC* waist circumference, *WERG* wheelchair ergometry.

* Note, authors refer to 70-80% HRR between studies as moderate⁴² and high-intensity⁵⁶, respectively. The terminology to describe exercise-intensity has been reclassified into moderate (40-60% HRR) and high-intensity (70-80% HRR).

† Unpublished data





Warm Up (10 minutes)

ACCEPTED MANUSCRIPT

Cardiovascular exercises (moving around independently in multi-directions)

- Marching with arm swings
- Walking backwards with knees straight
- Walking with leg curls
- Jogging
- Skipping
- Walking in slow motion (stepping with a one second pause before heel strike)
- Walking with longer strides
- Walking on heels
- Walking on toes


Upper body stretches

- Cervical rotation and side flexion (2 reps x 10 second hold bilaterally)
- Shoulder raises (2 reps x 5 second hold)
- Shoulder rolls (10 reps bilaterally)

Trunk stretches (with aqua noodle)

- Trunk rotation with arms abducted and externally rotated holding the aqua noodle (5 reps bilaterally)
- Arm raises reaching both arms overhead holding the noodle (5 reps bilaterally)
- Side bends pressing the aqua noodle into the water (5 reps x 5 second hold bilaterally)

Gait re-education (20 minutes)



Activity

- Continuous walking
- Stepping up and down off the steps

Progression

- Increase walking speed
- Stepping over steps
- Change of direction (turning)
- Walking with fins (as tolerated)

Strength exercises (10 minutes)

(2 minutes per exercise; 3 exercises selected per class with as many repetitions carried out as possible within the time)

Circuits

- Sit to stand (using pool chair)
- Step ups (progression: raising arms up and down holding the aqua noodle)
- Side step ups
- Trunk rotation (performed standing back to back with a partner, passing ball x 10 reps bilaterally)
- Squats with aqua noodle
- Lunges

Group

- Single leg stand (light finger hold at baseline progressed to 10 seconds with no hand support by session 12)
- Calf raises (10 reps at baseline progressed to 2 sets x 15 reps by session 12)
- Single leg calf raises (5 reps at baseline progressed to 15 reps by session 12)
- Push downs with aqua noodle (15 reps at baseline progressed to 30 reps by session 12)

Cool Down (5 minutes)

(Performed standing by pool wall at water depth level T8 (8th thoracic vertebrae), 30 second hold x 3 reps)

- Quadriceps, hamstring and calf stretches performed using aqua noodle