

Extent and ecological consequences of hunting in Central African rainforests in the twenty-first century

K. A. Abernethy, L. Coad, G. Taylor, M. E. Lee and F. Maisels

Phil. Trans. R. Soc. B 2013 **368**, 20120303, published 22 July 2013

Supplementary data

["Data Supplement"](#)

<http://rstb.royalsocietypublishing.org/content/suppl/2013/07/18/rstb.2012.0303.DC1.html>

["French Abstracts"](#)

<http://rstb.royalsocietypublishing.org/content/suppl/2013/07/23/rstb.2012.0303.DC2.html>

References

[This article cites 117 articles, 7 of which can be accessed free](#)

<http://rstb.royalsocietypublishing.org/content/368/1625/20120303.full.html#ref-list-1>

open access

This article is free to access

Erratum

An erratum has been published for this article, the contents of which has been appended at the end of this reprint. The erratum is available online at: [correction](#)
<http://rstb.royalsocietypublishing.org/content/368/1631/20130494.full.html>

Subject collections

Articles on similar topics can be found in the following collections

[ecology](#) (513 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Review



Cite this article: Abernethy KA, Coad L, Taylor G, Lee ME, Maisels F. 2013 Extent and ecological consequences of hunting in Central African rainforests in the twenty-first century. *Phil Trans R Soc B* 368: 20120303. <http://dx.doi.org/10.1098/rstb.2012.0303>

One contribution of 18 to a Theme Issue 'Change in African rainforests: past, present and future'.

Subject Areas:

ecology

Keywords:

Central Africa, hunting, future, wildlife, land-use change, ecological function

Author for correspondence:

K. A. Abernethy
e-mail: k.a.abernethy@stir.ac.uk

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rstb.2012.0303> or via <http://rstb.royalsocietypublishing.org>.

Extent and ecological consequences of hunting in Central African rainforests in the twenty-first century

K. A. Abernethy¹, L. Coad^{2,3}, G. Taylor³, M. E. Lee⁴ and F. Maisels^{1,5}

¹African Forest Ecology Group, School of Natural Sciences, University of Stirling, Stirling, UK

²School of Geography and Environmental Planning, University of Queensland, St Lucia Campus, Brisbane, Australia

³Environmental Change Institute, School of Geography, University of Oxford, Oxford, UK

⁴Wildlife Conservation Research Unit (WildCRU), University of Oxford, Oxford, UK

⁵Wildlife Conservation Society, Bronx, NY, USA

Humans have hunted wildlife in Central Africa for millennia. Today, however, many species are being rapidly extirpated and sanctuaries for wildlife are dwindling. Almost all Central Africa's forests are now accessible to hunters. Drastic declines of large mammals have been caused in the past 20 years by the commercial trade for meat or ivory. We review a growing body of empirical data which shows that trophic webs are significantly disrupted in the region, with knock-on effects for other ecological functions, including seed dispersal and forest regeneration. Plausible scenarios for land-use change indicate that increasing extraction pressure on Central African forests is likely to usher in new worker populations and to intensify the hunting impacts and trophic cascade disruption already in progress, unless serious efforts are made for hunting regulation. The profound ecological changes initiated by hunting will not mitigate and may even exacerbate the predicted effects of climate change for the region. We hypothesize that, in the near future, the trophic changes brought about by hunting will have a larger and more rapid impact on Central African rainforest structure and function than the direct impacts of climate change on the vegetation. Immediate hunting regulation is vital for the survival of the Central African rainforest ecosystem.

1. Introduction

Hunting is a ubiquitous part of daily life in rural Central Africa. Wild meat is part of the village subsistence economy, and commercial wildlife hunting—practised in Central Africa for at least two millennia—continues today [1,2]. Contemporary illegal wildlife trade, now one of the three most important types of crime on the planet [3], uses village hunters to secure tusks, meat and skins, but an increasing number of commercial hunters, using heavier-calibre weapons than those available to villagers, and particularly targeting ivory-bearing elephants, are also hunting in the region; meat and ivory then pass via highly organized trade chains to their destinations in the cities of the region and overseas [4–7]. The ecological consequences of wildlife hunting in the tropics are far-reaching, with knock-on effects disrupting ecological function over large areas and in key ecosystems [8–10].

Human shaping of the rainforest environment is a significant evolutionary force for the biome [1,11], and African rainforest species, unlike those endemic to other continents, have evolved in the constant presence of hominid hunters and their ancestors [12]. However, modern human hunting pressure, driven largely by increasing commercial trade, has resulted in the local extirpation of many larger African rainforest mammals [13–17] and is changing wildlife assemblages and species interactions [18,19]. Overhunting of wildlife is highly associated with loss of ecological integrity in tropical forests [20] and significant, deleterious trophic cascades in other ecosystems [9,10], and similar changes are almost certainly happening in the rainforests of

Central Africa, although empirical studies are few. Modern hunting practices in Central Africa are already modifying rainforest ecosystem function, and in synergy with changes in climate and land use may become increasingly influential in the future.

Within this special issue, the future of African rainforests is discussed in relation to likely climate change and land-use scenarios. In this paper, we

- collate and synthesize existing data on direct hunting impacts in the Central African region;
- review the factors driving human hunting in Central African forests and the empirical evidence for indirect ecological impacts of hunting, and discuss how the African rainforests of the twenty-first century are being shaped by current hunting activity; and
- consider how the future scenarios for land-use change (LUC) and climate change outlined in this special issue are likely to influence, and interact with, the drivers of wildlife hunting in Central Africa, to explore the potential long-term consequences for the region's rainforests.

2. The hunting footprint

Central Africa is one of the 'most remote' areas of tropical moist forest in the world, based on human settlement density, infrastructure and road location [21], yet empirical data from village hunting studies and ecological surveys in the region show that much of this remote forest is already accessed by hunters (table 1).

Humans, as central place foragers, gather food resources in a halo around the village [40,41], and village hunters in Central Africa will usually travel less than 10 km from the village during a day trip (table 1). However, most studies also document village hunters' use of hunting camps. These satellite 'central places', situated up to 50 km from the village, are used to catch larger bodied species favoured by the commercial trade [41,42], which are often highly depleted closer to the village where hunting pressure is more intense [31,43]. There have been no direct studies of forest use and offtake by purely commercial hunters in Central Africa: salaried hunters tend not to be based at a fixed point, and their activity is predominantly illegal and therefore concealed. However, ecological transect surveys often record hunting signs such as snares, gun cartridges and hunting paths, which suggests that hunters are penetrating up to 40 km into the forest from the nearest access point such as roads and rivers (table 1).

Even in 1998, more than 40% of Central African forest was within 10 km of a road, and more than 90% was within 50 km of a road [44]. Recent improvements in the road networks of many countries, expansion of logging and other industries and availability of mechanized transport in the region have increased the number of access points for hunters, who are likely to now be accessing the majority of the 'most remote' lands throughout the Congo Basin [44,45] (see the electronic supplementary material, figure S1).

3. Biological extent and impact of hunting

The biological impacts of hunting comprise both the direct impact on prey species (removal of individuals) and the

cascade effects of changing ecological function across the trophic web, as species declining under extreme hunting pressure change their ecological interactions with others [10].

(a) Prey species

An estimated 178 species are currently hunted and used in the wild meat industry in Central Africa [46]. The survival of over half (97) of these species is deemed threatened by this hunting [47]. Site-specific species lists compiled throughout the region typically show that the majority of local species are used in wild meat consumption and trade, with mammals dominating the harvest (see the electronic supplementary material, table S1). On average, over 60% of village hunting offtake comprises small ungulate and rodent species (see the electronic supplementary material, table S1), which are often caught using wire cable or tough plastic snares. However, escalating gun hunting facilitates the hunting of larger bodied animals, such as elephant or buffalo and primates, which can become the most common taxonomic group at market and village level in regions opened to gun hunting [18,48]. Ivory poaching is currently having exceptionally high impacts in Central Africa [49–51]. Forest elephants across the region declined by 62% between 2002 and 2011 and there is no sign of a fall in the rate of poaching [17]. Almost 90% of elephant carcasses found by guard patrols and on surveys within Central African protected areas in 2011 had been illegally killed [52,53]. Ape populations are also dropping very rapidly in the region, declining by 50% between 1984 and 2000 [54], targeted by hunters for commercial opportunity rather than for family meat [55].

(b) Changes in wildlife assemblages

Wildlife species are not equally affected by hunting, although some general ecological rules are clear: large, low-density, slow-reproducing and specialist species, such as elephants, will be more vulnerable to increases in predation pressure than smaller, fast-reproducing and high-density generalist species, such as rodents [56,57].

As forest elephants can represent between 33 and 89% of the animal biomass of intact Central African forests, and diurnal primates up to 30% [58,59], the dramatic declines recorded for these species will radically alter functional relationships in which they play a key role [19,60,61]. However, the detrimental and cascading effects of losing large fauna from an ecosystem are not always visible in tropical forests where forest cover and tree density are often used as proxy indicators of ecosystem health [20,62].

Although loss of the 'large forest architects' may cause the most obvious ecosystem changes, other shifts in species composition will also have important impacts on forest structure and function. Small species released from predation pressure and competition as their natural predators and competitors are hunted to low densities, yet themselves unattractive to human hunters, can find conditions of high hunting pressure favourable and densities may even locally increase, with knock-on consequences for the area's ecology [19,40]. In Central Africa, frugivorous medium-bodied duikers and primates make up a high proportion of hunting offtakes (see figure 1 and electronic supplementary material, table S1), and reductions in their abundance can be dramatic: Lahm

Table 1. Village and commercial hunting distances.

(a) evidence of village hunting distances, from village hunting surveys

source	site	method	maximum distance of traps and hunting camps from village
[22]	Ekou, Cameroon	GPS follow	hunting camps: 18 km
[23]	Oleme and Diba, ROC	GPS follow	traplines: 8.0 (Oleme) and 3.9 km (Diba)
[24]	Ituri, DRC	estimate	most hunting occurs within approximately 15 km of settlements
[25]	Mossapouna, CAR	GPS follow	day hunting: 10 km from village; hunting camps: 21 km from village
[26]	Sendje, Equatorial Guinea	GPS follow	hunting camps: 30 km, hunters trap a max of 3.2 km from the village or a hunting camp (1 day trip)
[27]	Zoulabot Ancien, Cameroon	GPS follow	hunting camps: 21.5 km from village, snares set in a radius of approximately 3 km from camp; every 2 years hunters stay in a hunting camp less than 50 km from the village, for 2 months or more
[28]	Mekas, Cameroon	estimate	village hunting: approximately 5–10 km; hunting camps: approximately 40 km
[29]	Nsiete, Gabon	hunter interviews	more than 10 km in the dry season. Less than 2 km in the wet season
[30]	Midyobo Anvom, Equatorial Guinea	GPS follow	hunting camps: 13.2 km, hunters travelled 2–3 km from hunting camps to place traps
[31] ^a	Dibouka and Kouagna, Gabon	GPS follow	village traplines: 6.5 km; hunting camps 12.7 km

(b) evidence of hunting presence in protected areas, from ecological transect surveys

source	site	furthest record of hunting sign, from nearest road or river (foot transect surveys; in kilometres)
[32]	Nouabale Ndoki, protected landscape ROC	<20
[32]	Ntokou Pikounda forest, ROC	<25
[33]	Mbam Djerem National Park, Cameroon	<27
[34]	Odzala-Koukoua National Park, ROC	<30
[35]	Sankuru landscape, DRC	<40
[36]	Lopé National Park, Gabon	<22
[37]	Mont de Cristal, Gabon	<10
[38]	Lac Tele National Park, ROC	<25
[39]	Waka National Park, Gabon	<10

^aMaximum trapping distances calculated for this paper; analysis of trapping distances and methods can be found in Coad *et al.* [31].

[63] recorded local extirpation of yellow-backed duiker around a village of northeastern Gabon less than a year from the onset of the unregulated hunting now typical of the entire region. The role of these species in the middle level of the trophic web, as browsers, seed dispersers and producers of food for the higher trophic levels, makes them a key part of the ecosystem: changes in the abundance of the entire guild are certain to have multiple consequences for the ecosystem [13,19,64]. Leopards, the apex predator in Central African forests, have already been lost from heavily hunted areas due to loss of the species which are their prey base, rather than direct persecution [15].

4. Current drivers of hunting

Globally, the need for both protein and income are well recognized as the ultimate drivers of hunting [31,65–67].

Wild meat is a traditional food for almost all ethnic groups in Central Africa, consumed by rich and poor, in

urban and rural communities [68,69]. While hunting is predominantly a poor man's activity [70], eating wild meat is not; hunting traditions are a source of pride in urban populations, who pronounce preferences for wild meat and will pay a premium for it, even when their protein requirement can be met with cheaper sources [67,68,71,72]. Vigorous trading of wild meat to satisfy urban demand is widespread in all major Central African cities [71,73–78], and the purchase of wild meat is common in even relatively small towns [41,68,79,80]. The certainty of demand, ease of entering the market and low risk of penalties have encouraged villagers in subsistence economies across the region to use local wildlife as a cash crop [66,81–83].

Commercial hunting has increased markedly in the last 50 years [18,63,81]. Recent studies at village subsistence level have found that more than 90% of resident village hunters sell at least part of their catch in all countries of the region [45,67,81,82,84,85] and that demand along the commodity chain is biased towards larger and rarer species [83,84,86–88]. In addition to village hunters, who are in




Figure 1. The impact of village hunting on trophic guilds of the Central African forests. Classification of species to these guilds can be found in the electronic supplementary material, table S1. Grey bars represent non-seed disperser, and black bars represent seed disperser.

part hunting to feed themselves, the purely commercial hunters, who often hunt species to order, are accessing more remote land in production forests or protected areas where village land rights do not apply and resources are *de facto* open access [62,69]. This type of hunting is particularly deleterious to the largest species, such as elephants or gorillas, which are typically less highly impacted by village subsistence activities [17,55].

(a) Sustainability of hunting offtakes

Considerable effort has been made in the last 20 years to develop indices and methods for evaluating hunting sustainability [89–93]. However, contemporary impacts depend enormously on the history of the local area and the dynamic interactions between hunters and the local wildlife community. Impacts of a given hunting pressure can only be predicted in the light of the area's recent past, which has already shaped the communities of both hunters and their available prey [31,94]. None of the sustainability indicators currently in use have been found to perform well [95], and empirical data on wildlife population trends remain the only valid measure of hunting impacts at the species level. To date, using either proxies or direct measures of population trends, only smaller species, such as blue duiker or large rodents, have shown any indication of being resilient to modern Central African hunting regimes sustained over several years [19,29,31,90,96]. If current hunting offtakes are unsustainable, both direct impacts and cascading ecological impacts will intensify over coming years unless hunting practices change.

5. Ecosystem function

Ecological systems are shaped by 'top-down' forces, such as predation, and 'bottom-up' forces, such as climate or land use [10]. The top-down force of overhunting seems already to be causing tropic cascade changes in Central Africa. A suite of empirical studies give evidence for relational changes between species and disruption of ecological

function in hunted tropical forests. Although much of the evidence available to date is from outside Central Africa, recent studies in the region confirm similar responses to overhunting to those documented in other tropical forests. Functional changes recorded relate to trophic roles, structural changes in forests, changes in species diversity and richness, seed dispersal, pollination and soil nutrient cycling. Table 2 collates existing empirical evidence for ecological change as a result of hunting, in all tropical forests and African rainforests. Central African forests still harbour large populations of extremely large mammalian 'forest architects' or 'ecological engineers', such as elephant, hippo, buffalo and gorilla. They are likely to experience significant changes in forest structure after their loss [117]. Importantly, significant implications of mesofaunal loss are also emerging from recent empirical studies [15,19,115].

Central African hunting systems are biased towards heavy offtakes of seed-dispersing frugivorous mammals; over 70% of animals in an average village hunting offtake have a seed dispersal role (see figure 1 and electronic supplementary material, table S1). Animals disperse the seeds of the majority of tree species in tropical forests [118] and seed dispersal by larger animal guilds particularly affected by hunting plays a key role in both the spatial pattern of tree recruitment and survival [19,61,102,107], and the relationship between tree and liana establishment [104,119,120]. In the absence of specialized dispersing fauna, recruitment and survival of seedlings is reduced for tree species that depend on them [19,121]. Animal-dispersed tropical trees are biased towards species with slower growth, longer life and higher wood density than abiotically dispersed species [105]. A positive (if weak) correlation between seed size and wood density also exists [122]. Animal-dispersed tree species, particularly those with large seeds dispersed by large mammals, therefore contribute a high proportion of the overall carbon-storage capacity of tropical forests. Carbon storage may therefore erode over time if tree regeneration is hampered by changes in faunal guilds, including extinction of large specialized disperser species [61,104,123,124] or increases in seed-predating species enjoying ecological release from their predators [99,125]. Recent data already show significant changes in the seedling layer of hunted West African forests [19,107] with regeneration in defaunated forests favouring faster growing, lower density plants and resulting in overall loss of diversity. This effect has also been recorded in Asian forests [104]. Across the tropics, forests suffering high levels of hunting over the past three decades have also recorded increases in the establishment of fast-growing pioneer species and lianas, and the loss of large hardwood trees [20].

6. Impacts of land-use change and climate change on Central African forest structure and function

Central African rainforests will be influenced by both climate change and LUC as well as by hunting over the coming decades. We summarize the climate change and LUC scenarios outlined in this special issue and the recent literature and go on to discuss their potential interaction with the ecological changes caused by hunting, already in motion.

Table 2. Empirical evidence of the impacts of hunting on ecological function in tropical forest ecosystems.

	disruption to ecological functioning as a result of hunting	examples from literature	empirical studies from African rainforests	empirical studies from other tropical forests
changes in wildlife assemblages	interspecific competition (between predator and prey)	apex and mesopredators inhabiting heavily hunted areas are forced to prey on smaller species through direct competition with hunters for larger bodied species; this reduces predator population sizes	[15]	[8,97]
	interspecific competition (between prey species)	heavy hunting pressure and selective hunting can change the competitive balance between sympatric species; many species have separate niches due to competition rather than habitat constraints	[96,98]	[40]
	ecological and predator release	as large herbivores and top predators are extirpated, an initial increase in secondary game species is witnessed; populations of small, non-target species increase but overall animal biomass decreases	[19,79]	[40]
changes in relative seed dispersal success	seed dispersal failure due to seed predators both increasing and switching to more abundant seeds	in heavily hunted areas, rodents and other small seed predators switch feeding behaviour to animal-dispersed species whose undispersed seeds are now found in clusters around the parent plant preference by small-bodied mammal species for small seeds; low predation on large seeds in heavily hunted area	[19]	[99,100]
	reduced seed dispersal, sapling recruitment and plant regeneration	in hunted areas where large, long distance dispersers are absent, rates of dispersal and regeneration of animal-dispersed trees are much lower than where these species are present	[101–103]	[104–107]
	increased kin competition between seedlings	seedlings, which remain under parent plant due to reduced dispersal in hunted areas, experience increased competition		[108]
	changes in relative pre-dispersion predation pressure for seeds	pre-dispersal seed predation by larger mammals is higher in protected areas than in hunted areas		[109,110]
changes in vegetation structure and composition	reduced plant species richness and diversity	tree species richness is lower in regenerating cohorts in hunted areas than in protected sites	[19]	[104,111–113]
	changes in tree spatial structure	increased clustering and densities of saplings of animal-dispersed plants due to loss of large herbivores	[102]	[104]
	increased proportion of lianas (and other wind-dispersed species) due to decrease in animal seed dispersers	wind-dispersed climbing species, which are overwhelmingly woody lianas, are found in higher proportions in the seedling bank at heavily hunted sites than at protected sites	[19]	[106]
	changes in vegetation composition	diversity of plants with large- and medium-sized seeds was significantly lower while diversity of plants with small-sized seeds increased significantly with hunting pressure	[114]	[100]

(Continued.)

Table 2. (Continued.)

	disruption to ecological functioning as a result of hunting	examples from literature	empirical studies from African rainforests	empirical studies from other tropical forests
soil quality	altered nutrient cycling and parasite suppression, disturbed soil fertilization and aeration	reduced abundance of Scarabaeinae beetles as a result of reduced availability of mammal dung alters ecosystem functioning		[113,115]
carbon balance	reduced carbon storage	reduced woody plant recruitment due to decreased large seed species will threaten carbon-storage capacity		[105,116]

(a) Land-use change: reduced habitat for forest species, more forests accessed

Central African economies rely on extractive industries, allocating a large part of their territories to formal sector oil, mining, agriculture and extensive timber use [126]. For example, in Gabon, 59% of land is currently allocated to oil and mining production, industrial agriculture and commercial logging, with logging comprising 54% [127]. In 2007, over 30% of all Central African forests were allocated to logging [45]. Despite this allocation, current levels of deforestation in Central Africa are low (0.08% per year from 2000 to 2010, [128]) and driven mainly by increases in population density and subsequent land conversion for small-scale agriculture [128].

Although land conversion and commercial logging are yet to result in large loss of forest canopy cover, the secondary impacts of extractive industries on forests have been substantial. In 2007, logging roads accounted for 38% of the road network in Central Africa, ranging from 13% in DRC to over 60% in Gabon and the Republic of Congo [45], significantly influencing forest disturbance and unregulated human access. Logging infrastructure and industrial roads usher in a domino effect of factors known to intensify hunting pressure, such as population growth from an immigrant workforce [129], increased income and demand for wild meat [130], increased forest access [131–133] and increased extraction to international markets for specialist products like ivory [6,17]. Although logging itself can affect animal densities by modifying habitat at landscape and local scales [134], evidence across the region indicates that secondary impacts of logging activity are currently of far greater ecological importance [135].

Future LUC in Central Africa is difficult to predict beyond the medium term in a region of political and economic instability; however, under current socio-economic conditions, land use may change rapidly, even in the short term [126]. The recent road network expansion in Central Africa, driven by global demand for raw natural resources, is expected to continue to increase access to the remaining remote tracts of forest [45,128], reducing the contiguous forest to smaller blocks both at local scales

[136,137] and across the Congo Basin [138]. International demand for agro-industrial products such as oil palm, rubber, sugar cane, coffee and cocoa, is already responsible for forest conversion in the region [128] with governments actively soliciting agro-industrial investment. For example, the government of Gabon recently created a 40/60 joint venture with a large multinational to develop agroindustry as well as other products and services like industrial parks, which ultimately support extractive activities and redistribute worker populations to forest areas [127]. Under scenarios of increasing agro-industrialization and forest access, coupled with intrinsic population growth, unless regulated, hunting off-takes in Central African forests are expected to increase in the short term, and the hunting footprint will be likely to spread into the last remnants of remote forest.

(b) Climate change impacts: resilient forests?

Climate models for Central Africa, presented in this special issue, suggest that direct impacts of climate change on the region's forest cover may be lower than previously thought by the Intergovernmental Panel on Climate Change. These models predict an increase in temperature, and increased dry signals for Central Africa [139], yet importantly also suggest that extreme precipitation or drought events are unlikely to increase [140,141]. Central African rainforests are also suggested to be more resilient to water deficits compared with moist tropical forests elsewhere [142].

Although extreme change linked to temperature and drying, such as forest die back, seems unlikely for the Central African region [142–144] and devastating forest fires are not predicted in the near future by any of the climate models for Central Africa, even the slight drying associated with higher temperatures, rainfall changes and increased human incursion predicted for this region [128,139,140] may make forests more vulnerable to fire in the future [145,146]. Forest resilience has been used to mean the resistance of the vegetation to change [144]; however, ecological function in even a seemingly 'resilient' forest may be significantly affected by the relatively small increases in temperature predicted. Tutin & Fernandez [147] record a guild of trees in Gabon relying on temperature triggers for flowering (and

thus fruiting) events. In Uganda, increases in annual temperatures over 30 years have been correlated with a decline in the fruiting and flowering of some tropical tree species, while increasing the fecundity of others [148]. How Central African ligneous species will respond to rapidly changing temperature is poorly understood, but significant impacts on forest species composition or tree productivity could potentially change food availability for animals, affecting animal ranging patterns and densities across the region [149] and initiating trophic cascades as prey distributions, seed dispersal functions and nutrient cycling are in turn changed. These changes would be likely to be additive to any trophic change initiated as a direct result of hunting.

7. Discussion

In this paper, we set out to assemble the current empirical knowledge on hunting in Central Africa, its extent, drivers and direct and indirect ecological consequences, and to consider how interactions between hunting and current scenarios for LUC and climate change, outlined in this special issue, might influence the future of the African rainforests.

The data reviewed show that village hunting has been studied in detail at several sites in the six main forested countries of the African rainforest region over the past two decades (table 1). Drivers of hunting are very similar across all countries [65] and direct impacts on species are broadly similar across the Central African forests (see the electronic supplementary material, table S1). Twenty years ago, Barnes *et al.* [150] first documented the deleterious effect of human access on elephant densities in Gabon. Ten years later, Wilkie *et al.* [151] postulated that hunting, rather than habitat loss, would be the greatest driver of wildlife declines across Central African forests, due to increasing road access, and Barnes [152] further predicted that the insidious effects of hunting on wildlife populations would not be realized until species were close to collapse. Barnes' and Wilkie's hypotheses have been upheld for large species [15,17,20,50,51,133]. However, the majority of studies on hunting have focused on hunters, using only proxies for impacts on prey species [88,153]. Data for wildlife population responses to hunting are only available for large species for which robust census methods are available [154,155]. Data on the true status of smaller species, which form the bulk of the wildmeat harvest (see the electronic supplementary material, table S1), are almost entirely absent. Thus, although we have now registered the catastrophic decline of megafauna in the Central African rainforests, the immense consequences of the resulting trophic cascades, on mesofauna densities, forest structure and overall ecosystem function, are only now becoming understood.

As the extent and drivers of the ecological changes underway across the world's ecosystems in the twenty-first century become apparent [9,20,121], research paradigms must widen to become more multidisciplinary. Research on individual impacts of climate change, LUC or hunting on Central African forests risks missing the interactions between these factors. Considering one without the others could lead to widely different conclusions on the future health of Central African rainforests, and conservation, management and research priorities. For example, if viewed solely through the climate change lens, the scenarios outlined in this issue

predict minimal change for African rainforests and reasonable resilience to ongoing change [156]. However, coupled with more certain likelihood of LUC, the picture becomes increasingly dynamic and threatening to intact forests and wildlife. LUC predictions, though limited to the medium term, broadly indicate loss of contiguous forest blocks [138], increased accessibility of remote areas, overall forest degradation and increased local populations [128] and permanent agricultural lands [126,128]. The drivers of hunting summarized in this paper are likely to intensify under these LUC scenarios, particularly in regard to the large-bodied, commercially valuable species, which are also more vulnerable to forest cover fragmentation and loss [62,157]. Without greatly improved regulation, intensified drivers will result in intensified hunting offtakes and an increase in the rate of ecological change that these offtakes are effecting. Central African forests now look highly threatened.

Assuming current levels of hunting are maintained, evidence from the studies we reviewed might indicate one possible future scenario for African rainforests: Overhunting severely reduces large 'forest architects' and mesofauna, releasing small-bodied seed predators. This promotes fast-growing abiotically dispersed tree species over large-seeded, animal-dispersed, slow-growing, shade-bearing species, reducing carbon storage, fruit availability and associated biodiversity. In turn, the forests' ability to support large frugivorous, seed-dispersing mammals declines, spurring a negative feedback loop whereby both large mammals and large-seeded, long-lived, hard-wooded trees further decline, further reducing carbon storage and thus global resilience to climate change. Although this scenario represents dramatic, permanent change for African rainforests, it may be one plausible, logical outcome that puts the risks of not regulating hunting into perspective.

As Central Africa moves into the twenty-first century, the challenges that LUC and climate change bring are sure to be complex. However, they will be unlikely to reduce the demand for hunting in the region, nor to mitigate any of its ecological impacts. As outlined above, anticipated LUC in the region will probably exacerbate the impacts of hunting and the ecological changes resulting from hunting will probably in turn exacerbate the effects of climate change. LUC and hunting effects are likely to combine to reduce overall carbon-storage potential and biodiversity, diminish overall forest area and, through the structural and compositional changes initiated by hunting, lessen the resilience of remaining forests to drought, and possibly fire, over the long term.

Good hunting management practices and planning are clearly vital to maintaining ecological function in the African tropical forests and must be incorporated into research priorities and overall land-use and climate change strategies [158], as well as impact assessments and private sector management practices on the ground. Conservation practitioners in Central Africa have shown that multiple-use landscapes under efficient management can maintain the large wide-ranging species critical for large tree seed dispersal [159], sustain game populations for hunting needs [160] and support threatened species [161], if a few design rules are applied: for example, including sufficient protected areas and maintaining large timber concessions with a mix of logging histories, including unlogged patches, and ensuring sufficient resources are allocated to controlling hunting and enforcing management plans [160,162]. Similar kinds of

landscape-level work are also needed to rethink and reinvent how agriculture might be designed and implemented in Central Africa: for example, to intensify local food production without converting new lands, siting large-scale concessions in already degraded areas, and designing and managing them to mitigate landscape-scale impacts.

The studies on trophic cascades reviewed here (table 2) already indicate that limiting the loss of megafauna and apex predators should become a first priority for conservation strategies that seek to sustain intact ecosystem function in tropical forests [9,121]. However, global demand for ivory, human–wildlife conflicts with megafauna and overall forest loss, coupled with wildlife population declines already suffered, may make retaining viable populations of these species impossible outside large protected areas [51]. Research to determine the responses of mesopredators and ecological niche adjustments, in both wildlife assemblages impacted by overhunting and the plant communities they live in, will be critical in determining the true resilience of

the ecosystem and thus in designing hunting management plans and multiple land-use scenarios that could mitigate spiralling and irreversible ecological change.

If future Central African human and wildlife communities are to rely on the range of ecosystem services currently provided by their rainforests—and if the value of these global goods is to be maintained—immediate management of hunting and integration of good hunting practices into large-scale land-use planning must be considered an urgent priority for rainforest preservation and thus an integral and important part of planning for climate change management and mitigation.

Acknowledgements. We would like to acknowledge three anonymous referees whose comments improved the manuscript.

Funding statement. L.C. and G.T. would like to acknowledge the support of the Oxford University John Fell Fund and the Oxford Martin School.

References

- Barton H, Denham T, Neumann K, Arroyo Kalin M. 2012 Long-term perspectives on human occupation of tropical rainforests: an introductory overview. *Quat. Int.* **249**, 1–3. (doi:10.1016/j.quaint.2011.07.044)
- Nasi R, Taber A, Van Vliet N. 2011 Empty forests, empty stomachs? Bushmeat and livelihoods in the Congo and Amazon Basins. *Int. For. Rev.* **13**, 355–368. (doi:10.1505/146554811798293872)
- INTERPOL. 2012 *Environmental crime*. See <http://www.interpol.int/Public/EnvironmentalCrime/Default.asp>.
- Milliken T, Burn RW, Underwood FM, Sangalakula L. 2012 The elephant trade information system (ETIS) and the illicit trade in ivory: a report to the 16th meeting of the Conference of the Parties to CITES (Cop 16). See <http://www.cites.org/eng/cop/16/doc/E-CoP16-53-02-02.pdf>.TRAFFIC.
- Roe D, Milliken T, Milledge S, Mremi J, Mosha S, Grien-Gran M. 2002 *Making a killing or making a living? Wildlife trade, trade controls and rural livelihoods*. Nairobi, Kenya: IIED, TRAFFIC International.
- Stiles D. 2011 Elephant meat and ivory trade in Central Africa. *Pachyderm* **50**, 26–36.
- UNEP, CITES, IUCN, TRAFFIC. 2013 Elephants in the dust—the African elephant crisis: United Nations Environment Programme, GRID-Arendal. See http://www.grida.no/_cms/OpenFile.aspx?s=1&id=1570 (accessed 15 May 2013.)
- Brashares J, Pruh L, Stoner CJ, Epps CW (eds) 2010 *Ecological and conservation implications of mesopredator release*, 1st edn. Washington, DC: Island Press.
- Estes JA *et al.* 2011 Trophic downgrading of planet earth. *Science* **333**, 301–306. (doi:10.1126/science.1205106)
- Terborgh J, Holt RD, Estes JA. 2010 Trophic cascades: what they are, how they work and why they matter. In *Trophic cascades* (eds J Terborgh, JA Estes), 1st edn, pp. 1–19. Washington, DC: Island Press.
- Bayon G, Dennielou B, Etoubleau J, Ponzevera E, Toucanne S, Bermell S. 2012 Intensifying weathering and land use in iron age Central Africa. *Science* **335**, 1219–1222. (doi:10.1126/science.1215400)
- Brunet M *et al.* 2002 A new hominid from the Upper Miocene of Chad, Central Africa. *Nature* **418**, 145–151. (doi:10.1038/nature00879)
- Jimoh SO, Ikyagba ET, Alarape AA, Adeyemi AA, Waltert M. 2012 Local depletion of two larger duikers in the Oban hills region, Nigeria. *Afr. J. Ecol.* **50**, 1–7. (doi:10.1111/j.1365-2028.2011.01285.x)
- Hearn GW, Morra WA. 2001 *The approaching extinction of monkeys and duikers in Bioko Island, Equatorial Guinea, Africa*. Glenside, PA: Bioko Biodiversity Protection Program, Arcadia University.
- Henschel P, Hunter LTB, Coad L, Abernethy KA, Muhlenberg M. 2011 Leopard prey choice in the Congo Basin rainforest suggests exploitative competition with human bushmeat hunters. *J. Zool.* **285**, 11–20. (doi:10.1111/j.1469-7998.2011.00826.x)
- Maisels F, Kesting E, Menei M, Toh C. 2001 The extirpation of large mammals and implications for montane forest conservation: the case of the Kilum-Ijim Forest, North-west Province, Cameroon. *Oryx* **35**, 322–331. (doi:10.1046/j.1365-3008.2001.00204.x)
- Maisels F *et al.* 2013 Devastating decline of forest elephants in Central Africa. *PLoS ONE* **8**, e59469. (doi:10.1371/journal.pone.0059469)
- Fa JE, Brown D. 2009 Impacts of hunting on mammals in African tropical moist forests: a review and synthesis. *Mamm. Rev.* **39**, 231–264. (doi:10.1111/j.1365-2907.2009.00149.x)
- Effiom OA, Nunez-Iturri G, Smith HG, Olsson O. 2013 Bushmeat hunting changes regeneration of African rainforests. *Proc. R. Soc. B* **280**, 20130246. (doi:10.1098/rspb.2013.0246)
- Laurance WF *et al.* 2012 Averting biodiversity collapse in tropical forest protected areas. *Nature* **489**, 290–294. (doi:10.1038/nature11318)
- WCS, CIESIN, Columbia University. 2005 *Last of the wild project, Version 2, 2005 (LWP-2): global human footprint dataset (IGHP)*. Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC). See <http://sedac.ciesin.columbia.edu/data/set/wildareas-v2-human-footprint-ighp> (accessed 24 September 2012).
- Dethier. 1995 *Etude chasse*. Brussels, Belgium: ECOFAC, AGRECO-CTFT. See <http://www.ecofac.org/Biblio/Download/EtudesChasse/Cameroun/DethierChasseCameroun.pdf>.
- Vanwijnsberghe S. 1996 *Etude sur la chasse villageoise aux environs du Parc d'Odzala*. Brussels, Belgium: ECOFAC.
- Wilkie DS, Curran B, Tshombe R, Morelli GA. 1998 Modeling the sustainability of subsistence farming and hunting in the Ituri forest of Zaire. *Conserv. Biol.* **12**, 137–147. (doi:10.1046/j.1523-1739.1998.96156.x)
- Noss AJ. 1998 The impacts of BaAka net hunting on rainforest wildlife. *Biol. Conserv.* **86**, 161–167. (doi:10.1016/S0006-3207(98)00013-5)
- Kumpel NF. 2006 Incentives for sustainable hunting of bushmeat in Rio Muni, Equatorial Guinea. PhD thesis, Imperial College, London, UK.
- Yasuoka H. 2006 The sustainability of duiker (*Cephalophus* spp.) hunting for the Baka hunter-gatherers in southeastern Cameroon. *Afr. Study Monogr.* **33**, 95–120.
- Solly H. 2004 *Bushmeat hunters and secondary traders: making the distinction for livelihood improvement*. London, UK: ODI.
- Van Vliet N, Nasi R. 2008 Hunting for livelihood in northeast Gabon: patterns, evolution, and

- sustainability. *Ecol. Soc.* **13**, 33. See <http://www.ecologyandsociety.org/vol13/iss2/art33/>.
30. Rist J. 2008 Bushmeat catch per unit effort in space and time: a monitoring tool for bushmeat hunting. PhD thesis, Imperial College, London, UK.
 31. Coad L *et al.* 2013 Social and ecological change over a decade in a village hunting system, central Gabon. *Conserv. Biol.* **27**, 270–280. (doi:10.1111/cobi.12012)
 32. Maisels F *et al.* 2012 *Great ape and human impact monitoring training, surveys, and protection in the Ndoki-Likouala Landscape, Republic of Congo. GACF Agreement: 96200–9-G247. Final report.* New York, NY: Wildlife Conservation Society.
 33. Malonga R. 2008 *Final report on elephants and great apes survey in the Ngombe FMU and Ntokou-Pikounda forest, Republic of Congo. Contribution to a status survey of Great Apes.* New York, NY: Wildlife Conservation Society.
 34. Maisels F, Ambahe E, Ambassa R, Nvemgah Yara C, Fosso B. 2009 *Great ape and human impact monitoring in the Mbam et Djerem National Park, Cameroon. Final report to USFWS-GACF Agreement 98210–7-G290.* New York, NY: Wildlife Conservation Society.
 35. Blake S. 2006 *Final report on elephant and ape surveys in the Odzala-Koukoua National Park, 2005.* New York, NY: Wildlife Conservation Society.
 36. Liengola I, Maisels F, KNkumu P, Bonvenge A. 2010 *Conserving bonobos in the Lokofa Block of the Salonga National Park, Democratic Republic of Congo. Report to the Beneficia Foundation.* New York, NY: Wildlife Conservation Society.
 37. Maisels F, Bechem M, Mihindou Y. 2006 *Lope National Park, Gabon. Large mammals and human impact. Monitoring program 2004–2005.* New York, NY: Wildlife Conservation Society.
 38. Aba'a R. 2006 *Abondance relative des grands mammifères et des activités humaines au parc National des Monts de Cristal et sa peripherie.* Libreville, Gabon: Wildlife Conservation Society.
 39. Abitsi G. 2006 *Inventaires de reconnaissance des grands mammifères et de l'impact des activités anthropiques. Parc National de Waka, Gabon. December 2005–Juillet 2006.* Libreville, Gabon: Wildlife Conservation Society.
 40. Peres CA, Palacios E. 2007 Basin-wide effects of game harvest on vertebrate population densities in Amazonian forests: implications for animal-mediated seed dispersal. *Biotropica* **39**, 304–315. (doi:10.1111/j.1744-7429.2007.00272.x)
 41. Coad L. 2007 Bushmeat hunting in Gabon: socioeconomics and hunter behaviour. PhD thesis, University of Cambridge, Cambridge, UK.
 42. Colom A. 2006 *Aspects Socio-economiques de la gestion des ressources naturelles dans le paysage Salonga-Lukenie-Sankuru, DRC.* Kinshasa, Democratic Republic of the Congo: WWF CARPO.
 43. Kumpel NF, Rowcliffe JM, Cowlshaw G, Milner-Gulland EJ. 2009 Trapper profiles and strategies: insights into sustainability from hunter behaviour. *Anim. Conserv.* **12**, 531–539. (doi:10.1111/j.1469-1795.2009.00279.x)
 44. WRI. 2009 *Earthtrends.* Washington, DC: World Resources Institute.
 45. Laporte NT, Stabach JA, Grosch R, Lin TS, Goetz SJ. 2007 Expansion of industrial logging in Central Africa. *Science* **316**, 1451. (doi:10.1126/science.1141057)
 46. Taylor G. 2012 A systematic review of the bushmeat trade in West and Central Africa. MSc thesis, University of Oxford, Oxford, UK.
 47. IUCN. 2012 *The red list of threatened species.* See <http://www.iucnredlist.org/> (accessed 27 September 2012).
 48. Fa JE, Johnson PJ, Dupain J, Lapuente J, Koster P, Macdonald DW. 2004 Sampling effort and dynamics of bushmeat markets. *Anim. Conserv.* **7**, 409–416. (doi:10.1017/S136794300400160X)
 49. Beyers RL, Hart JA, Sinclair ARE, Grossmann F, Klinkenberg B, Dino S, Somers M. 2011 Resource wars and conflict ivory: the impact of civil conflict on elephants in the Democratic Republic of Congo: the case of the Okapi Reserve. *PLoS ONE* **6**, e27129. (doi:10.1371/journal.pone.0027129)
 50. Bouche P, Douglas-Hamilton I, Wittemyer G, Nianogo AJ, Doucet JL, Lejeune P, Vermeulen C. 2011 Will elephants soon disappear from West African Savannas? *PLoS ONE* **6**, e20619. (doi:10.1371/journal.pone.0020619)
 51. Bouche P, Mange RNM, Tankalet F, Zowoya F, Lejeune P, Vermeulen C. 2012 Game over! Wildlife collapse in northern Central African Republic. *Environ. Monit. Assess.* **184**, 7001–7011. (doi:10.1007/s10661-011-2475-y)
 52. CITES. 2012 *Monitoring the illegal killing of elephants.* CoP16 Doc. 53.1. See <http://www.cites.org/eng/cop/16/doc/E-CoP16-53-01.pdf>.
 53. CITES. 2013 *Monitoring the illegal killing of elephants—Addendum.* CoP16 Doc. 53.1: Addendum. See <http://www.cites.org/eng/cop/16/doc/E-CoP16-53-01-Addendum.pdf>.
 54. Walsh PD *et al.* 2003 Catastrophic ape decline in western equatorial Africa. *Nature* **422**, 611–614. (doi:10.1038/nature01566)
 55. Kuëhl HS, Nzeingui C, Yeno SLD, Huijbregts B, Boesch C, Walsh PD. 2009 Discriminating between village and commercial hunting of apes. *Biol. Conserv.* **142**, 1500–1506. (doi:10.1016/j.biocon.2009.02.032)
 56. Berryman AA. 1992 The origins and evolution of predator prey theory. *Ecology* **73**, 1530–1536. (doi:10.2307/1940005)
 57. Bennett EL, Robinson JG. 1999 *Hunting for sustainability in tropical forests.* New York, NY: Columbia University Press.
 58. White LJT, Tutin CEG, Fernandez M. 1993 Group composition and diet of forest elephants, *Loxodonta africana cyclotis* Matschie 1900, in the Lopé Reserve, Gabon. *Afr. J. Ecol.* **31**, 181–199. (doi:10.1111/j.1365-2028.1993.tb00532.x)
 59. Prins HHT, Reitsma JM. 1989 Mammalian biomass in an African equatorial rain-forest. *J. Anim. Ecol.* **58**, 851–861. (doi:10.2307/5128)
 60. Tutin CEG, Williamson EA, Rogers ME, Fernandez M. 1991 A case-study of a plant–animal relationship: *Cola lizae* and lowland gorillas in the Lopé Reserve, Gabon. *J. Trop. Ecol.* **7**, 181–199. (doi:10.1017/S0266467400005320)
 61. Campos-Arceiz A, Blake S. 2011 Megagardeners of the forest: the role of elephants in seed dispersal. *Acta Oecol.* **37**, 542–553. (doi:10.1016/j.actao.2011.01.014)
 62. Wilkie DS, Bennett EL, Peres CA, Cunningham AA. 2011 The empty forest revisited. *Ann. NY Acad. Sci.* **1223**, 120–128. (doi:10.1111/j.1749-6632.2010.05908.x)
 63. Lahm S. 2001 Hunting and wildlife in Northeastern Gabon. Why conservation should extend beyond protected areas. In *African rain forest ecology and conservation: an interdisciplinary perspective* (eds W Weber, LJT White, A Vedder), pp. 344–354. New York, NY: Yale University Press.
 64. Kumpel NF, Milner-Gulland EJ, Rowcliffe JM, Cowlshaw G. 2008 Impact of gun-hunting on diurnal primates in continental Equatorial Guinea. *Int. J. Primatol.* **29**, 1065–1082. (doi:10.1007/s10764-008-9254-9)
 65. Brashares JS, Golden CD, Weinbaum KZ, Barrett CB, Okello GV. 2011 Economic and geographic drivers of wildlife consumption in rural Africa. *Proc. Natl Acad. Sci. USA* **108**, 13 931–13 936. (doi:10.1073/pnas.1011526108)
 66. Kumpel NF, Milner-Gulland EJ, Cowlshaw G, Rowcliffe JM. 2010 Incentives for hunting: the role of bushmeat in the household economy in rural Equatorial Guinea. *Hum. Ecol.* **38**, 251–264. (doi:10.1007/s10745-010-9316-4)
 67. Foerster S, Wilkie DS, Morelli GA, Demmer J, Starkey M, Telfer P, Steil M, Lewbel A. 2012 Correlates of bushmeat hunting among remote rural households in Gabon, Central Africa. *Conserv. Biol.* **26**, 335–344. (doi:10.1111/j.1523-1739.2011.01802.x)
 68. Wilkie DS, Starkey M, Abernethy K, Effa EN, Telfer P, Godoy R. 2005 Role of prices and wealth in consumer demand for bushmeat in Gabon, Central Africa. *Conserv. Biol.* **19**, 268–274. (doi:10.1111/j.1523-1739.2005.00372.x)
 69. Nasi R, Brown D, Wilkie D, Bennett EL, Tutin C, van Tol G, Christophersen T. 2008 *Conservation and use of wildlife-based resources: the bushmeat crisis.* Bogor, Indonesia: Secretariat of the Convention on Biological Diversity, Montreal, and Center for International Forestry Research (CIFOR).
 70. Wilkie DS, Carpenter JF. 1999 Bushmeat hunting in the Congo Basin: an assessment of impacts and options for mitigation. *Biodivers. Conserv.* **8**, 927–955. (doi:10.1023/A:1008877309871)
 71. Schenck M, Effa EN, Starkey M, Wilkie D, Abernethy K, Telfer P, Godoy R, Treves A. 2006 Why people eat bushmeat: results from two-choice, taste tests in Gabon, Central Africa. *Hum. Ecol.* **34**, 433–445. (doi:10.1007/s10745-006-9025-1)
 72. East T, Kumpel NF, Milner-Gulland EJ, Rowcliffe JM. 2005 Determinants of urban bushmeat consumption in Rio Muni, Equatorial Guinea. *Biol. Conserv.* **126**, 206–215. (doi:10.1016/j.biocon.2005.05.012)
 73. Hennessey AB, Rogers JA. 2008 A study of the bushmeat trade in Ouessou, Republic of Congo.

- Conserv. Soc.* **6**, 179–184. (doi:10.4103/0972-4923.49211)
74. Edderal D, Dame M. 2006 A census of the commercial bushmeat in Yaounde, Cameroon. *Oryx* **40**, 472–475. (doi:10.1017/S0030605306001256)
 75. Malonga R. 1996 *Dynamique socio-economique du circuit commercial de viande de chasse a Brazzaville*. New York, NY: Wildlife Conservation Society.
 76. Fa JE, Garcia-Yuste JE. 2001 Commercial bushmeat hunting in the Monte Mitra forests, Equatorial Guinea: extent and impact. *Anim. Biodivers. Conserv.* **24**, 31–52.
 77. Mbayma G. 2009 *Bushmeat consumption in Kinshasa, DRC. Analysis at the household level*. New York, NY: Wildlife Conservation Society.
 78. Mbete RA *et al.* 2011 Profile of bushmeat sellers and evaluation of biomass commercialized in the municipal markets of Brazzaville, Congo. *Trop. Conserv. Sci.* **4**, 203–217.
 79. Fa JE, Seymour S, Dupain J, Amin R, Albrechtsen L, Macdonald D. 2006 Getting to grips with the magnitude of exploitation: bushmeat in the Cross–Sanaga rivers region, Nigeria and Cameroon. *Biol. Conserv.* **129**, 497–510. (doi:10.1016/j.biocon.2005.11.031)
 80. Tieguhong JC, Zwolinski J. 2009 Supplies of bushmeat for livelihoods in logging towns in the Congo Basin. *J. Horticulture For.* **1**, 2–9.
 81. Starkey M. 2004 *Commerce and subsistence: the hunting, sale and consumption of bushmeat in Gabon*. Cambridge, UK: University of Cambridge.
 82. de Merode E, Homewood K, Cowlshaw G. 2004 The value of bushmeat and other wild foods to rural households living in extreme poverty in Democratic Republic of Congo. *Biol. Conserv.* **118**, 573–581. (doi:10.1016/j.biocon.2003.10.005)
 83. Coad L, Abernethy K, Balmford A, Manica A, Airey L, Milner-Gulland EJ. 2010 Distribution and use of income from bushmeat in a rural village, central Gabon. *Conserv. Biol.* **24**, 1510–1518. (doi:10.1111/j.1523-1739.2010.01525.x)
 84. Abernethy K, Coad L, Ilambu O, Makiloulita F, Easton J, Akiak J. 2010 *Wildlife hunting, consumption and trade in the Oshwe sector of the Salonga-Lukenie-Sankuru Landscape*. Kinshasa, Democratic Republic of the Congo: WWF CARPO.
 85. Auzel P, Wilkie D. (eds) 2000 *Wildlife use in northern Congo: hunting in a commercial logging concession*. New York, NY: Columbia University Press.
 86. Carpaneto GM, Fusarii A, Okongo H. 2007 Subsistence hunting and exploitation of mammals in the Haut-Ogooué province, south-eastern Gabon. *J. Anthropol. Sci.* **85**, 183–193.
 87. Lahm S. 1994 *Hunting and wildlife in northeastern Gabon: why conservation should extend beyond protected areas*. Makoukou, Gabon: Institut de Recherche en Ecologie Tropicale.
 88. Allebone-Webb SM, Kumpel NF, Rist J, Cowlshaw G, Rowcliffe JM, Milner-Gulland EJ. 2011 Use of market data to assess bushmeat hunting sustainability in Equatorial Guinea. *Conserv. Biol.* **25**, 597–606. (doi:10.1111/j.1523-1739.2011.01681.x)
 89. Robinson JG, Redford KH (eds) 1991 *Sustainable harvest of neotropical forest mammals*. Chicago, IL: University of Chicago Press.
 90. Cowlshaw G, Mendelson S, Rowcliffe JM. 2005 Evidence for post-depletion sustainability in a mature bushmeat market. *J. Appl. Ecol.* **42**, 460–468. (doi:10.1111/j.1365-2664.2005.01046.x)
 91. Kumpel NF, Milner-Gulland EJ, Cowlshaw G, Rowcliffe JM. 2010 Assessing sustainability at multiple scales in a rotational bushmeat hunting system. *Conserv. Biol.* **24**, 861–871. (doi:10.1111/j.1523-1739.2010.01505.x)
 92. Ling S, Milner-Gulland EJ. 2006 Assessment of the sustainability of bushmeat hunting based on dynamic bioeconomic models. *Conserv. Biol.* **20**, 1294–1299. (doi:10.1111/j.1523-1739.2006.00414.x)
 93. Robinson JG, Redford KH, Bennett EL. 1999 Conservation—Wildlife harvest in logged tropical forests. *Science* **284**, 595–596. (doi:10.1126/science.284.5414.595)
 94. Papworth SK, Rist J, Coad L, Milner-Gulland EJ. 2009 Evidence for shifting baseline syndrome in conservation. *Conserv. Lett.* **2**, 93–100. (doi:10.1111/j.1755-263X.2009.00049.x)
 95. Weinbaum KZ, Brashares JS, Golden CD, Getz WM. 2013 Searching for sustainability: are assessments of wildlife harvests behind the times? *Ecol. Lett.* **16**, 99–111. (doi:10.1111/ele.12008)
 96. Newing H. 2001 Bushmeat hunting and management: implications of duiker ecology and interspecific competition. *Biodivers. Conserv.* **10**, 99–118. (doi:10.1023/A:1016671524034)
 97. Treves A. 2009 Hunting for large carnivore conservation. *J. Appl. Ecol.* **46**, 1350–1356. (doi:10.1111/j.1365-2664.2009.01729.x)
 98. Linder JM, Oates JF. 2011 Differential impact of bushmeat hunting on monkey species and implications for primate conservation in Korup National Park, Cameroon. *Biol. Conserv.* **144**, 738–745. (doi:10.1016/j.biocon.2010.10.023)
 99. Dirzo R, Mendoza E, Ortiz P. 2007 Size-related differential seed predation in a heavily defaunated neotropical rain forest. *Biotropica* **39**, 355–362. (doi:10.1111/j.1744-7429.2007.00274.x)
 100. Terborgh J, Nunez-Iturri G, Pitman NCA, Valverde FHC, Alvarez P, Swamy V, Pringle EG, Paine CET. 2008 Tree recruitment in an empty forest. *Ecology* **89**, 1757–1768. (doi:10.1890/07-0479.1)
 101. Babweteera F, Savill P, Brown N. 2007 *Balanites wilsoniana*: regeneration with and without elephants. *Biol. Conserv.* **134**, 40–47. (doi:10.1016/j.biocon.2006.08.002)
 102. Blake S, Deem SL, Mossimbo E, Maisels F, Walsh P. 2009 Forest elephants: tree planters of the Congo. *Biotropica* **41**, 459–468. (doi:10.1111/j.1744-7429.2009.00512.x)
 103. Wang BC, Sork VL, Leong MT, Smith TB. 2007 Hunting of mammals reduces seed removal and dispersal of the afro-tropical tree *Antrocaryon klaineana* (Anacardiaceae). *Biotropica* **39**, 340–347. (doi:10.1111/j.1744-7429.2007.00275.x)
 104. Harrison RD, Tan S, Ploktin JB, Slik F, Detto M, Brenes T, Itoh A, Davies SJ. 2013 Consequences of defaunation for a tropical tree community. *Ecol. Lett.* **16**, 687–694. (doi:10.1111/ele.12102)
 105. Brodie JF, Gibbs HK. 2009 Bushmeat hunting as climate threat. *Science* **326**, 364–365. (doi:10.1126/science.326_364b)
 106. Wright SJ, Stoner KE, Beckman N, Corlett RT, Dirzo R, Muller-Landau HC, Nuñez-Iturri G, Peres CA, Wang BC. 2007 The plight of large animals in tropical forests and the consequences for plant regeneration. *Biotropica* **39**, 289–291. (doi:10.1111/j.1744-7429.2007.00293.x)
 107. Nunez-Iturri G, Olsson O, Howe HF. 2008 Hunting reduces recruitment of primate-dispersed trees in Amazonian Peru. *Biol. Conserv.* **141**, 1536–1546. (doi:10.1016/j.biocon.2008.03.020)
 108. Muller-Landau HC. 2007 Predicting the long-term effects of hunting on plant species composition and diversity in tropical forests. *Biotropica* **39**, 372–384. (doi:10.1111/j.1744-7429.2007.00290.x)
 109. Beckman NG, Muller-Landau HC. 2007 Differential effects of hunting on pre-dispersal seed predation and primary and secondary seed removal of two neotropical tree species. *Biotropica* **39**, 328–339. (doi:10.1111/j.1744-7429.2007.00273.x)
 110. Holbrook KM, Loiselle BA. 2009 Dispersal in a Neotropical tree, *Virola flexuosa* (Myristicaceae): Does hunting of large vertebrates limit seed removal? *Ecology* **90**, 1449–1455. (doi:10.1890/08-1332.1)
 111. Nunez-Iturri G, Howe HF. 2007 Bushmeat and the fate of trees with seeds dispersed by large primates in a lowland rain forest in western Amazonia. *Biotropica* **39**, 348–354. (doi:10.1111/j.1744-7429.2007.00276.x)
 112. Stevenson PR. 2011 The abundance of large ateline monkeys is positively associated with the diversity of plants regenerating in neotropical forests. *Biotropica* **43**, 512–519. (doi:10.1111/j.1744-7429.2010.00708.x)
 113. Nichols E, Larsen T, Spector S, Davis AL, Escobar F, Favila M, Vulinec K. 2008 Global dung beetle response to tropical forest modification and fragmentation: a quantitative literature review and meta-analysis. *Biol. Conserv.* **141**, 1932–1939. (doi:10.1016/j.biocon.2008.06.004)
 114. Vanthomme H, Belle B, Forget PM. 2010 Bushmeat hunting alters recruitment of large-seeded plant species in Central Africa. *Biotropica* **42**, 672–679. (doi:10.1111/j.1744-7429.2010.00630.x)
 115. Andresen E, Laurance SGW. 2007 Possible indirect effects of mammal hunting on dung beetle assemblages in Panama. *Biotropica* **39**, 141–146. (doi:10.1111/j.1744-7429.2006.00239.x)
 116. Jansen PA, Muller-Landau HC, Wright SJ. 2010 Bushmeat hunting and climate: an indirect link. *Science* **327**, 330. (doi:10.1126/science.327.5961.30-a)
 117. Guimaraes PR, Galetti M, Jordano P. 2008 Seed dispersal anachronisms: rethinking the fruits extinct megafauna ate. *PLoS ONE* **3**, e1745. (doi:10.1371/journal.pone.0001745)
 118. Marquis R. 2010 The role of herbivores in terrestrial trophic cascades. In *Trophic cascades* (eds

- J Terborgh, JA Estes), 1st edn, pp. 109–124. Washington, DC: Island Press.
119. Phillips OL *et al.* 2002 Increasing dominance of large lianas in Amazonian forests. *Nature* **418**, 770–774. (doi:10.1038/nature00926)
 120. van der Heijden GMF, Phillips OL. 2009 Liana infestation impacts tree growth in a lowland tropical moist forest. *Biogeosciences* **6**, 2217–2226. (doi:10.5194/bg-6-2217-2009)
 121. Terborgh J. In press. Using Janzen–Connell to predict the consequences of defaunation and other disturbances of tropical forests. *Biol. Conserv.* (doi:10.1016/j.biocon.2013.01.015)
 122. Wright IJ *et al.* 2007 Relationships among major dimensions of plant trait variation in seven Neotropical forests. *Ann. Bot. Lond.* **99**, 1003–1015. (doi:10.1093/aob/mcl066)
 123. Voysey BC, McDonald KE, Rogers ME, Tutin CEG, Parnell RJ. 1999 Gorillas and seed dispersal in the Lopé Reserve, Gabon. II: survival and growth of seedlings. *J. Trop. Ecol.* **15**, 39–60. (doi:10.1017/S026646749900668)
 124. Cochrane EP. 2003 The need to be eaten: *Balanites wilsoniana* with and without elephant seed-dispersal. *J. Trop. Ecol.* **19**, 579–589. (doi:10.1017/S0266467403003638)
 125. Stoner KE, Riba-Hernandez P, Vulinec K, Lambert JE. 2007 The role of mammals in creating and modifying seed shadows in tropical forests and some possible consequences of their elimination. *Biotropica* **39**, 316–327. (doi:10.1111/j.1744-7429.2007.00292.x)
 126. Rudel TK. 2013 The national determinants of deforestation in sub-Saharan Africa. *Phil. Trans. R. Soc. B* **368**, 20120405. (doi:10.1098/rstb.2012.0405)
 127. Oxford Business Group. 2011 *Gabon: the report*. Oxford, UK: Oxford Business Group.
 128. Mayaux P *et al.* 2013 State and evolution of the African rainforests between 1990 and 2010. *Phil. Trans. R. Soc. B* **368**, 20120300. (doi:10.1098/rstb.2012.0300)
 129. Poulsen JR, Clark CJ, Mavah G, Elkan PW. 2009 Bushmeat supply and consumption in a tropical logging concession in Northern Congo. *Conserv. Biol.* **23**, 1597–608. (doi:10.1111/j.1523-1739.2009.01251.x)
 130. Thibault M, Blaney S. 2003 The oil industry as an underlying factor in the bushmeat crisis in Central Africa. *Conserv. Biol.* **17**, 1807–1813. (doi:10.1111/j.1523-1739.2003.00159.x)
 131. Blake S *et al.* 2008 Roadless wilderness area determines forest elephant movements in the Congo Basin. *PLoS ONE* **3**, e3546. (doi:10.1371/journal.pone.0003546)
 132. Wilkie D, Shaw E, Rotberg F, Morelli G, Auzel P. 2000 Roads, development, and conservation in the Congo basin. *Conserv. Biol.* **14**, 1614–1622. (doi:10.1046/j.1523-1739.2000.99102.x)
 133. Yackulic CB, Strindberg S, Maisels F, Blake S. 2011 The spatial structure of hunter access determines the local abundance of forest elephants (*Loxodonta africana cyclotis*). *Ecol. Appl.* **21**, 1296–1307. (doi:10.1890/09-1099.1)
 134. Poulsen JR, Clark CJ, Bolker BM. 2011 Decoupling the effects of logging and hunting on an Afrotropical animal community. *Ecol. Appl.* **21**, 1819–1836. (doi:10.1890/10-1083.1)
 135. Nasi R, Billand A, Vanvliet N. 2012 Managing for timber and biodiversity in the Congo Basin. *For. Ecol. Manag.* **268**, 103–111. (doi:10.1016/j.foreco.2011.04.005)
 136. Mertens B, Lambin EF. 2000 Land-cover-change trajectories in southern Cameroon. *Ann. Assoc. Am. Geogr.* **90**, 467–494. (doi:10.1111/0004-5608.00205)
 137. Galford GL, Soares-Filho B, Cerri CEP. 2013 Prospects for land-use sustainability on the agricultural frontier of the Brazilian Amazon. *Phil. Trans. R. Soc. B* **368**, 20120171. (doi:10.1098/rstb.2012.0171)
 138. Zhang QF, Justice CO, Jiang MX, Brunner J, Wilkie DS. 2006 A GIS-based assessment on the vulnerability and future extent of the tropical forests of the Congo Basin. *Environ. Monit. Assess.* **114**, 107–121. (doi:10.1007/s10661-006-2015-3)
 139. James R, Washington R, Rowell DP. 2013 Implications of global warming for the climate of African rainforests. *Phil. Trans. R. Soc. B* **368**, 20120298. (doi:10.1098/rstb.2012.0298)
 140. Otto FEL, Jones RG, Halladay K, Allen MR. 2013 Attribution of changes in precipitation patterns in African rainforests. *Phil. Trans. R. Soc. B* **368**, 20120299. (doi:10.1098/rstb.2012.0299)
 141. Fisher JB *et al.* 2013 African tropical rainforest net carbon dioxide fluxes in the twentieth century. *Phil. Trans. R. Soc. B* **368**, 20120376. (doi:10.1098/rstb.2012.0376)
 142. Asefi-Najafabady S, Saatchi S. 2013 Response of African humid tropical forests to recent rainfall anomalies. *Phil. Trans. R. Soc. B* **368**, 20120306. (doi:10.1098/rstb.2012.0306)
 143. Malhi Y, Aragao LEOC, Galbraith D, Huntingford C, Fisher R, Zelazowski P, Sitch S, McSweeney C, Meir P. 2009 Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest. *Proc. Natl Acad. Sci. USA* **106**, 20 610–20 615. (doi:10.1073/pnas.0804619106)
 144. Huntingford C *et al.* 2013 Simulated resilience of tropical rainforests to CO₂-induced climate change. *Nat. Geosci.* **6**, 268–273. (doi:10.1038/ngeo1741)
 145. Cochrane MA. 2003 Fire science for rainforests. *Nature* **421**, 913–919. (doi:10.1038/nature01437)
 146. Bowman DMJS *et al.* 2011 The human dimension of fire regimes on earth. *J. Biogeogr.* **38**, 2223–2236. (doi:10.1111/j.1365-2699.2011.02595.x)
 147. Tutin CEG, Fernandez M. 1993 Relationships between minimum temperature and fruit production in some tropical forest trees in Gabon. *J. Trop. Ecol.* **9**, 241–248. (doi:10.1017/S0266467400007239)
 148. Chapman CA, Chapman LJ, Struhsaker TT, Zanne AE, Clark CJ, Poulsen JR. 2005 A long-term evaluation of fruiting phenology: importance of climate change. *J. Trop. Ecol.* **21**, 31–45. (doi:10.1017/S0266467404001993)
 149. Tutin CEG, White LJ. 1998 Primates, phenology and frugivory: present past and future patterns in the Lopé Reserve, Gabon. In *Dynamics of tropical communities* (eds DH Newbury, HT Prins, N Brown), pp. 309–337. Oxford, UK: Blackwell Science.
 150. Barnes RFW, Barnes KL, Alers MPT, Blom A. 1991 Man determines the distribution of elephants in the rain-forests of northeastern Gabon. *Afr. J. Ecol.* **29**, 54–63. (doi:10.1111/j.1365-2028.1991.tb00820.x)
 151. Wilkie D, Sidle JG, Boundzanga GC, Blake S, Auzel P. 2001 Defaunation or deforestation: commercial logging and market hunting in northern Congo. In *The cutting edge: conserving wildlife in logged tropical forests* (eds R Fimbel, JG Robinson, A Grajal), pp. 375–399. New York, NY: Columbia University Press.
 152. Barnes RFW. 2002 The bushmeat boom and bust in West and Central Africa. *Oryx* **36**, 236–242. (doi:10.1017/S0030605302000443)
 153. Rist J, Milner-Gulland EJ, Cowlshaw G, Rowcliffe M. 2010 Hunter reporting of catch per unit effort as a monitoring tool in a bushmeat-harvesting system. *Conserv. Biol.* **24**, 489–499. (doi:10.1111/j.1523-1739.2010.01470.x)
 154. Hedges S. 2012 *Monitoring elephants and assessing threats: a manual for researchers, managers and conservationists*. Himayatnagar, India: Hyderabad Universities.
 155. Kuél H, Maisels F, Ancrenaz M, Williamson EA (eds) 2008 *Best practice guidelines for surveys and monitoring of great ape populations*. Gland, Switzerland: IUCN.
 156. Lewis SL *et al.* 2013 Above-ground biomass and structure of 260 African tropical forests. *Phil. Trans. R. Soc. B* **368**, 20120295. (doi:10.1098/rstb.2012.0295)
 157. Prugh LR, Hodges KE, Sinclair ARE, Brashares JS. 2008 Effect of habitat area and isolation on fragmented animal populations. *Proc. Natl Acad. Sci. USA* **105**, 20 770–20 775. (doi:10.1073/pnas.0806080105)
 158. Chazdon RL *et al.* 2009 Beyond reserves: a research agenda for conserving biodiversity in human-modified tropical landscapes. *Biotropica* **41**, 142–153. (doi:10.1111/j.1744-7429.2008.00471.x)
 159. Stokes EJ *et al.* 2010 Monitoring great ape and elephant abundance at large spatial scales: measuring effectiveness of a conservation landscape. *PLoS ONE* **5**, e10294. (doi:10.1371/journal.pone.0010294)
 160. Clark CJ, Poulsen JR, Malonga R, Elkan PW. 2009 Can wildlife management in logging concession extend the ‘conservation estate’ of tropical forests? *Conserv. Biol.* **23**, 1281–1293. (doi:10.1111/j.1523-1739.2009.01243.x)
 161. Tranquilli S *et al.* 2011 Lack of conservation effort rapidly increases African great ape extinction risk. *Conserv. Lett.* **5**, 48–55. (doi:10.1111/j.1755-263X.2011.00211.x)
 162. Stokes EJ. 2010 Improving effectiveness of protection efforts in tiger source sites: developing a framework for law enforcement monitoring using MIST. *Integr. Zool.* **5**, 363–377. (doi:10.1111/j.1749-4877.2010.00223.x)



CrossMark
[click for updates](#)

Correction

Cite this article: Abernethy KA, Coad L, Taylor G, Lee ME, Maisels F. 2013 Extent and ecological consequences of hunting in Central African rainforests in the twenty-first century. *Phil Trans R Soc B* 368: 20130494.
<http://dx.doi.org/10.1098/rstb.2013.0494>

Extent and ecological consequences of hunting in Central African rainforests in the twenty-first century

K. A. Abernethy, L. Coad, G. Taylor, M. E. Lee and F. Maisels

Phil. Trans. R. Soc. B **368**, 20120303 (5 September 2013; Published online 22 July 2013) (doi:10.1098/rstb.2012.0303)

Authors would like to acknowledge some additional supporters in their funding statement:

Funding statement. L.C. and G.T. would like to acknowledge the support of the Oxford University John Fell Fund and the Oxford Martin School. F.M. would like to acknowledge the support of WCS, and that of the US Fish and Wildlife Service and CARPE (USAID) for supporting ecological surveys that informed part of this paper.