

Norambuena F, Morais S, Estevez A, Bell JG, Tocher DR, Navarro JC, Cerda J & Duncan N (2013) Dietary modulation of arachidonic acid metabolism in senegalese sole (*Solea Sengalensis*) broodstock reared in captivity, *Aquaculture*, 372-375, pp. 80-88.

**This is the peer reviewed version of this article**

*NOTICE: this is the author's version of a work that was accepted for publication in Aquaculture. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Aquaculture, [VOL 372-375 (2013)] DOI: <http://dx.doi.org/10.1016/j.aquaculture.2012.10.035>*

**DIETARY MODULATION OF ARACHIDONIC ACID METABOLISM IN  
SENEGALESE SOLE (*SOLEA SENEGALENSIS*) BROODSTOCK REARED IN  
CAPTIVITY**

Fernando Norambuena<sup>1\*</sup>, Sofia Morais<sup>1</sup>, Alicia Estévez<sup>1</sup>, J Gordon Bell<sup>2</sup>, Douglas R. Tocher<sup>2</sup>,  
Juan C. Navarro<sup>3</sup>, Joan Cerdà<sup>4</sup> and Neil Duncan<sup>1</sup>

<sup>1</sup>IRTA-Sant Carles de la Ràpita, Ctra. Poble Nou Km 6, 43540-Sant Carles de la Ràpita,  
Tarragona, Spain

<sup>2</sup>Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling FK9 4LA,  
Scotland, UK

<sup>3</sup>Institute of Aquaculture of Torre la Sal, Spanish Council for Scientific Research (CSIC), Torre  
la Sal s/n, 12595-Cabanes, Castellón, Spain

<sup>4</sup>IRTA-Institute of Marine Sciences (CSIC), 08003 Barcelona, Spain

\*Corresponding author: Fernando Norambuena, Phone: +34- 977 745427, Fax +34-977744138  
email: [norambuena52@hotmail.com](mailto:norambuena52@hotmail.com)

**Abstract**

Previous studies have shown higher levels of arachidonic acid (20:4n-6, ARA) in testis, liver, and muscle of wild Senegalese sole (*Solea senegalensis*) compared to fish reared in captivity (first generation, G1). The present study was conducted to establish the optimal level of dietary ARA for G1 Senegalese sole broodstock, using as a reference the fatty acid profile of wild broodstock (gonads, liver and muscle). A total of 120 Senegalese sole broodstock were randomly distributed into 12 tanks and (1:1 male and female), fed in duplicate with six experimental diets containing increasing amounts of ARA (0.7%, 1.6%, 2.3%, 3.2%, 5.0%, and 6.0 % of total fatty acids) for nine months. The relative ARA levels in liver, muscle and male and female gonads at the end of the feeding period increased in a dose dependent manner. Dietary ARA was mainly incorporated and stored in testis or ovary, followed by liver and muscle. Fish fed 2.3% and 3.2% ARA showed no differences in the ARA content of testis, ovary and liver when compared to

wild fish. In male fish, a significant increase in the levels of 22:4n-6 and 22:5n-6 fatty acids was also observed, which was consistent with the up-regulation of fatty acyl elongase (*elovl5*) and desaturase (*d4fad*) transcript levels in the liver of fish fed 0.7%, 2.3% and 6% ARA. These results suggest that dietary inclusion of 3.2% ARA during periods shorter than nine months, or of 2.3% ARA for prolonged periods, can maintain optimal levels of tissue ARA in captive Senegalese sole broodstock. In addition, the data indicate that male Senegalese sole is able to elongate and desaturate ARA to 22:4n-6 and 22:5n-6, suggesting that these fatty acids may be important for male reproduction.

**Keywords:** Arachidonic acid, broodstock fish, fish nutrition, fatty acids.

## 1. Introduction

One of the most important nutritional factors for successful fish reproduction, and among the most studied, is the fatty acid arachidonic acid (20:4n-6, ARA) [1-6]. Arachidonic acid is the main precursor for production of 2-series prostaglandins (PGs) [7,8], which stimulate ovarian and testicular steroidogenesis, triggering oocyte maturation in females and milt production in males, and are involved in female sexual behaviour [9-13]. Arachidonic acid itself and its metabolites regulate cholesterol (CHOL) transfer from the outer to inner mitochondrial membrane where the P450 enzyme resides to initiate steroid hormone synthesis [11,14]. Moreover, ARA had differential effects on steroid biosynthesis. Although it stimulates testosterone production by elevating cAMP levels in a dose-dependent manner, ARA at high doses can also inhibit steroidogenesis by affecting the availability of CHOL [11,15].

Senegalese sole (*Solea senegalensis*) is a promising species for aquaculture in Southern Europe. However, an important problem encountered in this species is the fact that first generation (G1) of reared fish often fail to spawn viable eggs, contrary to wild animals, which produce eggs of sufficient quality and quantity after variable times of acclimation in captivity [16]. This is hindering the expansion of Sole aquaculture and hence several studies have been recently performed in an attempt to understand the underlying causes. Studies on wild sole broodstock showed higher levels of ARA and ARA-derived fatty acids in different tissues compared to those observed in G1 fish [17], similarly to that has been reported in other fish species [18-27]. High accumulation of ARA in sperm has also been reported in rainbow trout (*Oncorhynchus mykiss*)

57 fed diets low in docosahexaenoic acid (22:6n-3, DHA) [28], and in wild European seabass  
58 (*Dicentrarchus labrax*) [29]. In addition, previous studies on Senegalese sole showed that  
59 differences in ARA tissue content resulted in differences in cyclooxygenase (COX-2) gene  
60 expression, which was significantly up-regulated in the sperm-duct, oviduct and gills of males  
61 from wild origin compared to G1 fish [30]. Thus, wild fish showed significantly higher levels of  
62 2-series PGs compared to cultured fish, especially in testis, whereas G1 Senegalese sole, with a  
63 lower ARA tissue content, exhibited significantly higher levels of 3-series PGs and lower levels  
64 of CHOL [17], the precursor of steroid hormones in vertebrates [31]. On the other hand,  
65 Senegalese sole fed artificial diets formulated with graded ARA levels showed an increase in  
66 ARA levels in circulating blood, which in turn may induce an increase in CHOL and steroid  
67 production, especially in males [32]. Higher levels of ARA in the tissues of wild fish and  
68 increased levels in the blood of G1 fish previously fed ARA-enriched diets resulted in an  
69 increase in ARA-derived fatty acids, 22:4n-6 and 22:5n-6 [32]. A similar increase in these n-6  
70 long-chain polyunsaturated fatty acids (LC-PUFA) was observed in the sperm of wild European  
71 seabass [29]. These fatty acids are present in the cells of reproductive (i.e., seminiferous tubules,  
72 sperm) and nervous tissues in larger quantities (human, bull, boar and rabbit) [33-36] than those  
73 reported in any other fish tissue and mammals [37,38]. On the other hand, although the  
74 physiological function of these LC-PUFAs in sperm is not well known, in mammals they are  
75 considered indicators of normal testicular development, spermatogenesis, germ cell populations  
76 and fertility [36,39-42] as well as in sperm formation and transportation in the rat testicle  
77 [36,38].

78 Biologically active essential fatty acids such as ARA, eicosapentaenoic acid (20:5n-3, EPA) and  
79 DHA can be synthesized to some extent by some mammals and freshwater fish through  
80 elongation and desaturation of dietary shorter chain precursors. Carnivores and marine fish have  
81 only negligible biosynthetic capacity and hence require preformed LC-PUFA in the diet [6,43].  
82 However, it was recently demonstrated that desaturation of 22:4n-6 to 22:5n-6 may be carried  
83 out by a direct pathway involving a delta 4 desaturase in both a marine herbivorous fish [44], as  
84 well as in Senegalese sole larvae [45]. This indicates that there is more than one possible  
85 pathway for the synthesis of 22:5n-6 and DHA in vertebrates, i.e., not only the classical  
86 ‘Sprecher pathway’ [43].

Based on these latest observations, in the present study we conducted a nine-month feeding trial on broodstock G1 Senegalese sole using a standard commercial feed formulation with six graded levels of dietary ARA. The objectives were: (1) to determine the optimal dietary level of ARA for G1 Senegalese sole, using as a reference the fatty acid profile in gonads, liver and muscle of wild broodstock [17]; and (2) to investigate the regulation of fatty acyl desaturase (*d4fad*) and elongase (*elovl5*) gene expression in the liver of G1 sole fed different amounts of ARA.

## **2. Materials and Methods**

Research involving animal experimentation conformed to the principles for the use and care of laboratory animals, in agreement with the Spanish and European regulations on animal welfare (Federation of Laboratory Animal Science Associations, FELASA).

### **2.1. Fish and Diets**

One hundred and twenty Senegalese sole (four year old and  $524 \pm 11$  g average weight), reared in captivity were PIT tagged (AVID, UK) and sexed using a heterologous vitellogenin ELISA for European seabass (*Dicentrarchus labrax*) and validated for Senegalese sole [46]. The fish were distributed among twelve experimental tanks (10 fish per tank, 5 males and 5 females) and fed in duplicate standard commercial (extruded) diet with six graded ARA contents (Tables 1 and 2) for nine months (from September 2009 until May 2010). The fish were held in a recirculation system with simulated natural photoperiod and temperature (40° 37' and 40° 48' N and between 0° 21' and 0° 40' E., Tarragona, Spain), with minimum temperature observed during two weeks in January - February (13°C) and maximum temperature over twelve weeks during June - September (21°C). The fish were fed six days per week at a daily ration of 0.15- 0.3% body weight.

### **2.2. Fish Sampling**

In May 2010, seventy two fish were sacrificed by pithing the spinal cord (12 fish per dietary treatment, 6 males and 6 females) after anaesthesia with 0.3 ml L<sup>-1</sup> Aqui-S® (Scan Aqua A.S, Årnes, Norway) [47]. Gonads, liver and muscle were collected, and 2 g of liver frozen immediately in liquid nitrogen and subsequently stored at -70°C until RNA extraction, whereas the rest of the tissues for lipid and fatty acid profile were stored at -20 °C. All the fish used for

these analyses were in advanced stages of sexual maturation, females with vitellogenic oocytes and males containing spermatozoa in the seminiferous tubules.

### 2.3. Lipid and Fatty Acid Analyses

Samples of tissues and feeds were homogenized and total lipids extracted [48] and quantified gravimetrically. Tissue samples of six males and six females for each diet treatment were analyzed, and feeds were analyzed in triplicate every three months during the experiment. Fatty acid methyl esters were prepared by acid-catalyzed transmethylation [49], and extracted and purified following [50]. Methyl esters were separated and quantified by gas-liquid chromatography (Thermo Trace GC, Thermo Finningan, Milan, Italy) using a 30 m x 0.25 mm ID capillary column (BPX 70, SGE Europe Ltd., UK) with on-column injection and flame ionization detection using helium as carrier gas (1.2 mL min<sup>-1</sup> constant flow rate). Individual methyl esters were identified by comparison with known standards (Supelco Inc., Madrid) and a well-characterized fish oil, and quantified in relation to the internal standard, 21:0. The results are presented as percentage of the total fatty acids (TFA) as mean  $\pm$  standard error of the mean (SEM). Water content was calculated by drying samples at 105°C until a constant weight was obtained [51].

### 2.4. Tissue RNA Extraction and Quantitative Real-Time (qRT-PCR)

In order to study the expression of fatty acyl desaturase (*d4fad*) and elongase (*elovl5*) in liver, which is the main metabolic organ where LC-PUFA biosynthesis occurs, total RNA was extracted by organic solvent (Tri-reagent), according to the manufacturer's instructions (Ambion, Applied Biosystems). RNA quality and quantity were assessed by gel electrophoresis and spectrophotometry (NanoDrop ND-1000, Thermo Scientific, Wilmington, USA), respectively. One microgram of total RNA per sample was reverse-transcribed into cDNA using a Verso™ cDNA kit (ABgene, Surrey, UK), following the manufacturer's instructions, using a mixture of random hexamers and anchored oligo-dT (3:1, v/v). The cDNA was then diluted 50-fold with water, after a similar amount of cDNA was pooled from all samples. The expression levels of *d4fad* and *elovl5* transcripts were determined by real-time quantitative (qRT-PCR) and normalized using ubiquitin (*UBQ*) and ribosomal protein S4 (*RPS4*) expression using primers described previously by Morais et al. [45] and normalized using ubiquitin (*UBQ*) and ribosomal

protein S4 (*RPS4*) [52]. Details of the primers used can be found in Table 3. The amplification efficiency of the primer pairs was assessed using serial dilutions of cDNA pooled from the samples. All amplifications were carried out in duplicate using an Eppendorf qPCR Cycler (Stanford) in a final volume of 20  $\mu$ l containing 2  $\mu$ l (for reference genes) or 5  $\mu$ l (for *d4fad* and *elovl5*) diluted cDNA (1/50), 0.5  $\mu$ M of each primer and 10  $\mu$ l of Absolute<sup>TM</sup> qPCR SYBR<sup>®</sup> Green mix (ABgene). Every amplification experiment also included non-template controls (NTC). The RT-qPCR profiles contained an initial activation step at 95°C for 15 min, followed by 35 cycles: 15 s at 95°C, 15 s at the specific primer pair annealing  $T_m$  (Table 3), and 30 s at 72°C. After the amplification phase, a melt curve of 0.5°C increments from 75°C to 90°C was performed, enabling confirmation of the amplification of a single product in each reaction. The RT-qPCR product sizes were checked by agarose gel electrophoresis and their identity was confirmed by sequencing. No primer–dimer formation occurred in the NTC.

## 2.5. Statistical Analysis

Statistical differences in lipid and fatty acid compositions among the six experimental groups and wild fish measured previously [17] were analysed separately in males and females by one-way ANOVA followed by the post-hoc multiple comparison test Tukey's HSD, at a significance level ( $P$ ) of 0.05. Moreover, correlations among ARA and C22 fatty acids of the n-6 series were calculated at  $P = 0.05$ . The compliance of data with normality and homogeneity of variance were tested using the Kolmogorov–Smirnov and Bartlett (Chi-Sqr) tests and, when necessary, log-transformation was carried out. Fatty acid content was expressed as mean % TFA  $\pm$  SEM. Statistical analysis was performed using the Statistica<sup>®</sup> package for windows (version 6.0; StatSoft Inc, Tulsa, USA).

The relative expression of *d4fad* and *elovl5* in fish from groups A (control), C and F was normalized by the expression of *UBQ* and *RPS4* using the normalization factor calculated by geNorm<sup>®</sup> Software version 3.5 [53] and then analyzed for statistical significance using the relative expression software tool (REST-MCS<sup>®</sup>, version 2, <http://www.gene-quantification.info/>), which employs a pairwise fixed reallocation randomization test (10,000 randomizations) with efficiency correction [54].

### 3. Results

After 9 months of feeding the experimental diets the weight of females increased 1.8-fold (from  $533 \pm 13\text{g}$  to  $950 \pm 25\text{g}$ ,  $\text{SGR} = 0.20\% \text{ day}^{-1}$ ) whereas weights of males increased 1.5-fold, (from  $515 \pm 16\text{g}$  to  $755 \pm 25\text{g}$ ,  $\text{SGR} = 0.13\% \text{ day}^{-1}$ ), with a feed conversion ratio of  $1.3 \pm 0.04$ . No significant differences were noted among the six experimental groups in either growth or feed conversion.

#### 3.1. Fatty acid composition of tissues

The total fatty acid compositions of the different male and female tissues are shown in Tables 4-9. All the tissues analyzed showed a significant accumulation of ARA in a dose dependent manner. Thus, in testis the fish in groups C, D, E and F showed a significantly higher ARA content compared to the control group A, particularly group F which had 13% ARA, 2.7-fold higher than group A with 5% ARA. In the ovary, fish from group F showed a 3.2-fold higher ARA content than group A, whereas groups B, C, D and E did not show significant differences compared to A. The liver of the fish from groups fed diets C, D, E and F had higher ARA levels than control diet A both in males and females. Additionally, males from group B also had significantly higher liver ARA than those from group A, while in females there were no significant differences. The muscle of male fish from groups C, D, E and F showed significantly higher ARA levels than those from group A, whereas in females only those from groups E and F had significantly higher ARA levels than group A.

In the case of 22:4n-6 and 22:5n-6 contents, fish fed the six experimental diets showed a dose dependent and significant increase in testis (Table 4) and liver (Table 6) of male fish, whereas no differences could be found in females, although a similar trend was observed but with much smaller differences (Table 5 and 7). The relative level of 22:4n-6 in testis was significantly higher in groups B (2.2-fold), C (4.2-fold), D (4.5-fold), E (6.1-fold) and F (8.1-fold) compared with the lowest levels found in the control group A (0.2% TFA) (Table 4). The 22:5n-6 content in testis was also higher in groups C (1.5-fold), D (1.8-fold), E (2.2-fold) and F (2.4-fold) compared with the control group A (0.4% TFA). The 22:4n-6 levels in the liver of males were also significantly higher in groups E (7.1-fold) and F (9.5-fold) compared to control group A



(0.1% TFA), while 22:5n-6 was 5-fold higher in group F compared with group A (0.2% TFA) (Table 6).

As a consequence of the increasing ARA levels, the EPA/ARA ratio was significantly reduced in similar dose-dependent manner in gonads (Tables 4 and 5), liver (Tables 6 and 7) and muscle (Tables 8 and 9) of both males and females. The increase of ARA in gonad, liver and muscle of males resulted in a concomitant significant increase in total n-6 PUFA whereas a significant reduction in total n-3 PUFA was only observed in testis.

### **3.2. Comparison of tissue ARA levels with wild fish**

The tissue ARA levels in the present study were compared those of wild broodstock reported previously [17], and different results were obtained depending on the tissue (Fig. 1). In gonads ARA was significantly higher in males compared with females however in liver and muscle no differences in ARA content between males and females were observed. The levels of ARA in testis of fish from groups E and F showed significantly higher accumulation of ARA compared with those from wild fish. However, ARA levels in testis from groups A, B, C, and D were similar and not significantly different to those from wild fish. In ovary no differences in ARA between the experimental groups and wild fish were observed. ARA content in liver (males and females) from groups A and B were significantly lower than the wild group. However, groups C, D, E and F showed no differences compared to the wild group. In muscle of males the ARA levels in all the fish groups were significantly lower compared to wild fish, and the same as in females from groups A, B, C and D. However, females from groups E and F showed similar ARA levels to that found in the wild.

### **3.3. Elongase and desaturase gene expression (RT-qPCR)**

The expression of *d4fad* and *elovl5* increased in a dose dependent manner in fish fed diets C and F in comparison to the control (A) treatment, but only in males (Fig. 2). Levels of *d4fad* transcripts were significantly higher in the liver of male fish from groups C (3.2-fold) and F (6.8-fold). On the other hand, expression of *elovl5* increased significantly in group F, being 5.3-fold higher than in group A, but not in group C (1.8-fold increase).

#### 4. Discussion

One of the main bottlenecks to the culture of Senegalese sole, a species with high market value and interest for aquaculture diversification in Southern Europe, is the poor reproductive performance of G1 fish. Previous studies have indicated that the problem is more likely associated with male sperm production and quality than female performance [55-58]. When comparing the fatty acid profile of cultured and wild fish major differences were found in the contents of ARA which, due to its important multiple roles in reproduction, was deemed a potential factor associated with reproductive failure of fish produced in captivity. Hence, the objective of the present study was to test a gradient of dietary ARA to determine the levels that would raise ARA tissue contents in G1 fish to those similar to wild fish. In addition, given previous observations that increased ARA in the tissue and blood of fish raise the levels of its elongated and desaturated products, 22:4n-6 and 22:5n-6, [17,32], the pathway of LC-PUFA biosynthesis was also studied by assessing the expression levels of two genes coding for Elovl5 and d4Fad, which are two key enzymes in this pathway.

The results obtained show that the relative content of ARA in tissues of G1 Senegalese sole generally correlated with dietary ARA levels, increasing in a dose-dependent manner in both sexes. Feeding fish with diets containing 2.3% and 3.2% ARA levels (diets C and D, respectively) resulted in similar ARA contents in testis, ovary and liver compared to wild fish. Muscle, on the other hand, did not accumulate ARA at the same rate as liver and gonads and only the female fish fed diets containing 5% or higher ARA achieved similar ARA levels in muscle as the wild fish. A previous study on dietary ARA preference of Senegalese sole broodstock using a self-feeding system over 16 months revealed that fish regulate the ingestion of ARA to around 3.0% TFA in diet and the resulting ARA content in the tissues was 8.9% TFA in testis, 4% in ovary and 2.5% in liver [59]. These values were similar to those found previously in wild Senegalese sole [17] and also to those obtained in the present study in the testis, ovary and liver of G1 fish in groups C and D. Thus, the ARA accumulation in the muscle was lower than in the self-feeding experiment, except for females of groups E and F. Considering that the self-feeding experiment was conducted for 16 months and the present study lasted only 9 months, optimal dietary ARA level seems to be dependent on the duration of the feeding period. If fish are fed for a period shorter than nine months it might be appropriate to use at least 3.2%

ARA in the diet. For longer feeding periods, 2.3% ARA in the diet might be sufficient for Senegalese sole G1, such that no differences in ARA content could be found in the testis and ovary of the G1 fish compared to wild fish, or fish in the self-feeding experiment. Alorend [4], studied ARA requirements for G1 Atlantic halibut over 3 years and suggested 2.3% as the optimal dietary content, as this level led to the longest milt production period and highest fecundity. This result is likely due to ARA being rapidly incorporated into the reproductive tissues [4].

Previous studies on ARA requirements for broodstock fish showed that the optimal dietary level is species-specific, with higher or lower ARA levels producing detrimental effects in reproductive physiology [2-4,11,12,15,30,60]. A significant increase in the production of steroids was observed in Senegalese sole males fed 3.2% ARA [32], and the highest egg production was obtained using diets with 3.6% ARA for Japanese flounder (*Paralichthys olivaceus*) [3]. On the other hand, a study with Atlantic cod (*Gadus morhua*) found that a diet with 4% ARA, increased the production of estradiol and extended the length of the spawning season [60]. However, negative effects on steroid production, fecundity, egg and larval quality have been observed when fish were fed high ARA levels. Japanese flounder fed 7.3% ARA exhibited a significant reduction in egg and larval quality [3] whereas Atlantic halibut fed 3.2% ARA showed a delay in the spawning season [4]. Other effects such as an earlier estradiol peak in Senegalese sole [32], and an earlier peak in estradiol and vitellogenesis in Atlantic cod [60] were also observed. More research is required to establish the effects of dietary ARA on sperm and oocyte quality in *S. senegalensis* and the feeding time required to incorporate ARA into the reproductive organs, but the present study suggests that dietary levels between 2.3% and 3.2%, depending on the duration of the feeding, might improve the reproductive performance of Senegalese sole G1.

The results obtained in the present study also showed that dietary ARA was preferentially transferred to and accumulated in the gonads (testis and ovary), followed by the liver and muscle, similarly to what had been previously observed in Atlantic halibut, white seabream, black seabream and silver pomfret [4,22,23,61,62].

A positive correlation between the level of ARA and the concentrations of 22:4n-6 and 22:5n-6 in testis, liver and muscle of wild fish had been shown previously [17], and it was suggested that this probably occurred from metabolism (elongation and desaturation) of ARA. A similar pattern

of 22:4n-6 and 22:5n-6 concentration in the tissues after graded dietary ARA was observed in the testis and liver of males in the present study. Accumulation of these fatty acids in the muscle of wild black seabream [23] and Senegalese sole [17], in the gonads of seabass [63], seabream [21], silver pomfret [62] and in the sperm of seabass [29] and rainbow trout [28,29] have been reported previously, with the accumulation of 22:5n-6 being suggested as deriving from local production and uptake from the circulatory system [64-66]. These fatty acids are found in storage lipids in testis of mammals [38] and are considered indicators of normal testicular and sperm condition [36-42]. Although the physiological function of these LC-PUFA in sperm is not well understood, Lenzie et al. [36] suggested that they are involved in sperm formation and in fertilization and, in mammals, there is an increase in the degree of fatty acid metabolism and desaturation during spermatogenesis and sperm maturation [42]. However, the function of 22:4n-6 and 22:5n-6 in fish and their effects on reproduction and spermatogenesis have not been established. It has been suggested that 22:5n-6 is accumulated in the testis as an ARA reservoir, being retro-converted into ARA by hydrogenation and subsequent oxidation [37,64,65,67]. Cultured fish fed commercial extruded diets show a significantly lower accumulation of 22:5n-6 and 22:4n-6 compared to wild fish [17]. Nonetheless, in the present study, when fish were fed increasing ARA levels, a parallel increase in these ARA-derived fatty acids was observed, especially in the liver and testis. Recently, the cloning and characterization of *elovl5* and *d4fad* transcripts in Senegalese sole revealed that this species is able to elongate ARA to 22:4n-6 and then directly desaturate this substrate to 22:5n-6 [45]. Although the activity of this pathway could not be assessed *in vivo*, results from this experiment suggest that it is physiologically relevant given that ARA induced a marked dose-dependent up-regulation in the expression of both *elovl5* and *d4fad* in liver, with *elovl5* transcripts increasing 5.3-fold in males fed 6.0% ARA in relation to those fed 0.7%, and *d4fad* being up-regulated 3.2- and 6.8-fold in fish fed diets containing 2.3% and 6.0% ARA, respectively. However, an interesting observation was that the up-regulation of these two genes was only observed in males, which was also clearly reflected in the fatty acid composition of the tissues, where significant changes in 22:4n-6 and 22:5n-6 contents between treatments were only observed in testis, liver and flesh of males whereas in female tissues only ARA levels were significantly different between fish fed the experimental diets. Gender differences in liver desaturase expression have also been observed in Wistar rats fed an n-3 PUFA enriched diet [68]. In that case, desaturation activity was significantly increased

in the females compared to males. It was also shown that female rats have higher plasma DHA concentrations than males [69]. It is conceivable that the LC-PUFA biosynthesis pathway in female fish liver is also more directed to the provision of DHA for later incorporation into eggs and, at least in post-larvae, the expression of *d4fad* was up-regulated at lower dietary levels of n-3 LC-PUFA (mainly DHA and EPA) [45]. However, in the present experiment only the levels of ARA varied significantly between diets and hence *elovl5* and *d4fad* expression was associated with the conversion of ARA into 22:4n-6 and subsequently to 22:5n-6 in G1 male fish, both of which are important fatty acids involved in testis and sperm composition in mammals [39-42]. Further studies are necessary to understand the importance of these two fatty acids and their physiological function in male fish reproduction, but the present results suggest that they may be as important in Senegalese sole as in higher vertebrates.

## 5. Conclusion

Based on previous data on ARA content in wild Senegalese sole [17], and considering the results presented here, fish fed either 2.3% or 3.2% ARA enriched diets showed levels of ARA in testis, ovary and liver compared to those of wild fish. Thus, diets with 3.2% ARA for feeding periods up to nine months or 2.3% ARA diets for prolonged feeding periods are suggested. The ARA was preferentially transferred and conserved in the gonads (testis and ovary), followed by the liver and muscle. The increase in the expression of *elovl5* and *d4fad* transcripts in liver in response to dietary ARA content and a parallel increase in tissue 22:4n-6 and 22:5n-6 levels suggest the ability of Senegalese sole to elongate ARA to 22:4n-6 followed by desaturation to 22:5n-6, suggesting an important role of these ARA-derived fatty acids in male fish reproduction. Further studies are required to establish the dietary ARA effect on reproductive performance of Senegalese sole and the time required for effective incorporation of ARA into reproductive organs.

## Acknowledgments

Thanks to N. Gras, G. Macia and J.R. Dick for technical support and G. Rosenlund from Skretting ARC (Norway) for formulating and providing the diets used in the study. This research was funded by Ministerio de Agricultura, Pesca, Alimentación y Medio Ambiente -JACUMAR Project Lenguado coordinated nationally by Pedro Cañavate and in IRTA by N.D. and (INIA)-

FEDER project RTA2005-00113-00-00 coordinated by N.D. F.N. thanks Agència de Gestió d'Ajuts Universitaris i de Recerca (AGAUR) for the PhD grant he was awarded.

## 6. Reference

- [1] Sargent J, Bell G, McEvoy L, Tocher D, Estevez A. Recent developments in the essential fatty acid nutrition of fish. *Aquaculture* 1999; **177**: 191-199.
- [2] Mazorra C, Bruce M, Bell JG, Davie A, Alorend E, Jordan N *et al.* Dietary lipid enhancement of broodstock reproductive performance and egg and larval quality in Atlantic halibut (*Hippoglossus hippoglossus*). *Aquaculture* 2003; **227**: 21-33.
- [3] Furuita H, Yamamoto T, Shima T, Suzuki N, Takeuchi T. Effect of arachidonic acid levels in broodstock diet on larval and egg quality of Japanese flounder *Paralichthys olivaceus*. *Aquaculture* 2003; **220**: 725-735.
- [4] Alorend E. The effect of dietary arachidonic acid concentration on Atlantic Halibut (*Hippoglossus hippoglossus*) broodstock performance, assessment of egg, milt and larval quality. *PhD Thesis, Institute of Aquaculture University of Stirling, UK*; 2004. p. 176.
- [5] Meunpol O, Meejing P, Piyatiratitivorakul S. Maturation diet based on fatty acid content for male *Penaeus monodon* (Fabricius) broodstock. *Aquac Res* 2005; **36**: 1216-1225.
- [6] Tocher DR. Fatty acid requirements in ontogeny of marine and freshwater fish. *Aquac Res* 2010; **41**: 717-732.
- [7] Smith WL, Murphy RC, Dennis E, Vance JEV. The eicosanoids: cyclooxygenase, lipoxygenase, and epoxygenase pathways. *New Compr Biochem* 2002; **36**: 341-371.
- [8] Tocher DR. Metabolism and functions of lipids and fatty acids in teleost fish. *Rev Fish Sci* 2003; **11**: 107-184.
- [9] Van der Kraak G, Chang JP. Arachidonic acid stimulates steroidogenesis in goldfish preovulatory ovarian follicles. *Gen Comp Endocr* 1990; **77**: 221-228.
- [10] Wade MG, Van der Kraak G. Arachidonic acid and prostaglandin E2 stimulate testosterone production by goldfish testis in vitro. *Gen Comp Endocr* 1993; **90**: 109-118.
- [11] Mercure F, Van der Kraak G. Inhibition of gonadotropin-stimulated ovarian steroid production by polyunsaturated fatty acids in teleost fish. *Lipids* 1995; **30**: 547-554.
- [12] Sorbera LA, Asturiano JF, Carrillo M, Zanuy S. Effects of polyunsaturated fatty acids and prostaglandins on oocyte maturation in a marine teleost, the European Sea bass (*Dicentrarchus labrax*). *Biol Reprod* 2001; **64**: 382-389.
- [13] Sorensen PW, Stacey NE. Brief review of fish pheromones and discussion of their possible uses in the control of non-indigenous teleost fishes. *New Zeal J Mar Fresh* 2004; **38**: 399-417.
- [14] Wang X, Stocco DM. The decline in testosterone biosynthesis during male aging: A consequence of multiple alterations. *Mol Cell Endocrinol* 2005; **238**: 1-7.
- [15] Mercure F, Van der Kraak G. Mechanisms of Action of Free Arachidonic Acid on Ovarian Steroid Production in the Goldfish. *Gen Comp Endocr* 1996; **102**: 130-140.
- [16] Carazo I, Martin I, Chereguini O, Mañanós E, Duncan N. The absence of reproductive behaviour in cultured (G1 generation) Senegalese sole (*Solea senegalensis*) explains poor reproductive performance. *Proceeding of 5th Workshop on the Cultivation of Soles* Centre of Marine Sciences, University of the Algarve, Faro, Portugal, 5-7 April; 2011. p. 12
- [17] Norambuena F, Estevez A, Bell G, Carazo I, Duncan N. Proximate and fatty acid composition in muscle, liver and gonads of wild versus cultured broodstock of Senegalese sole (*Solea senegalensis*). *Aquaculture* 2012; **356-357**: 176-185.

- [18] Harrell RM, Woods LC. Comparative fatty acid composition of eggs from domesticated and wild striped bass (*Morone saxatilis*). *Aquaculture* 1995; **133**: 225-233.
- [19] Sheikh-Eldin M, De Silva SS, Anderson TA, Gooley G. Comparison of fatty acid composition of muscle, liver, mature oocytes, and diets of wild and captive Macquarie perch, *Macquaria australasica*, broodfish. *Aquaculture* 1996; **144**: 201-216.
- [20] Silversand C, Norberg B, Haux C. Fatty-acid composition of ovulated eggs from wild and cultured turbot (*Scophthalmus maximus*) in relation to yolk and oil globule lipids. *Mar Biol* 1996; **125**: 269-278.
- [21] Cejas JR, Almansa E, Villamandos JE, Badía P, Bolaños A, Lorenzo A. Lipid and fatty acid composition of ovaries from wild fish and ovaries and eggs from captive fish of white sea bream (*Diplodus sargus*). *Aquaculture* 2003; **216**: 299-313.
- [22] Cejas JR, Almansa E, Jerez S, Bolaños A, Samper M, Lorenzo A. Lipid and fatty acid composition of muscle and liver from wild and captive mature female broodstocks of white seabream, *Diplodus sargus*. *Comp Biochem Physiol B* 2004; **138**: 91-102.
- [23] Rodríguez C, Acosta C, Badía P, Cejas JR, Santamaría FJ, Lorenzo A. Assessment of lipid and essential fatty acids requirements of black seabream (*Spondylusoma cantharus*) by comparison of lipid composition in muscle and liver of wild and captive adult fish. *Comp Biochem Physiol B* 2004; **139**: 619-629.
- [24] Salze G, Tocher DR, Roy WJ, Robertson DA. Egg quality determinants in cod (*Gadus morhua* L.): egg performance and lipids in eggs from farmed and wild broodstock. *Aquac Res* 2005; **36**: 1488-1499.
- [25] Blanchet C, Lucas M, Julien P, Morin R, Gingras S, Dewailly E. Fatty acid composition of wild and farmed Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*). *Lipids* 2005; **40**: 529-531.
- [26] Aslan SS, Guven KC, Gezgin T, Alpaslan M, Tekinay A. Comparison of fatty acid contents of wild and cultured rainbow trout *Oncorhynchus mykiss* in Turkey. *Fisheries Sci* 2007; **73**: 1195-1198.
- [27] Lund I, Steenfeldt SJ, Suhr KI, Hansen BW. A comparison of fatty acid composition and quality aspects of eggs and larvae from cultured and wild broodstock of common sole (*Solea solea* L.). *Aquacult Nutr* 2008; **14**: 544-555.
- [28] Vassallo-Agius R, Watanabe T, Yoshizaki G, Satoh S, Takeuchi Y. Quality of eggs and spermatozoa of rainbow trout fed an n-3 essential fatty acid-deficient diet and its effects on the lipid and fatty acid components of eggs, semen and livers. *Fisheries Sci* 2001; **67**: 818-827.
- [29] Bell MV, Dick JR, Thrush M, Navarro JC. Decrease 20:4n-6/20:5n-3 ratio in sperm from cultured sea bass, *Dicentrarchus labrax*, broodstock compared with wild fish *Aquaculture* 1996; **144**: 189-199.
- [30] Norambuena F, Mackenzie S, Bell JG, Callol A, Estevéz A, Duncan N. Prostaglandin (F and E, 2- and 3-series) production and cyclooxygenase (COX-2) gene expression of wild and cultured broodstock of Senegalese sole (*Solea senegalensis*). *Gen Comp Endocr* 2012; **177**: 256-262.
- [31] Baron SF, Hylemon PB. Biotransformation of bile acids, cholesterol, and steroids hormones. In: White RL MaBA (ed). *Gastrointestinal microbiology*. London, England: Chapman and Hall; 1997. p. 663.
- [32] Norambuena F. Senegalese sole (*Solea senegalensis*) broodstock nutrition: arachidonic acid (20:4n-6, ARA) and reproductive physiology. *PhD Thesis, Institute of Animal Biology, Vegetal and Ecology*: Autonomous University of Barcelona, Spain; 2012. p. 187.
- [33] Ahaluwalia B, Holman RT. Fatty acid composition of lipid of bull, boar, rabbit and human semen. *J Reprod Fertil* 1969; **18**: 431-437.
- [34] Tinoco J. Dietary requirements and functions of  $\alpha$ -linolenic acid in animals. *Prog Lipid Res* 1982; **21**: 1-45.

- [35] Picardo M, Gramatico P, De Luca C, Passi S, Nazzaro-Porro M. Differential effect of azelaic acid in melanoma cell cultures. *Pigm Cell Res* 1990; **20**: 142-149.
- [36] Lenzi A, Picardo M, Gandini L, Dondero F. Lipids of the sperm plasma membrane: from polyunsaturated fatty acids considered as markers of sperm function to possible scavenger therapy. *Hum Reprod Update* 1996; **2**: 246-256.
- [37] Bridges RB, Coniglio JG. The metabolism of 4,7,10,13,16-[5-14c]docosapentaenoic acid in the testis of the rat. *BBA-Lipid Lipid Met* 1970; **218**: 29-35.
- [38] Ayala S, Gaspar G, Brenner RR, Peluffo RIO, Kunau W. Fate of linoleic, arachidonic, and docosa-7,10,13,16-tetraenoic acids in rat testicles. *J Lipid Res* 1973; **14**: 296-305.
- [39] Leat WMF, Northrop CA, Harrison FA, Cox RW. Effect of dietary linoleic and linolenic acids on testicular development in the rat. *Exp Physiol* 1983; **68**: 221-231.
- [40] MacDonald ML, Rogers QR, Morris JG, Cupps PT. Effects of linoleate and arachidonate deficiencies on reproduction and spermatogenesis in the cat. *J Nutr* 1984; **114**: 719-726.
- [41] Lenzi A, Gandini L, Maresca V, Rago R, Sgro P, Dondero F *et al.* Fatty acid composition of spermatozoa and immature germ cells. *Mol Hum Reprod* 2000; **6**: 226-231.
- [42] Furland NE, Maldonado EN, Aresti PA, Avelaño MI. Changes in lipids containing long- and very long-chain polyunsaturated fatty acids in cryptorchid rat testes. *Biol Reprod* 2007; **77**: 181-188.
- [43] Sprecher H. Metabolism of highly unsaturated n-3 and n-6 fatty acids. *Biochim Biophys Acta* 2000; **1486**: 219-231.
- [44] Li Y, Monroig O, Zhang L, Wang S, Zheng X, Dick JR *et al.* Vertebrate fatty acyl desaturase with  $\Delta 4$  activity. *P Natl Acad Sci USA* 2010; **107**: 16840-16845.
- [45] Morais S, Castanheira F, Martinez-Rubio L, Conceição LEC, Tocher DR. Long chain polyunsaturated fatty acid synthesis in a marine vertebrate: Ontogenetic and nutritional regulation of a fatty acyl desaturase with delta 4 activity. *BBA- Mol Cell Biol L* 2012; **1821**: 660-671.
- [46] Mañanós E, Núñez J, Zanuy S, Carrillo M, Le Menn F. Sea bass (*Dicentrarchus labrax* L.) vitellogenin. II-Validation of an enzyme-linked immunosorbent assay (ELISA). *Comp Biochem Physiol B* 1994; **107**: 217-223.
- [47] Norambuena F, Carazo I, Duncan N, Estevez A, Roque A. Anesthesia of broodstock mediterranean fish. *Humane Slaughter Association Centenary International Symposium: Recent Advances in the Welfare of Livestock at Slaughter* Portsmouth Historic Dockyard, Portsmouth, UK, 30 June - 1 July; 2011. p. 30.
- [48] Folch JM, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 1957; **226**: 497-509.
- [49] Christie WW. Lipid Analysis. Pergamon, Oxford, UK; 1982. p. 207
- [50] Tocher DR, Harvie DG. Fatty acid compositions of the major phosphoglycerides from fish neural tissues; (n-3) and (n-6) polyunsaturated fatty acids in rainbow trout (*Salmo gairdneri*) and cod (*Gadus morhua*) brains and retinas. *Fish Physiol Biochem* 1988; **5**: 229-239.
- [51] AOAC. Crude protein content by the Dumas procedure, by means of a Nitrogen/protein FP-528 determinator and moisture. *Official Methods of Analysis*. Washington, USA: Association of Official Analytical Chemist; 2000.
- [52] Infante C, Matsuoka M, Asensio E, Canavate J, Reith M, Manchado M. Selection of housekeeping genes for gene expression studies in larvae from flatfish using real-time PCR. *BMC Mol Biol* 2008; **9**: 28.
- [53] Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A *et al.* Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Gen Biol* 2002; **3**: 34.31-34.11.




- [54] Pfaffl MW, Horgan GW, Dempfle L. Relative expression software tool (REST©) for group-wise comparison and statistical analysis of relative expression results in real-time PCR. *Nucleic Acids Res* 2004.
- [55] Guzmán JM, Cal R, García-López A, Chereguini O, Kight K, Olmedo M *et al.* Effects of in vivo treatment with the dopamine antagonist pimozide and gonadotropin-releasing hormone agonist (GnRHa) on the reproductive axis of Senegalese sole (*Solea senegalensis*). *Com Biochem Phys A* 2011; **158**: 235-245.
- [56] Agulleiro MJ, Anguis V, Cañavate JP, Martinez-Rodriguez G, Mylonas CC, Cerda J. Induction of spawning of captive-reared Senegal sole (*Solea senegalensis*) using different administration methods for gonadotropin-releasing hormone agonist. *Aquaculture* 2006; **257**: 511-524.
- [57] Agulleiro MJ, Scott AP, Duncan N, Mylonas CC, Cerdà J. Treatment of GnRHa-implanted Senegalese sole (*Solea senegalensis*) with 11-ketoandrostenedione stimulates spermatogenesis and increases sperm motility. *Comp Biochem Phys A* 2007; **147**: 885-892.
- [58] Forné I, Agulleiro MJ, Asensio E, Abián J, Cerdà J. 2-D DIGE analysis of Senegalese sole (*Solea senegalensis*) testis proteome in wild-caught and hormone-treated F1 fish. *Proteomics* 2009; **9**: 2171-2181.
- [59] Norambuena F, Sánchez-Vásquez J, Estevez A, Carazo I, Duncan N. Self-selection of diets with different content of arachidonic acid by Senegalese sole (*Solea senegalensis*) broodstock. *Aquaculture*, submitted 2012.
- [60] Norberg B, Hamre K, Araujo P, van der Meeren T, Rosenlund G. Effect of graded levels of arachidonic acid on reproductive physiology, spawning performance, and egg quality of atlantic cod *5th Fish & shellfish larviculture symposium, European Aquaculture Society* Oostende, Belgium, 7-11 September; 2009.
- [61] Pérez MJ, Rodríguez C, Cejas JR, Martín MV, Jerez S, Lorenzo A. Lipid and fatty acid content in wild white seabream (*Diplodus sargus*) broodstock at different stages of the reproductive cycle. *Comp Biochem Physiol B* 2007; **146**: 187-196.
- [62] Huang X, Yin Y, Shi Z, Li W, Zhou H, Lv W. Lipid content and fatty acid composition in wild-caught silver pomfret (*Pampus argenteus*) broodstocks: Effects on gonad development. *Aquaculture* 2010; **310**: 192-199.
- [63] Bell JG, Farndale BM, Bruce MP, Navas JM, Carillo M. Effects of broodstock dietary lipid on fatty acid compositions of eggs from sea bass (*Dicentrarchus labrax*). *Aquaculture* 1997; **149**: 107-119.
- [64] Rosenthal MD, Garcia MC, Jones MR, Sprecher H. Retroconversion and D4 desaturation of docosatetraenoate (22:4(n-6)) and docosapentaenoate (22:5(n-3)) by human cells in culture. *BBA-Lipid Lipid Met* 1991; **1083**: 29-36.
- [65] Tam P, Umeda-Sawada R, Yaguchi T, Akimoto K, Kiso Y, Igarashi O. The metabolism and distribution of docosapentaenoic acid (n-6) in rats and rat hepatocytes. *Lipids* 2000; **35**: 71-75.
- [66] Tam PS, Sawada R, Cui Y, Matsumoto A, Fujiwara Y. The metabolism and distribution of docosapentaenoic acid (n-6) in the liver and testis of growing rats. *Biosci Biotechnol Biochem* 2008; **72**: 2548-2554.
- [67] Kunau W. Partial degradation of 4.7.10.13.16-docosapentaenoic acid in rat liver. *Febs Lett* 1971; **16**: 54-56.
- [68] Extier A, Langelier Bnd, Perruchot M-Hln, Guesnet P, Van Veldhoven PP, Lavialle M *et al.* Gender affects liver desaturase expression in a rat model of n-3 fatty acid repletion. *J Nutr Biochem* 2010; **21**: 180-187.
- [69] Childs CE, Romeu-Nadal M, Burdge GC, Calder PC. Gender differences in the n-3 fatty acid content of tissues. *P Nut Soc* 2008; **67**: 19-27.


**Fig. 1** Arachidonic acid content (20:4n-6, ARA) in a ) gonads, b) liver and c) muscle of wild fish (Norambuena et al., 2012a) compared with cultured fish fed with different dietary ARA levels (A= 0.7% ARA, B=1.6% ARA, C= 2.3% ARA, D= 3.2% ARA, E= 5.0% ARA and F= 6.0% ARA. Different letters indicate significant differences (ANOVA,  $P<0.05$ ,  $N=6$ ) between fish groups. Capital letters are used for males and small letters for females. (\*) Indicate significant differences ( $P<0.05$ ) between males and females.

**Fig. 2** Relative expression (RT-qPCR) of fatty acyl (a) elongase (elovl5) and (b) desaturase (d4fad) in the liver of female and male Senegalese sole in relation to group A (control), normalized by the expression of UBQ and RPS4 (reference genes). (\*) Denote significant differences of groups C and F with respect to control (group A), calculated by REST ( $P<0.05$ ,  $N=6$ ).

Fig. 1



592 Fig. 2



593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619

620 **Table 1.** *Ingredients and proximate composition of the experimental diets (A, B, C, D, E and F)*

Ingredients (g/Kg)	A	B	C	D	E	F
Fish meal <sup>1</sup>	645.0	645.0	645.0	645.0	645.0	645.0
Wheat gluten <sup>2</sup>	120.0	120.0	120.0	120.0	120.0	120.0
Wheat <sup>3</sup>	125.8	125.8	125.8	125.8	125.8	125.8
Fish oil <sup>4</sup>	80.0	76.0	71.8	67.6	63.2	59.0
Vevodar <sup>5</sup>	0.0	4.0	8.2	12.4	16.8	21.0
Premixes <sup>6</sup>	29.2	29.2	29.2	29.2	29.2	29.2
Analysed values						
Moisture, %	8.0	7.8	8.3	8.4	8.6	8.3
Crude protein, % DM <sup>7</sup>	61,2	61,4	61,6	61,7	61,8	62,2
Crude fat, % DM	13.8	14.1	14.4	13.7	14.1	14.3

<sup>1</sup> LT fish meal, Skretting, Stavanger, Norway

<sup>2</sup> Cargill Nordic, Charlottenlund, Denmark

<sup>3</sup> Skretting, Stavanger, Norway

<sup>4</sup> Scandinavian fish oil, Skretting, Stavanger, Norway

<sup>5</sup> Contains 35% arachidonic acid, DSM Food Specialities, Delft, The Netherlands

<sup>6</sup> Include micronutrients, vitamin and mineral supplementation. Trouw Nutrition, Boxmeer, Netherlands, proprietary composition Skretting ARC

<sup>7</sup> Dry matter

**Table 2.** Lipid, fatty acid content and fatty acid composition (% TFA  $\pm$  SEM) of the diets used (A, B, C, D, E and F) for feeding G1 Senegalese sole (*Solea senegalensis*). Columns assigned different letters were significantly different (ANOVA,  $P < 0.05$ ,  $N = 3$ )

	A	B	C	D	E	F
TFA ( $\mu\text{g mg}^{-1}$ L)	880 $\pm$ 87	702 $\pm$ 34	820 $\pm$ 74	861 $\pm$ 82	961 $\pm$ 19	911 $\pm$ 94
Fatty acid composition (%TFA)						
14:0	2.5 $\pm$ 1.2	3.6 $\pm$ 2.8	3.8 $\pm$ 2.1	3.6 $\pm$ 2.3	3.7 $\pm$ 2.8	4.0 $\pm$ 2.9
16:0	14.9 $\pm$ 1.1	15.3 $\pm$ 3.7	17.6 $\pm$ 2.6	15.8 $\pm$ 1.8	15.0 $\pm$ 2.8	16.1 $\pm$ 2.7
18:0	1.8 $\pm$ 0.9	2.4 $\pm$ 0.9	2.6 $\pm$ 0.7	2.6 $\pm$ 0.6	3.0 $\pm$ 0.7	3.2 $\pm$ 0.4
Total SFA	19.4 $\pm$ 1.5	21.7 $\pm$ 5.7	24.4 $\pm$ 4.2	22.3 $\pm$ 3.6	21.9 $\pm$ 4.8	23.4 $\pm$ 5.2
16:1n-7	4.9 $\pm$ 0.8	4.8 $\pm$ 2.0	5.0 $\pm$ 0.8	4.9 $\pm$ 0.8	4.3 $\pm$ 1.8	4.4 $\pm$ 1.2
18:1n-9	15.0 $\pm$ 1.3	15.7 $\pm$ 2.4	16.9 $\pm$ 1.9	15.9 $\pm$ 2.0	15.1 $\pm$ 2.7	15.7 $\pm$ 0.6
18:1n-7	1.2 $\pm$ 2.0	0.9 $\pm$ 1.6	1.0 $\pm$ 1.7	0.8 $\pm$ 1.4	0.8 $\pm$ 1.4	1.0 $\pm$ 1.7
20:1n-9	7.2 $\pm$ 1.5	7.1 $\pm$ 0.3	7.2 $\pm$ 1.1	6.3 $\pm$ 0.5	6.9 $\pm$ 0.7	6.6 $\pm$ 0.4
22:1n-9	3.5 $\pm$ 6.0	3.9 $\pm$ 6.7	2.9 $\pm$ 5.0	3.0 $\pm$ 5.2	3.9 $\pm$ 6.7	2.9 $\pm$ 5.1
Total MUFA	32.3 $\pm$ 10.5	32.7 $\pm$ 4.7	33.4 $\pm$ 4.4	31.3 $\pm$ 4.9	31.4 $\pm$ 5.5	31.1 $\pm$ 6.0
18:2n-6	5.9 $\pm$ 0.7	6.4 $\pm$ 0.7	6.0 $\pm$ 0.9	6.6 $\pm$ 0.4	5.9 $\pm$ 0.1	7.2 $\pm$ 0.7
20:4n-6, ARA	0.7 $\pm$ 0.3 <sup>c</sup>	1.6 $\pm$ 0.6 <sup>c</sup>	2.3 $\pm$ 0.8 <sup>bc</sup>	3.2 $\pm$ 0.7 <sup>b</sup>	5.0 $\pm$ 0.6 <sup>a</sup>	6.0 $\pm$ 0.1 <sup>a</sup>
22:4n-6	0.1 $\pm$ 0.1	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0	0.0 $\pm$ 0.0
22:5n-6	0.3 $\pm$ 0.6	0.3 $\pm$ 0.2	0.2 $\pm$ 0.3	0.2 $\pm$ 0.3	0.2 $\pm$ 0.2	0.3 $\pm$ 0.4
Total n-6 PUFA	9.3 $\pm$ 3.4	8.4 $\pm$ 1.3	8.9 $\pm$ 1.5	10.6 $\pm$ 0.8	12.4 $\pm$ 1.0	14.0 $\pm$ 1.6
18:3n-3	1.3 $\pm$ 0.2	1.4 $\pm$ 0.2	1.3 $\pm$ 0.3	1.3 $\pm$ 0.2	1.2 $\pm$ 0.1	1.2 $\pm$ 0.1
18:4n-3	2.2 $\pm$ 0.3	2.0 $\pm$ 0.1	1.8 $\pm$ 0.3	2.0 $\pm$ 0.1	1.8 $\pm$ 0.2	1.7 $\pm$ 0.2
20:4n-3	0.7 $\pm$ 0.1	0.7 $\pm$ 0.1	0.6 $\pm$ 0.1	0.7 $\pm$ 0.0	0.6 $\pm$ 0.1	0.6 $\pm$ 0.1
20:5n-3, EPA	13.0 $\pm$ 8.4	16.8 $\pm$ 5.3	15.9 $\pm$ 5.4	16.4 $\pm$ 5.4	14.8 $\pm$ 6.0	14.7 $\pm$ 4.9
22:5n-3, DPA	1.6 $\pm$ 0.4	3.0 $\pm$ 2.3	2.0 $\pm$ 1.0	2.3 $\pm$ 1.3	4.6 $\pm$ 5.7	2.0 $\pm$ 0.9
22:6n-3, DHA	14.4 $\pm$ 2.0	13.0 $\pm$ 2.1	11.3 $\pm$ 2.5	13.0 $\pm$ 1.6	11.1 $\pm$ 0.6	11.3 $\pm$ 2.3
Total n-3 PUFA	39.0 $\pm$ 11.0	37.1 $\pm$ 0.7	33.4 $\pm$ 2.9	35.8 $\pm$ 2.1	34.3 $\pm$ 1.9	31.5 $\pm$ 1.4
Total PUFA	48.3 $\pm$ 11.2	45.5 $\pm$ 1.4	42.3 $\pm$ 2.6	46.4 $\pm$ 1.4	46.6 $\pm$ 2.0	45.5 $\pm$ 1.7
EPA/ARA	23.6 $\pm$ 18.2 <sup>a</sup>	12.4 $\pm$ 6.6 <sup>a</sup>	7.8 $\pm$ 4.2 <sup>ab</sup>	5.5 $\pm$ 2.5 <sup>ab</sup>	3.0 $\pm$ 1.3 <sup>b</sup>	2.4 $\pm$ 0.8 <sup>b</sup>
EPA/DHA	0.9 $\pm$ 0.5	1.3 $\pm$ 0.6	1.5 $\pm$ 0.8	1.3 $\pm$ 0.5	1.3 $\pm$ 0.6	1.4 $\pm$ 0.7
DHA/ARA	23.5 $\pm$ 9.0 <sup>a</sup>	8.8 $\pm$ 1.6 <sup>b</sup>	5.1 $\pm$ 1.1 <sup>bc</sup>	4.1 $\pm$ 0.5 <sup>bc</sup>	2.2 $\pm$ 0.3 <sup>bc</sup>	1.9 $\pm$ 0.4 <sup>c</sup>
n-3/n-6	4.6 $\pm$ 2.3	4.5 $\pm$ 0.7	3.8 $\pm$ 0.8	3.4 $\pm$ 0.4	2.8 $\pm$ 0.3	2.3 $\pm$ 0.3

L: lipids, DW: dry weight, TFA: total fatty acids, ARA: arachidonic acid, DPA: docosapentaenoic acid, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid.

**Table 3. Sequences of PCR primers utilized in this study [45,52]**

Transcript	Primer name	Sequence	Amplicon	Tm <sup>4</sup>	Accession no.
<i>d4fad</i>	Δ4fad -Solea-F8	AAGCCTCTGCTGATTGGAGA	131 bp <sup>3</sup>	60	JN673546
	Δ4fad-Solea-R5	GGCTGAGCTTGAAACAGACC			
<i>Elov15</i>	Elov15-Solea-F3	TTTCATGTTTTTGCACTGC	161 bp	60	JN793448
	Elov15-Solea-R3	GACACCTTTAGGCTCGGTTTT			
<i>UBQ</i> <sup>1</sup>	qUBQ-F	AGCTGGCCCAGAAATATAACTGCGACA	93 bp	70	AB291588
	qUBQ-R	ACTTCTTCTTGCGGCAGTTGACAGCAC			
<i>RPS4</i> <sup>2</sup>	qRPS4-F	GTGAAGAAGCTCCTTGTCGGCACCA	83 bp	70	AB291557
	qRPS4-R	AGGGGGTCGGGGTAGCGGATG			

<sup>1</sup>UBQ: Ubiquitin

<sup>2</sup>RPS4: 40S ribosomal protein S4

<sup>3</sup>bp: base pairs

<sup>4</sup>Tm: annealing temperature.

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

**Table 4.** Lipid, fatty acid content and fatty acid composition (% TFA  $\pm$  SEM) of testis of Senegalese sole fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,  $P < 0.05$ ,  $N = 6$ )

	A	B	C	D	E	F
TL (mg g <sup>-1</sup> DW)	78 $\pm$ 7	68 $\pm$ 22	66 $\pm$ 20	81 $\pm$ 3	76 $\pm$ 22	81 $\pm$ 17
TFA ( $\mu$ g mg <sup>-1</sup> L)	503 $\pm$ 46	533 $\pm$ 52	546 $\pm$ 39	561 $\pm$ 56	538 $\pm$ 34	520 $\pm$ 13
Fatty acid composition (%TFA)						
14:0	0.7 $\pm$ 0.1	0.6 $\pm$ 0.1	0.9 $\pm$ 0.1	0.7 $\pm$ 0.1	0.6 $\pm$ 0.1	0.6 $\pm$ 0.1
16:0	17.8 $\pm$ 0.9	17.8 $\pm$ 0.6	18.9 $\pm$ 0.6	19.8 $\pm$ 0.4	17.7 $\pm$ 1.1	19.0 $\pm$ 0.9
18:0	7.0 $\pm$ 0.5	6.4 $\pm$ 0.4	6.3 $\pm$ 0.8	6.7 $\pm$ 0.7	6.8 $\pm$ 0.6	6.4 $\pm$ 0.3
Total SFA	26.3 $\pm$ 0.8	26.3 $\pm$ 1.7	26.9 $\pm$ 1.2	28.0 $\pm$ 0.8	25.4 $\pm$ 1.7	26.5 $\pm$ 0.9
16:1n-7	3.3 $\pm$ 0.4	2.6 $\pm$ 0.1	3.2 $\pm$ 0.4	3.2 $\pm$ 0.3	2.4 $\pm$ 0.3	3.6 $\pm$ 0.3
18:1n-9	15.8 $\pm$ 1.4	14.2 $\pm$ 1.6	15.1 $\pm$ 1.8	14.9 $\pm$ 0.6	13.0 $\pm$ 1.0	17.3 $\pm$ 1.2
18:1n-7	6.1 $\pm$ 1.7	6.5 $\pm$ 0.6	6.7 $\pm$ 0.7	5.2 $\pm$ 1.5	7.4 $\pm$ 0.6	6.0 $\pm$ 1.6
20:1n-9	1.9 $\pm$ 0.3	1.7 $\pm$ 0.4	1.8 $\pm$ 0.2	1.3 $\pm$ 0.2	1.8 $\pm$ 0.2	1.5 $\pm$ 0.1
22:1n-9	0.5 $\pm$ 0.1	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.1	0.3 $\pm$ 0.1	0.3 $\pm$ 0.1
Total MUFA	27.8 $\pm$ 1.8	24.9 $\pm$ 2.5	26.1 $\pm$ 1.1	24.2 $\pm$ 1.3	24.6 $\pm$ 0.9	28.9 $\pm$ 1.1
18:2n-6	8.5 $\pm$ 0.5	8.0 $\pm$ 0.2	6.9 $\pm$ 0.2	8.1 $\pm$ 0.3	7.0 $\pm$ 0.4	7.0 $\pm$ 0.4
20:4n-6, ARA	5.0 $\pm$ 0.5 <sup>d</sup>	7.1 $\pm$ 0.8 <sup>cd</sup>	8.6 $\pm$ 1.4 <sup>c</sup>	9.5 $\pm$ 0.3 <sup>bc</sup>	12.5 $\pm$ 1.0 <sup>ab</sup>	13.4 $\pm$ 0.5 <sup>a</sup>
22:4n-6	0.2 $\pm$ 0.1 <sup>d</sup>	0.4 $\pm$ 0.1 <sup>c</sup>	0.8 $\pm$ 0.1 <sup>bc</sup>	0.8 $\pm$ 0.1 <sup>b</sup>	1.1 $\pm$ 0.1 <sup>ab</sup>	1.5 $\pm$ 0.1 <sup>a</sup>
22:5n-6, DPA	0.4 $\pm$ 0.1 <sup>d</sup>	0.5 $\pm$ 0.1 <sup>cd</sup>	0.6 $\pm$ 0.0 <sup>bcd</sup>	0.7 $\pm$ 0.1 <sup>abc</sup>	0.9 $\pm$ 0.1 <sup>ab</sup>	1.0 $\pm$ 0.1 <sup>a</sup>
Total n-6 PUFA	16.1 $\pm$ 0.5 <sup>c</sup>	16.6 $\pm$ 0.8 <sup>c</sup>	18.8 $\pm$ 1.5 <sup>bc</sup>	19.6 $\pm$ 0.5 <sup>ab</sup>	22.2 $\pm$ 0.6 <sup>ab</sup>	22.3 $\pm$ 0.2 <sup>a</sup>
18:3n-3	0.4 $\pm$ 0.0	0.4 $\pm$ 0.0	0.4 $\pm$ 0.1	0.4 $\pm$ 0.0	0.3 $\pm$ 0.0	0.3 $\pm$ 0.1
18:4n-3	0.2 $\pm$ 0.1	0.1 $\pm$ 0.0	0.2 $\pm$ 0.1	0.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0
20:4n-3	1.5 $\pm$ 0.7	0.8 $\pm$ 0.3	0.4 $\pm$ 0.1	0.1 $\pm$ 0.1	0.4 $\pm$ 0.0	0.2 $\pm$ 0.0
20:5n-3, EPA	4.1 $\pm$ 0.5	5.1 $\pm$ 0.6	3.6 $\pm$ 0.2	3.4 $\pm$ 0.8	3.9 $\pm$ 0.4	2.9 $\pm$ 0.3
22:5n-3	3.6 $\pm$ 0.3	4.4 $\pm$ 0.5	3.7 $\pm$ 0.4	3.9 $\pm$ 0.4	3.5 $\pm$ 0.3	2.9 $\pm$ 0.2
22:6n-3, DHA	16.5 $\pm$ 1.2	18.8 $\pm$ 0.5	18.9 $\pm$ 1.7	19.2 $\pm$ 0.5	17.8 $\pm$ 0.5	15.1 $\pm$ 0.8
Total n-3 PUFA	27.4 $\pm$ 1.7 <sup>a</sup>	30.7 $\pm$ 1.6 <sup>a</sup>	27.3 $\pm$ 2.0 <sup>a</sup>	27.4 $\pm$ 0.4 <sup>a</sup>	26.1 $\pm$ 0.6 <sup>a</sup>	20.7 $\pm$ 0.8 <sup>b</sup>
Total PUFA	43.5 $\pm$ 1.5 <sup>ab</sup>	47.3 $\pm$ 2.3 <sup>ab</sup>	46.1 $\pm$ 0.7 <sup>ab</sup>	46.9 $\pm$ 0.6 <sup>ab</sup>	48.4 $\pm$ 0.6 <sup>a</sup>	43.0 $\pm$ 0.8 <sup>b</sup>
EPA/ARA	0.8 $\pm$ 0.1 <sup>a</sup>	0.7 $\pm$ 0.1 <sup>ab</sup>	0.5 $\pm$ 0.1 <sup>bc</sup>	0.4 $\pm$ 0.1 <sup>c</sup>	0.3 $\pm$ 0.1 <sup>c</sup>	0.2 $\pm$ 0.0 <sup>c</sup>
EPA/DHA	0.2 $\pm$ 0.0	0.3 $\pm$ 0.0	0.2 $\pm$ 0.0	0.2 $\pm$ 0.0	0.2 $\pm$ 0.0	0.2 $\pm$ 0.0
DHA/ARA	3.5 $\pm$ 0.5	2.3 $\pm$ 0.6	3.0 $\pm$ 1.3	2.0 $\pm$ 0.1	1.5 $\pm$ 0.1	0.9 $\pm$ 0.2
n-3/n-6	1.7 $\pm$ 0.1 <sup>a</sup>	1.8 $\pm$ 0.1 <sup>a</sup>	1.5 $\pm$ 0.3 <sup>ab</sup>	1.4 $\pm$ 0.0 <sup>ab</sup>	1.2 $\pm$ 0.1 <sup>b</sup>	0.9 $\pm$ 0.0 <sup>c</sup>

TL: Total lipids, L: lipids, DW: dry weight, TFA: total fatty acids, ARA: arachidonic acid, DPA: docosapentaenoic acid, EPA: eicosapentaenoic acid, DHA: docosahexaenoic acid. Diet A= 0.7, B= 1.6, C= 2.3, D= 3.2, E= 5.0 and F= 6.0% TFA. Data within a row assigned different letters were significantly different (ANOVA,  $P < 0.05$ ,  $N = 6$ ).



**Table 5.** Lipid, fatty acid content and fatty acid composition (% TFA  $\pm$  SEM) of ovary of Senegalese sole fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,  $P < 0.05$ ,  $N = 6$ )

	A	B	C	D	E	F
TL (mg g <sup>-1</sup> DW)	108 $\pm$ 20	100 $\pm$ 22	125 $\pm$ 10	127 $\pm$ 6	131 $\pm$ 9	130 $\pm$ 8
TFA ( $\mu$ g mg <sup>-1</sup> L)	503 $\pm$ 46	533 $\pm$ 52	546 $\pm$ 39	561 $\pm$ 56	538 $\pm$ 34	520 $\pm$ 13
Fatty acid composition (%TFA)						
14:0	2.0 $\pm$ 0.2	1.8 $\pm$ 0.3	1.7 $\pm$ 0.4	2.0 $\pm$ 0.2	2.0 $\pm$ 0.1	1.9 $\pm$ 0.4
16:0	19.1 $\pm$ 1.0	19.8 $\pm$ 1.5	18.2 $\pm$ 0.9	19.0 $\pm$ 1.0	18.7 $\pm$ 0.7	19.6 $\pm$ 0.5
18:0	4.2 $\pm$ 0.4	4.3 $\pm$ 1.4	4.4 $\pm$ 0.8	3.2 $\pm$ 0.3	3.5 $\pm$ 0.1	4.8 $\pm$ 0.6
Total SFA	25.5 $\pm$ 0.9	28.1 $\pm$ 2.9	25.3 $\pm$ 1.1	24.5 $\pm$ 1.4	24.9 $\pm$ 1.0	27.4 $\pm$ 0.2
16:1n-7	5.3 $\pm$ 0.5	4.8 $\pm$ 0.7	4.6 $\pm$ 0.6	6.2 $\pm$ 0.4	5.0 $\pm$ 0.2	4.6 $\pm$ 0.7
18:1n-9	16.5 $\pm$ 1.1	14.9 $\pm$ 1.0	15.7 $\pm$ 1.3	13.3 $\pm$ 2.8	14.9 $\pm$ 0.8	16.2 $\pm$ 0.8
18:1n-7	3.8 $\pm$ 0.4	3.8 $\pm$ 0.3	4.4 $\pm$ 0.7	3.1 $\pm$ 0.1	3.8 $\pm$ 0.4	3.3 $\pm$ 1.1
20:1n-9	2.9 $\pm$ 0.3	2.5 $\pm$ 0.5	2.6 $\pm$ 0.5	2.9 $\pm$ 0.2	3.1 $\pm$ 0.3	2.6 $\pm$ 0.4
22:1n-9	1.6 $\pm$ 0.3	1.8 $\pm$ 0.2	1.8 $\pm$ 0.6	1.6 $\pm$ 0.2	1.4 $\pm$ 0.5	1.4 $\pm$ 0.4
Total MUFA	30.7 $\pm$ 1.4	26.6 $\pm$ 2.1	28.9 $\pm$ 1.8	25.1 $\pm$ 2.7	28.4 $\pm$ 1.1	27.4 $\pm$ 1.2
18:2n-6	7.8 $\pm$ 0.4	8.4 $\pm$ 0.4	7.1 $\pm$ 0.7	7.9 $\pm$ 0.5	7.1 $\pm$ 0.3	7.5 $\pm$ 0.5
20:4n-6, ARA	2.0 $\pm$ 0.9 <sup>b</sup>	2.9 $\pm$ 1.0 <sup>b</sup>	3.8 $\pm$ 1.3 <sup>ab</sup>	4.1 $\pm$ 0.1 <sup>ab</sup>	5.1 $\pm$ 0.2 <sup>ab</sup>	6.4 $\pm$ 1.6 <sup>a</sup>
22:4n-6	0.3 $\pm$ 0.1	0.4 $\pm$ 0.0	0.5 $\pm$ 0.1	0.5 $\pm$ 0.0	0.5 $\pm$ 0.2	0.7 $\pm$ 0.2
22:5n-6, DPA	0.3 $\pm$ 0.1	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	0.5 $\pm$ 0.0	0.6 $\pm$ 0.0	0.6 $\pm$ 0.1
Total n-6 PUFA	11.3 $\pm$ 1.3	12.9 $\pm$ 1.1	14.1 $\pm$ 2.0	13.2 $\pm$ 0.8	14.1 $\pm$ 0.5	15.9 $\pm$ 1.8
18:3n-3	0.6 $\pm$ 0.2	0.6 $\pm$ 0.2	0.6 $\pm$ 0.2	1.0 $\pm$ 0.1	0.9 $\pm$ 0.0	0.8 $\pm$ 0.1
18:4n-3	0.6 $\pm$ 0.1	0.9 $\pm$ 0.4	0.5 $\pm$ 0.1	0.8 $\pm$ 0.1	0.8 $\pm$ 0.0	0.6 $\pm$ 0.1
20:4n-3	1.4 $\pm$ 0.7	1.1 $\pm$ 0.5	0.4 $\pm$ 0.1	0.5 $\pm$ 0.1	0.6 $\pm$ 0.0	0.6 $\pm$ 0.1
20:5n-3, EPA	2.8 $\pm$ 0.5	2.0 $\pm$ 0.8	2.9 $\pm$ 0.7	2.8 $\pm$ 0.2	2.3 $\pm$ 0.0	2.7 $\pm$ 0.4
22:5n-3	4.1 $\pm$ 0.4	4.3 $\pm$ 0.6	3.9 $\pm$ 0.3	4.3 $\pm$ 0.4	4.3 $\pm$ 0.1	4.0 $\pm$ 0.3
22:6n-3, DHA	20.5 $\pm$ 1.5	21.2 $\pm$ 3.4	22.8 $\pm$ 2.9	23.3 $\pm$ 1.3	22.7 $\pm$ 0.8	20.0 $\pm$ 0.8
Total n-3 PUFA	31.0 $\pm$ 1.5	30.8 $\pm$ 2.0	31.1 $\pm$ 2.0	35.8 $\pm$ 1.4	31.6 $\pm$ 0.8	28.0 $\pm$ 1.2
Total PUFA	42.4 $\pm$ 1.6	43.6 $\pm$ 1.7	45.2 $\pm$ 1.9	49.0 $\pm$ 1.4	45.7 $\pm$ 0.9	43.9 $\pm$ 1.3
EPA/ARA	1.9 $\pm$ 0.0 <sup>a</sup>	0.9 $\pm$ 0.2 <sup>b</sup>	0.8 $\pm$ 0.1 <sup>b</sup>	0.7 $\pm$ 0.0 <sup>b</sup>	0.4 $\pm$ 0.0 <sup>b</sup>	0.5 $\pm$ 0.0 <sup>b</sup>
EPA/DHA	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0
DHA/ARA	17.3 $\pm$ 0.2	12.3 $\pm$ 5.1	12.0 $\pm$ 5.8	6.0 $\pm$ 0.2	4.5 $\pm$ 0.3	3.4 $\pm$ 0.9
n-3/n-6	2.9 $\pm$ 0.2	2.5 $\pm$ 0.3	2.5 $\pm$ 0.4	2.8 $\pm$ 0.2	2.3 $\pm$ 0.1	1.9 $\pm$ 0.3

Abbreviations as in Table 4.

**Table 6.** Lipid, fatty acid content and fatty acid composition (% TFA  $\pm$  SEM) of liver of male Senegalese sole fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,  $P < 0.05$ ,  $N = 6$ )

	A	B	C	D	E	F
TL (mg g <sup>-1</sup> DW)	485 $\pm$ 39	472 $\pm$ 41	477 $\pm$ 29	455 $\pm$ 43	457 $\pm$ 36	450 $\pm$ 22
TFA ( $\mu$ g mg <sup>-1</sup> L)	562 $\pm$ 60	538 $\pm$ 109	555 $\pm$ 59	566 $\pm$ 66	504 $\pm$ 69	539 $\pm$ 50
Fatty acid composition (%TFA)						
14:0	3.9 $\pm$ 0.7	2.8 $\pm$ 0.8	3.8 $\pm$ 0.5	2.2 $\pm$ 1.1	2.9 $\pm$ 0.7	3.4 $\pm$ 0.7
16:0	19.9 $\pm$ 0.4	17.8 $\pm$ 0.8	18.6 $\pm$ 0.5	19.4 $\pm$ 1.2	16.6 $\pm$ 0.9	16.7 $\pm$ 1.5
18:0	3.7 $\pm$ 0.7	3.5 $\pm$ 0.5	2.7 $\pm$ 0.2	2.5 $\pm$ 0.4	2.5 $\pm$ 0.3	3.0 $\pm$ 0.3
Total SFA	28.8 $\pm$ 1.1	24.8 $\pm$ 1.0	25.7 $\pm$ 1.1	24.4 $\pm$ 1.7	22.5 $\pm$ 1.1	23.5 $\pm$ 2.3
16:1n-7	7.3 $\pm$ 0.6	6.5 $\pm$ 1.0	8.8 $\pm$ 0.5	7.4 $\pm$ 0.4	6.2 $\pm$ 0.7	6.1 $\pm$ 1.7
18:1n-9	20.9 $\pm$ 1.3	17.1 $\pm$ 2.2	21.9 $\pm$ 1.3	21.1 $\pm$ 1.4	18.3 $\pm$ 1.0	20.4 $\pm$ 1.2
18:1n-7	3.7 $\pm$ 0.0	3.3 $\pm$ 0.3	3.6 $\pm$ 0.3	3.8 $\pm$ 0.1	3.3 $\pm$ 0.2	3.5 $\pm$ 0.9
20:1n-9	3.5 $\pm$ 0.6	3.2 $\pm$ 0.7	3.7 $\pm$ 0.4	3.8 $\pm$ 0.6	3.8 $\pm$ 0.4	2.7 $\pm$ 0.3
22:1n-9	3.0 $\pm$ 0.4	2.9 $\pm$ 0.4	2.9 $\pm$ 0.5	3.1 $\pm$ 0.8	3.4 $\pm$ 0.6	2.8 $\pm$ 0.4
Total MUFA	29.9 $\pm$ 5.8	30.7 $\pm$ 3.8	40.5 $\pm$ 2.6	45.8 $\pm$ 4.5	33.7 $\pm$ 2.1	33.7 $\pm$ 2.0
18:2n-6	6.7 $\pm$ 0.5	6.7 $\pm$ 1.0	6.4 $\pm$ 0.5	6.8 $\pm$ 0.7	7.9 $\pm$ 0.4	7.5 $\pm$ 0.4
20:4n-6, ARA	0.5 $\pm$ 0.1 <sup>d</sup>	1.2 $\pm$ 0.2 <sup>c</sup>	1.5 $\pm$ 0.1 <sup>c</sup>	1.8 $\pm$ 0.1 <sup>bc</sup>	2.4 $\pm$ 0.2 <sup>ab</sup>	2.8 $\pm$ 0.4 <sup>a</sup>
22:4n-6	0.1 $\pm$ 0.0 <sup>c</sup>	0.1 $\pm$ 0.1 <sup>c</sup>	0.5 $\pm$ 0.0 <sup>bc</sup>	0.5 $\pm$ 0.2 <sup>bc</sup>	0.9 $\pm$ 0.1 <sup>ab</sup>	1.2 $\pm$ 0.1 <sup>a</sup>
22:5n-6, DPA	0.2 $\pm$ 0.1 <sup>b</sup>	0.3 $\pm$ 0.1 <sup>b</sup>	0.6 $\pm$ 0.1 <sup>ab</sup>	0.6 $\pm$ 0.1 <sup>ab</sup>	0.9 $\pm$ 0.1 <sup>ab</sup>	1.1 $\pm$ 0.1 <sup>a</sup>
Total n-6 PUFA	7.8 $\pm$ 0.6	8.4 $\pm$ 1.1	9.0 $\pm$ 0.6	9.0 $\pm$ 1.1	12.6 $\pm$ 0.6	12.8 $\pm$ 0.8
18:3n-3	0.8 $\pm$ 0.1	0.8 $\pm$ 0.1	0.8 $\pm$ 0.0	0.6 $\pm$ 0.1	0.9 $\pm$ 0.1	0.8 $\pm$ 0.1
18:4n-3	0.4 $\pm$ 0.0	0.5 $\pm$ 0.1	0.3 $\pm$ 0.1	0.3 $\pm$ 0.1	0.5 $\pm$ 0.0	0.3 $\pm$ 0.1
20:4n-3	0.6 $\pm$ 0.1	0.4 $\pm$ 0.1	0.6 $\pm$ 0.0	0.4 $\pm$ 0.1	0.6 $\pm$ 0.1	0.6 $\pm$ 0.1
20:5n-3, EPA	2.7 $\pm$ 0.7	2.2 $\pm$ 0.8	2.7 $\pm$ 0.8	1.7 $\pm$ 0.8	2.3 $\pm$ 0.6	3.2 $\pm$ 0.7
22:5n-3	4.3 $\pm$ 0.4	4.8 $\pm$ 0.8	4.2 $\pm$ 0.7	4.0 $\pm$ 0.8	5.2 $\pm$ 0.2	4.7 $\pm$ 0.4
22:6n-3, DHA	20.2 $\pm$ 3.7	23.8 $\pm$ 4.1	18.8 $\pm$ 1.9	17.3 $\pm$ 2.5	22.0 $\pm$ 2.0	23.0 $\pm$ 4.8
Total n-3 PUFA	29.3 $\pm$ 3.3	33.5 $\pm$ 3.3	24.2 $\pm$ 2.2	24.4 $\pm$ 4.3	30.7 $\pm$ 2.1	29.6 $\pm$ 3.8
Total PUFA	39.4 $\pm$ 3.0	41.7 $\pm$ 3.6	34.8 $\pm$ 2.4	29.9 $\pm$ 2.6	42.5 $\pm$ 1.8	42.5 $\pm$ 3.9
EPA/ARA	7.4 $\pm$ 0.9 <sup>a</sup>	2.6 $\pm$ 0.9 <sup>b</sup>	1.7 $\pm$ 0.7 <sup>b</sup>	1.4 $\pm$ 0.7 <sup>b</sup>	1.3 $\pm$ 0.2 <sup>b</sup>	1.4 $\pm$ 0.1 <sup>b</sup>
EPA/DHA	0.2 $\pm$ 0.1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.1	0.1 $\pm$ 0.0	0.1 $\pm$ 0.1
DHA/ARA	32.9 $\pm$ 2.5 <sup>a</sup>	20.0 $\pm$ 5.6 <sup>b</sup>	7.1 $\pm$ 2.9 <sup>b</sup>	5.8 $\pm$ 2.3 <sup>b</sup>	6.2 $\pm$ 1.7 <sup>b</sup>	6.0 $\pm$ 1.9 <sup>b</sup>
n-3/n-6	3.8 $\pm$ 0.5	3.5 $\pm$ 1.1	2.7 $\pm$ 0.2	2.2 $\pm$ 0.1	2.3 $\pm$ 0.2	2.3 $\pm$ 0.4

Abbreviations as in Table 4.

**Table 7.** Lipid, fatty acid content and fatty acid composition (% TFA  $\pm$  SEM) of liver of female Senegalese sole fed with six different diets (A, B, C, D, E and F) for nine months (ANOVA,  $P < 0.05$ ,  $N = 6$ )

	A	B	C	D	E	F
TL (mg g <sup>-1</sup> DW)	442 $\pm$ 83	436 $\pm$ 35	419 $\pm$ 27	365 $\pm$ 30	361 $\pm$ 40	406 $\pm$ 33
TFA ( $\mu$ g mg <sup>-1</sup> L)	482 $\pm$ 94	417 $\pm$ 85	542 $\pm$ 20	534 $\pm$ 27	456 $\pm$ 25	427 $\pm$ 94
Fatty acid composition (%TFA)						
14:0	4.3 $\pm$ 0.6	4.2 $\pm$ 0.7	3.7 $\pm$ 0.7	4.3 $\pm$ 0.7	3.9 $\pm$ 0.6	3.9 $\pm$ 1.0
16:0	21.9 $\pm$ 0.7	19.6 $\pm$ 1.4	18.5 $\pm$ 1.3	21.0 $\pm$ 0.9	21.5 $\pm$ 0.6	19.8 $\pm$ 0.8
18:0	3.7 $\pm$ 1.0	2.8 $\pm$ 0.2	1.9 $\pm$ 0.1	2.1 $\pm$ 0.2	2.9 $\pm$ 0.2	3.2 $\pm$ 0.5
Total SFA	31.1 $\pm$ 1.5	26.9 $\pm$ 2.0	24.2 $\pm$ 1.7	28.3 $\pm$ 1.3	28.8 $\pm$ 1.0	27.0 $\pm$ 2.3
16:1n-7	9.5 $\pm$ 1.7	7.8 $\pm$ 0.6	7.3 $\pm$ 1.6	9.8 $\pm$ 0.4	7.8 $\pm$ 0.4	7.3 $\pm$ 1.3
18:1n-9	18.2 $\pm$ 4.4	19.4 $\pm$ 2.0	21.0 $\pm$ 2.0	21.4 $\pm$ 0.6	22.5 $\pm$ 0.1	20.2 $\pm$ 1.8
18:1n-7	4.2 $\pm$ 0.4	4.1 $\pm$ 0.3	4.7 $\pm$ 0.2	4.7 $\pm$ 0.4	5.1 $\pm$ 0.2	4.9 $\pm$ 0.2
20:1n-9	2.9 $\pm$ 0.5	3.5 $\pm$ 0.8	3.2 $\pm$ 0.6	2.5 $\pm$ 0.2	3.8 $\pm$ 0.4	2.6 $\pm$ 0.5
22:1n-9	2.2 $\pm$ 0.7	2.3 $\pm$ 0.3	2.4 $\pm$ 0.6	2.0 $\pm$ 0.7	2.2 $\pm$ 0.5	2.5 $\pm$ 0.3
Total MUFA	32.2 $\pm$ 5.9	35.2 $\pm$ 3.4	39.8 $\pm$ 2.8	38.8 $\pm$ 1.1	39.7 $\pm$ 0.2	33.7 $\pm$ 3.7
18:2n-6	8.0 $\pm$ 1.1	8.0 $\pm$ 0.1	9.6 $\pm$ 0.5	9.7 $\pm$ 0.6	9.1 $\pm$ 0.2	8.7 $\pm$ 0.6
20:4n-6, ARA	0.6 $\pm$ 0.3 <sup>c</sup>	1.1 $\pm$ 0.2 <sup>c</sup>	1.7 $\pm$ 0.1 <sup>b</sup>	1.8 $\pm$ 0.2 <sup>b</sup>	2.3 $\pm$ 0.1 <sup>ab</sup>	3.0 $\pm$ 0.1 <sup>a</sup>
22:4n-6	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	0.4 $\pm$ 0.0	0.4 $\pm$ 0.0	0.4 $\pm$ 0.1	0.5 $\pm$ 0.1
22:5n-6, DPA	0.3 $\pm$ 0.0	0.3 $\pm$ 0.0	0.5 $\pm$ 0.0	0.5 $\pm$ 0.1	0.5 $\pm$ 0.1	0.6 $\pm$ 0.1
Total n-6 PUFA	9.0 $\pm$ 0.9 <sup>b</sup>	9.6 $\pm$ 0.4 <sup>b</sup>	12.6 $\pm$ 0.4 <sup>a</sup>	12.6 $\pm$ 0.6 <sup>a</sup>	13.9 $\pm$ 0.4 <sup>a</sup>	13.0 $\pm$ 0.6 <sup>a</sup>
18:3n-3	0.9 $\pm$ 0.1	0.9 $\pm$ 0.0	0.8 $\pm$ 0.2	0.9 $\pm$ 0.1	0.9 $\pm$ 0.1	0.8 $\pm$ 0.1
18:4n-3	0.8 $\pm$ 0.2	0.5 $\pm$ 0.0	0.6 $\pm$ 0.2	0.9 $\pm$ 0.1	0.9 $\pm$ 0.2	0.7 $\pm$ 0.1
20:4n-3	0.5 $\pm$ 0.1	0.5 $\pm$ 0.1	0.6 $\pm$ 0.0	0.3 $\pm$ 0.1	0.5 $\pm$ 0.1	0.5 $\pm$ 0.0
20:5n-3, EPA	1.2 $\pm$ 0.3	1.9 $\pm$ 0.4	1.2 $\pm$ 0.2	1.2 $\pm$ 0.2	0.8 $\pm$ 0.3	1.1 $\pm$ 0.1
22:5n-3	2.8 $\pm$ 0.4	3.8 $\pm$ 0.4	3.7 $\pm$ 0.5	3.0 $\pm$ 0.7	3.5 $\pm$ 0.4	3.7 $\pm$ 0.7
22:6n-3, DHA	16.9 $\pm$ 3.9	14.4 $\pm$ 1.4	15.7 $\pm$ 2.3	13.4 $\pm$ 1.9	12.0 $\pm$ 1.3	16.4 $\pm$ 6.0
Total n-3 PUFA	22.6 $\pm$ 4.0	22.9 $\pm$ 1.7	23.1 $\pm$ 3.0	19.5 $\pm$ 1.9	17.2 $\pm$ 0.8	22.3 $\pm$ 5.8
Total PUFA	31.7 $\pm$ 4.6	37.3 $\pm$ 5.4	35.7 $\pm$ 3.3	32.1 $\pm$ 2.3	31.1 $\pm$ 1.2	38.9 $\pm$ 5.9
EPA/ARA	3.3 $\pm$ 1.0 <sup>a</sup>	2.0 $\pm$ 0.6 <sup>ab</sup>	0.7 $\pm$ 0.2 <sup>b</sup>	0.6 $\pm$ 0.1 <sup>b</sup>	0.4 $\pm$ 0.1 <sup>b</sup>	0.3 $\pm$ 0.0 <sup>b</sup>
EPA/DHA	0.1 $\pm$ 0.0	0.2 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0	0.1 $\pm$ 0.0
DHA/ARA	45.1 $\pm$ 17 <sup>a</sup>	13.4 $\pm$ 3.6 <sup>ab</sup>	9.6 $\pm$ 2.0 <sup>b</sup>	7.5 $\pm$ 1.3 <sup>b</sup>	5.4 $\pm$ 1.1 <sup>b</sup>	4.5 $\pm$ 2.1 <sup>b</sup>
n-3/n-6	2.5 $\pm$ 0.3	2.0 $\pm$ 0.5	1.8 $\pm$ 0.2	1.5 $\pm$ 0.1	1.2 $\pm$ 0.0	0.9 $\pm$ 0.3

Abbreviations as in Table 4.