

The role of phospholipids in nutrition and metabolism of teleost fish

Douglas R. Tocher^{1*}, Eldar Å. Bendiksen², Patrick J. Campbell² and J. Gordon

Bell¹

¹Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK.

²BioMar AS, Nordregt. 11, N-7484 Trondheim, Norway.

Keywords: Teleosts; fish; phospholipids; requirements; nutrition; metabolism; dietary supply; plant meals; cholesterol

^{*}, Correspondence to: Douglas Tocher, Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, United Kingdom. Tel. No: +44 1786 467996; Fax No: +44 1786 472133; Email:

d.r.tocher@stir.ac.uk

Abstract

It has been known for almost 25 years now that inclusion of intact phospholipids in the diet could improve culture performance of various freshwater and marine fish species. The primary beneficial effect was improved growth in both larvae and early juveniles, but also increased survival rates and decreased incidence of malformation in larvae, and perhaps increased stress resistance. Determination of absolute dietary requirements has been hampered by the use, in different dietary trials, of a wide range of phospholipid preparations that can vary greatly both in phospholipid content and class composition. Larval studies have been compromised further by the need on many occasions to supply phospholipid through enrichment of live feeds with subsequent re-modelling of the phospholipid and fatty acid compositions. Generally, the levels of phospholipid requirement are around 2 - 4% of diet for juvenile fish and probably higher in larval fish. The effects were restricted to young fish, as a requirement for dietary phospholipids has not been established for adult fish, although this has been virtually unstudied. As the majority of studies have used crude mixed phospholipid preparations, particularly soybean lecithin, but also other plant phospholipids and egg yolk lecithin, that are enriched in several phospholipids, it has been difficult to elucidate which specific phospholipid classes impart beneficial effects. Based on the few studies where single pure phospholipid species have been used, the rank order for efficacy appears to be phosphatidylcholine > phosphatidylinositol > phosphatidylethanolamine > phosphatidylserine. The efficacy of other phospholipid classes or sphingolipids is not known. The mechanism underpinning the role of the phospholipids in larval and early juvenile fish must also explain their lack of effect in adult fish. The role of phospholipids appears to be independent of fatty acid requirements although the presence of an unsaturated fatty acid at the *sn*-2 position may be important. Similarly, the phospholipid requirement is not related to the delivery of other essential dietary components such as the bases choline and inositol. Studies also suggested that the phospholipid effect was not due to generally enhanced emulsification and digestion of lipids. Rather the evidence led to the hypothesis that early developing stages of fish had impaired ability to transport dietary lipids away from the intestine possibly through limitations in lipoprotein synthesis. The current hypothesis is that the enzymic location of the limitation is actually in phospholipid biosynthesis, perhaps the production of the glycerophosphobase backbone and that dietary supplementation with intact phospholipids in larvae and juvenile fish compensated for this. Thus, dietary phospholipids increase the efficiency of transport of dietary fatty acids and lipids from the gut to the rest of the body possibly through enhanced lipoprotein synthesis.

59	Contents
60	
61	1. Introduction
62	2. Roles of phospholipids
63	2.1 Structural
64	2.2 Regulation of metabolism and physiology
65	2.3 Eicosanoids
66	2.4 Phosphoinositides
67	2.5 Platelet-activating factor
68	2.6 Energy provision
69	3. Phospholipid biosynthesis
70	3.1 De novo synthesis
71	3.2 Turnover and remodelling
72	4. Phospholipid digestion, absorption and transport
73	4.1 Digestion
74	4.2 Absorption
75	4.3 Transport
76	5. Phospholipid requirements in fish
77	5.1 Early studies
78	5.2 Limitations in studies
79	5.3 Requirement criteria
80	5.4 Range of species and developmental stages
81	5.5 Quantitative requirements
82	5.6 Qualitative requirements
83	6. Mechanisms of requirement for intact phospholipids in larvae and juveniles
84	6.1 Improved diet qualities
85	6.2 Improved digestion
86	6.3 Provision of essential components
87	6.4 Fatty acids (energy and essential fatty acids)
88	6.5 Phosphorous
89	6.6 Choline and inositol
90	6.7 Provision of intact phospholipid
91	7. Dietary phospholipids and sustainability

92	7.1 Sustainability
93	7.2 Dietary sources of phospholipids
94	7.2.1 Oils
95	7.2.2 Meals
96	7.3 Analysis of phospholipid content and composition in fish diets and ingredients
97	7.4 Dietary cholesterol
98	8. Conclusions
99	References
100	
101	

1. Introduction

Phospholipid is a general term that includes all lipids containing phosphorus. However, it is a term often mistakenly equated with phosphoglycerides, the most common of the phospholipids. Phosphoglycerides are characterised by a common backbone of phosphatidic acid (PA), formed from L-glycerol 3-phosphate with two fatty acids esterified on positions 1 and 2. The major phosphoglycerides of animal including fish tissues phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylserine (PS) and phosphatidylinositol (PI), respectively, are formed by the esterification of the “bases” choline, ethanolamine, serine and inositol to the phosphate group of PA. A common generalisation has been that saturated and monounsaturated fatty acids are preferentially esterified on position *sn*-1 of phosphoglycerides with polyunsaturated fatty acids (PUFA) preferentially esterified on position *sn*-2. However, there are many exceptions to this, such as the di-docosahexaenoyl phosphoglycerides that are abundant in the retina of fish, specifically in rod outer membrane segments (Bell and Tocher, 1989; Bell and Dick, 1991). Another phosphorus-containing polar lipid is the sphingolipid, sphingomyelin, a complex lipid based on the long-chain amino alcohol sphingosine. All sphingolipids contain a long chain, generally saturated or monounsaturated fatty acid, linked to the amino group of sphingosine (forming a ceramide), with different polar groups, such as, phosphocholine in the case of sphingomyelin, attached to the primary alcohol group.

For the remainder of this review we will exclusively use the term “phospholipid” when, in actual fact, phosphoglyceride would be the more correct term, as described above. However, the use of phospholipid will be consistent with the vast majority of studies that have been reported in the literature. Similarly, the abbreviations PC, PE, PS, PI and PA, as used above, are not strictly correct but are by far the most commonly used for the major phosphoglyceride classes and so will be used throughout this review.

2. Roles of phospholipids

2.1 Structural

The amphipathic structure of phospholipids, having both hydrophilic (*sn*-3 phosphate/head-group) and hydrophobic (*sn*-1 and -2 fatty acids) regions means they have the same central role in the structure of cell membrane bilayers in fish as they do in mammals. Although not well studied in fish, there is evidence that the asymmetric distribution of phospholipids in cell membranes

established in mammals, with choline-containing phospholipids, PC and sphingomyelin concentrated in the outer leaflet and PE, PS and, to a lesser extent, PI concentrated in the inner leaflet, is also a feature of fish cell membranes (Kagan et al., 1984). The dynamic changes in the composition and metabolism of the phospholipids in biomembranes in response to environmental factors, especially temperature, have been reviewed previously (Hazel and Williams, 1990; Hochachka and Mommsen, 1995). Their amphipathicity is also central to the role phospholipids have in the structure of the lipoproteins that are important in the extracellular transport of lipids in the blood and lymph. The lipoproteins are described in more detail in section 4.3 but, essentially, phospholipids enable hydrophobic lipids such as triacylglycerols and steryl esters to be transported in aqueous environments by forming, along with cholesterol and proteins, the lipid/water interfaces (Tocher, 1985). It can be argued that phospholipids also have an important structural role in the digestion of lipids as they are essential in forming intra-luminal mixed micelles along with bile salts and dietary lipids (see section 4.2) (Olsen and Ringø, 1997). This phospholipid is not entirely of dietary origin as fish bile can contain variable amounts of phospholipid (Moschetta et al., 2005). It is thought that biliary phospholipid has two roles in the bile as the formation of mixed micelles with bile salts not only solubilises biliary cholesterol, but also has a cytoprotective effect, protecting biliary tract epithelium from the cytotoxic effects of bile salts (Moschetta et al., 2005). PC is usually the major phospholipid in fish bile but sphingomyelin can also be a major component, particularly in species with lower biliary phospholipid levels and phospholipid:bile salt ratios (Moschetta et al., 2005).

2.2 Regulation of metabolism and physiology

Phospholipids are important precursors for a range of highly biologically active mediators of metabolism and physiology including eicosanoids, diacylglycerol (DAG), inositol phosphates and platelet activating factors (PAFs). With the possible exception of eicosanoids, these areas of phospholipid metabolism are relatively poorly studied in fish. However, there is evidence that all these pathways occur in fish and that the phospholipid-derived mediators play similar roles in fish as they do in mammals (Tocher, 1995, 2003).

2.2.1 Eicosanoids

Phospholipids are the source of the substrate fatty acids for the formation of eicosanoids, a range of highly bioactive derivatives of, in particular C₂₀ highly unsaturated fatty acids (HUFA),

especially arachidonic acid (ARA, 20:4n-6) and eicosapentaenoic acid (EPA, 20:5n-3). Fatty acids released from membrane phospholipids by the action of phospholipase A₂ are converted by either cyclooxygenase enzymes, which produces cyclic oxygenated derivatives, collectively called prostanoids, including prostaglandins, prostacyclins and thromboxanes, or lipoxygenase enzymes which produce linear oxygenated derivatives including hydroperoxy- and hydroxy fatty acids, leukotrienes and lipoxins. Eicosanoids are produced by many tissues in response to various extracellular stimuli and are autocrines, hormone-like compounds of short half-life that act in the immediate vicinity of their production. They are implicated in many physiological processes including blood clotting and cardiovascular tone, immune and inflammatory responses, reproduction, renal and neural functions. The distribution and production of eicosanoids in fish species and tissues and their possible roles have been reviewed previously (Sargent et al., 1989, 2002; Tocher, 1995, 2003).

2.2.2 Phosphoinositides

Phosphorylated derivatives of PI (e.g. PIP₂) are produced by the action of various kinases and phosphatases on the inositol ring. The phosphorylated derivatives of PI may themselves have important roles in metabolism including golgi/lysosome/endosome trafficking, cytoskeleton regulation and cell survival. The most studied pathway though is the cleavage of PIP₂ by phospholipase C- β to produce two intracellular second messengers, DAG, and the inositol phosphate, IP₃, in response to various hormones and effectors (Berridge, 1987). IP₃ induces calcium mobilisation from the endoplasmic reticulum, and increased intracellular Ca²⁺ and DAG are activators of protein kinase C, a threonine/serine kinase and important regulator of metabolism (Kuo et al., 1980). DAG can also be phosphorylated by DAG kinase to form PA and lyso-PA, increasingly being recognised as other important intracellular regulators of a growing number of signalling proteins (Hao et al., 2007). Other phospholipids, including PC and PS, also have important metabolic functions in mammals as specific activators of protein kinase C, along with DAG and Ca²⁺ ions (Kuo et al., 1980). Protein kinase C and PS-activation has been reported in rainbow trout (*Oncorhynchus mykiss*) spleen and dogfish (*Scyliorhinus canicula*) rectal gland (Bell and Sargent, 1987), and a role for protein kinase C has been implicated in the stimulation of steroidogenesis in goldfish (*Carassius auratus*) gonadal tissues (Van Der Kraak, 1990; Wade and Van Der Kraak, 1991). Inositol phospholipid metabolism including the roles of phospholipase C

and Li-sensitive phosphatases have been studied in metabolically active electrocytes from the electric ray (*Discopyge tschudii*) (Arias and Barrantes, 1990).

2.2.3 Platelet-activating factor (PAF)

PAF, 1-*O*-alkyl-2-acetyl-*sn*-glycero-3-phosphocholine, is a biologically active phospholipid synthesised by inflammatory cells and a potent mediator of many leukocyte functions, including platelet aggregation, inflammation, and anaphylaxis (Snyder, 1987). PAF is biosynthesized from lyso-PAF and acetyl CoA by the enzyme lyso-PAF acetyltransferase and degraded (thereby terminating activity) by a group of enzymes called PAF acetylhydrolases that are related to phospholipase A₂. In mammals, the reacylation of lyso-PAF is highly specific for ARA, and it is the 1-alkyl-2-arachidonyl-glycerophosphocholine species that is the substrate for the synthesis of PAF *via* an ARA-specific phospholipase A₂ that also produces ARA for eicosanoid synthesis (Snyder, 1987). Synthesis of PAF has been demonstrated in rainbow trout (Turner and Lumb, 1989) and PAF acetylhydrolase activity has been reported in fish serum (Cabot et al., 1984).

2.3 Energy production

Any lipid class containing fatty acids can act as a source of energy, which is released through β -oxidation of the acyl chains producing acetyl-CoA and NADH that are further metabolised via the tricarboxylic acid cycle and oxidative phosphorylation, respectively. Triacylglycerols are the primary class for lipid storage and energy provision, but phospholipids can serve as a source of energy in fish in certain circumstances, such as embryonic and early larval development.

Upon hatching, the larvae of many fish species have immature mouthparts and intestinal tract and are unable to commence exogenous feeding immediately. Therefore, throughout embryogenesis in the egg and then during early larval development up to first feeding, the larvae gain nutrition from endogenous energy reserves in the yolk. Essentially there are two main types of fish egg, those with oil globule(s) having relatively high levels of neutral lipids (20-50% of egg total lipid), and those without an oil globule, low levels of neutral lipids (< 15% of total lipid) and, consequently, high levels of phospholipids, predominantly PC (Wiegand, 1996; Salze et al., 2006). In eggs of the latter type, including those from Atlantic herring (*Clupea harengus*), cod (*Gadus morhua*), halibut (*Hippoglossus hippoglossus*) and plaice (*Pleuronectes platessa*), phospholipid, predominantly PC, is utilised during both embryogenesis and early larval development (Tocher et al., 1985a; Fraser et al., 1988; Rainuzzo et al., 1992; Finn et al., 1995;

Rønnestad et al., 1995 (Finn et al., 1995; Rønnestad et al., 1995). In contrast, neutral lipid-rich eggs such as those from turbot (*Psetta maximus*), utilise primarily triacylglycerols, but also sterol and wax esters where present (Sargent et al., 1989, 2002; Rainuzzo et al., 1992; Weigand, 1996).

Another relatively common feature observed during development is the conservation and/or synthesis of PE, as reported in both the phospholipid-rich eggs of cod (Fraser et al., 1988), plaice and halibut (Rainuzzo et al., 1992; Rønnestad et al., 1995), and also the neutral lipid-rich eggs of Atlantic salmon (*Salmo salar*) and turbot (Cowey et al., 1985; Rainuzzo et al., 1992), Senegal sole (*Solea senegalensis*) (Mourete and Vazquez, 1996) and dentex (*Dentex dentex*) (Mourete et al., 1999). This results in a decrease and normalisation of the PC : PE ratio as development proceeds, from the high values seen in most marine fish eggs to values normally observed in fish tissues. This is particularly the case in the phospholipid - rich eggs, dominated by PC, where PC is catabolised during embryogenesis, but has also been observed in Atlantic salmon where PC, originally accounting for over 94% of the phospholipid in the egg, was continuously metabolized so that in the fry a PC : PE ratio approaching that of salmon muscle was obtained (Cowey et al., 1985).

These data show a decrease in the absolute amount of phospholipid, usually PC, during embryogenesis and early larval development, particularly in phospholipid-rich marine fish eggs. A possible consequence of complete catabolism of phospholipid for energy would be the loss of important PUFA. However, in an early study in Atlantic herring, much of the PUFA liberated by the catabolism of PC was selectively retained in the neutral lipid pool (Tocher et al., 1985b). About one third of the docosahexaenoic acid (DHA, 22:6n-3) released during PC catabolism in cod eggs was incorporated into neutral lipids (Fraser et al., 1988). Similarly, in halibut, over 60% of the DHA from PC hydrolysis was selectively retained in PE and neutral lipids (Rønnestad et al., 1995). However, DHA was also a quantitatively important fuel in halibut eggs with almost 40% of the DHA being catabolised (Rønnestad et al., 1995). Similarly, in cod the fatty acids in PC were catabolised non-selectively with both EPA and DHA being utilised (Finn et al., 1995).

3. Phospholipid biosynthesis

3.1 De novo synthesis

In mammals, phospholipids can be synthesised de novo by two main mechanisms (see Lykidis, 2007). One utilizes a cytidine diphosphate (CDP)-activated polar head group for attachment to 1,2-DAG, and the other utilizes CDP-activated DAG and an inactivated polar head group (Fig.1).

They both initially proceed by esterification of glycerol-3-phosphate with two activated fatty acids (acyl-CoA) to form PA. Action of PA phosphatase on PA results in the production of DAG, which can then react with CDP-choline or CDP-ethanolamine to produce PC and PE, respectively. Alternatively, CTP reacts with PA to form CDP-DAG, which can then combine with serine and inositol to form PS and PI, respectively. Reaction of CDP-DAG with glycerol-3-phosphate leads to phosphatidylglycerol (PG) and combination of PG with a further CDP-DAG leads to the formation of diphosphatidylglycerols that include cardiolipin. PS can also be formed in the membrane bilayer by exchange reactions of serine for ethanolamine or choline in PE and PC, respectively. PS can be converted back to PE by decarboxylation and PC can also be formed by three successive methylations of PE utilizing S-adenosylmethionine.

The pathways of *de novo* phospholipid biosynthesis have not been extensively studied in fish, but all the necessary phospholipid biosynthetic enzymes and protein families can be found in the fugu (*Takifugu rubripes*) and zebrafish (*Danio rerio*) genomes (Lykidis, 2007). The existing biochemical evidence also suggests that pathways are essentially the same as in mammals (Tocher, 1995). Incubation of trout liver microsomes with ^{14}C -glycerol-3-phosphate in the presence of palmitoyl-CoA, resulted in three-quarters of the radioactivity being recovered in total phospholipids, predominantly PA and lyso-PA but also other phospholipid classes, demonstrating glycerol-3-phosphate acyltransferase activity and supporting the conclusion that phospholipid biosynthesis in fish proceeded *via* a PA intermediate (Holub et al., 1975a). Activity of α -glycerophosphate acyltransferase in fish was later confirmed, although it was reported to be extremely low compared to other animals (Iritani et al., 1984). Studies on the incorporation of ^{14}C -palmitate and ^{14}C -glycerol-3-phosphate into lipids in carp (*Cyprinus carpio*) intestinal homogenates in the presence of CTP, CDP-choline and CDP-ethanolamine showed phospholipid biosynthesis from PA and DAG intermediates, indicating synthesis from moieties other than lyso-phospholipids (Iijima et al., 1983). However, little is known about what determines the fate of newly-synthesised DAG (PC or triacylglycerol synthesis) in fish, but it has been suggested that fatty acid composition of the DAG may differentially affect the activities of the biosynthetic enzymes (Oxley et al., 2007). Early studies demonstrated the presence of CDP-choline:DAG choline phosphotransferase activity in trout liver microsomes (Holub et al., 1975b), and brain and liver from goldfish (Leslie and Buckley, 1975), establishing that the choline phosphotransferase (CPT) pathway for the biosynthesis of PC operated in fish. Similarly, synthesis of PE from DAG and the presence of CDP-ethanolamine phosphotransferase (i.e. the EPT pathway) were subsequently demonstrated in trout hepatocytes (Hazel, 1990). The activity of CTP in

microsomes from intestinal mucosa in adult Atlantic salmon (~ 1 kg) was comparable to the reported mammalian activity (Oxley et al., 2005). Thus, synthesis via DAG and CPT/EPT was confirmed as the predominant pathway for phospholipid synthesis in fish as in mammals (Hazel, 1990). Although not directly studied or demonstrated in fish, the presence in fish of PI, PG and cardiolipin, that cannot be synthesised via the phosphotransferase pathways, implies the presence of the alternative CDP-DAG pathway in fish. Similarly, not all the phospholipid inter-conversion pathways have been studied in fish, but the activities of PE-methyltransferase (PE to PC) and PS-decarboxylase (PS to PE) have been demonstrated in trout hepatocytes (Hazel, 1990).

3.2 Phospholipid turnover and remodelling

Remodelling and turnover of phospholipids can involve both head groups and fatty acyl chains. The mechanisms whereby phospholipid classes (PC, PE, PS and PI, etc.) could be inter-converted are described above (see Fig.1), and phospholipid remodelling at this level is certainly involved in homeoviscous adaptation of biological membranes, especially in relation to temperature adaptation. Thus, cold acclimation in fish has been associated with increased proportions of PE and decreased proportions of PC (Hazel and Williams, 1990). Although the activities of CDP-choline and CDP-ethanolamine phosphotransferase were both reduced at lower temperature, synthesis of PC was more dependent upon temperature than that of PE and, in addition, PS decarboxylase activity was increased (Hazel, 1990). As a result, the PC : PE synthesis ratio was positively correlated with temperature and so these mechanisms may underpin the increased PE : PC ratio in cold-acclimated fish (Hazel, 1990).

Fatty acid retailoring of phospholipids through acyl exchange reactions has been relatively more studied in fish, again most often in relation to homeoviscous adaptation to environmental changes. The enzymes required for deacylation of phospholipids and reacylation of lyso-phospholipids, and thus for the turnover of phospholipids, have been demonstrated in fish. Intracellular phospholipase A activities have been reported in muscle tissue from a variety of fish species including rainbow trout and cod (see Tocher, 2003), and a review of muscle lipase activities indicated that hydrolysis of phospholipids in fish was primarily under the control of phospholipases A₁ and A₂ (Shewfelt, 1981). The activity of liver microsomal phospholipase A₂ has been studied in trout (Neas and Hazel, 1985), whereas cytosolic phospholipase A₂ was demonstrated to be the principal phospholipase activity during zebrafish embryogenesis (Farber et al., 1999). AcylCoA:1-acyl-*sn*-glycero-3-phosphorylcholine acyltransferase activity has also

been demonstrated in trout liver microsomes (Holub et al., 1976). It is presumed that the fatty acyl and head group specificities of the enzymes involved in the deacylation/reacylation turnover processes have important roles in maintaining the characteristic fatty acyl distribution among the phospholipid classes (Tocher, 2003). Thus, PC is characterized by having high 16:0 (higher than any other class) and relatively lower PUFA (i.e. lower than PE and PS); PE generally displays intermediate levels of saturated and monounsaturated fatty acids and high levels of PUFA which are relatively evenly split between C₂₀ and C₂₂ PUFA; PS is characterized by high 18:0 and high PUFA, predominantly C₂₂ PUFA; PI also has high 18:0, but relatively lower PUFA, which is predominantly C₂₀, particularly ARA, and a low n-3 to n-6 ratio (Tocher, 1995).

4. Phospholipid digestion, absorption and transport

4.1 Digestion

Digestion of dietary phospholipids is relatively unstudied in fish, but it is presumed that the mechanisms are generally similar to those in mammals whereby phospholipids are digested by intestinal phospholipase A₂, secreted by the pancreas, resulting in the formation of 1-acyl lyso-phospholipids and free fatty acids that are absorbed by the intestinal mucosal cells (Henderson and Tocher, 1987; Sargent et al., 1989). Lyso-phospholipids have a detergent action and so aid the digestion of other lipids and, indeed, some phospholipid is secreted in the bile as substrate for phospholipase A₂. The lack of a discrete pancreas in most teleost species has hampered studies on intestinal lipolysis in fish in general and, although several studies have demonstrated phospholipase A₂ activities in intestinal tissue of fish, it has been difficult to establish conclusively if they were intracellular activities or secreted (see Tocher, 2003). However, two phospholipase A₂ isoforms were purified from red sea bream hepatopancreas and characterisation confirmed them as low molecular weight, Ca²⁺-dependent group I (secretory) forms (Ono and Iijima, 1998). Phospholipase A₂ activity was only detected several days after mouth opening in marine fish larvae (Zambonino-Infante and Cahu, 2001). There are few data on rates of phospholipid hydrolysis, but digesta from turbot rectum was shown to hydrolyse PC, although at slower rates than triacylglycerol and sterol esters (Koven et al., 1994).

4.2 Absorption

The mechanisms of absorption of the products of phospholipid digestion have not been extensively studied in fish but are assumed to be generally similar to that in mammals. Thus, the hydrolytic products, 1-acyl lyso-phospholipids and free fatty acids, will be associated with all the other products of fat digestion in mixed micelles with bile salts, that diffuse to the intestinal mucosa where uptake into the enterocytes occurs, probably mainly by passive diffusion.

The concentration of lyso-phospholipids is very low in fish plasma and so it is assumed that in the intestinal mucosa the majority of the 1-acyl lyso-phospholipids are re-esterified with activated free fatty acids in the microsomes before export from the enterocytes into the circulatory system (Sargent et al., 1989). This is supported by a study measuring the absorption of ^{14}C -dioleoyl PC force-fed to carp (Iijima et al., 1990). Radioactivity in plasma lipoproteins 20 - 28 h after force-feeding was associated with various lipid classes including PC, and the radioactivity associated with the *sn*-1 position was more than twice that associated with the *sn*-2 position (Iijima et al., 1990). There appears to be some specificity in the reacylation processes as a study of fatty acid uptake by isolated enterocytes from rainbow trout showed that the recovery of HUFA and 16:0 in phospholipids was higher than that of other fatty acids (Perez et al., 1999).

4.3 Transport

Phospholipids are transported in the blood of most fish as lipoproteins as they are in mammals (see Tocher, 1995). Lipoproteins include chylomicrons, very-low density lipoproteins (VLDL), low density lipoprotein (LDL) and high density lipoprotein (HDL) containing 8, 21, 25 and 29 % of total weight as phospholipids, respectively (Chapman et al., 1978). However, as a percentage of the total lipids, the proportion of phospholipids in trout lipoproteins ranges from around 9% in chylomicrons to 23 % in VLDL, 35 % in LDL and 53 % in HDL (Chapman et al., 1978), although carp lipoproteins contained proportionally more phospholipid than trout, salmon or sardine (*Sardinops*) lipoproteins with the total lipid from VLDL, LDL and HDL containing 30 %, 58 % and 82 % phospholipid, respectively (Iijima et al., 1989). The relative proportions of the plasma lipoproteins in fish are dependent upon age, nutrition and reproductive cycle and also vary with species, but is a constant characteristic of each species depending upon dietary status (Fremont and Leger, 1981).

The re-esterification reactions in the intestine occur primarily in the endoplasmic reticulum leading to the production of chylomicrons and, to a lesser extent, VLDL (see Tocher, 1995). Transport from the intestine to the liver is primarily via the lymphatic system as in trout and

tench (*Tinca tinca*) although, in carp, intestinal lipoproteins may be transported directly to the liver via the portal system (Noaillac-Depeyre and Gas, 1974; Sire et al., 1981). Although the effects of phospholipid content and composition on the production of intestinal lipoproteins in fish have not been studied, as chylomicrons and VLDL have very different phospholipid contents, variable proportions of dietary phospholipid could be accommodated by varying the relative proportions of the intestinal lipoproteins produced. In common with most vertebrates, PC appears to be the predominant phospholipid class in fish lipoproteins, although the precise phospholipid class composition in fish plasma lipoproteins has been rarely reported (Nelson and Shore, 1974; Chapman, 1980).

The processes of lipoprotein metabolism and remodelling are essentially the same in fish as in mammals and are described in detail previously (Tocher, 1995, 2003) and so are only summarised here. Although some may be produced in the intestine, the majority of VLDL is synthesized in the liver. Triacylglycerols in chylomicrons and VLDL are hydrolyzed by lipoprotein lipase (LPL) and hepatic lipase (HL) at tissue sites with the hydrolysis products being absorbed. Excess surface constituents, including phospholipids, dissociate as nascent HDL particles, leading to the formation of intermediate density lipoprotein (IDL) and LDL. Nascent HDL can take up free cholesterol from peripheral tissues which is then esterified by the action of lecithin:cholesterol acyl transferase (LCAT), a plasma enzyme which catalyzes the esterification of cholesterol using fatty acid from PC, resulting in the production of mature HDL. All the major enzymes of lipoprotein metabolism and remodelling, including LPL, HL and LCAT, have been demonstrated in the plasma of various fish (Tocher, 1995).

Although virtually unstudied in fish, by analogy with the system characterized in mammals, phospholipids can probably be taken up into the tissues by two or three main mechanisms. In liver and some other tissues, receptor-mediated endocytosis, *via* receptors for B/E and E apoproteins, are important pathways for LDL (apo B), VLDL- and chylomicron-remnants (apo B and E) and a subfraction of HDL rich in apo E. Non-specific pinocytosis may be an important pathway for all lipoproteins, but particularly LDL and HDL. In liver, around 30 % of LDL uptake is *via* a non-receptor-mediated pathway in mammals. Finally, phospholipids and other surface components of VLDL and chylomicrons may be taken up or exchanged *via* direct interaction with the endothelial cell membranes in the tissues.

5. Phospholipid requirements of fish

All the available evidence suggests that phospholipid metabolic pathways, including those of *de novo* phospholipid biosynthesis, are essentially the same in fish as in mammals. However, there is now substantial evidence that at least some species of fish, both freshwater and marine, may have only a limited capacity to synthesise phospholipids *de novo* both at larval and early juvenile stages (see Coutteau et al., 1997). This may reflect the fact that many fish larvae receive an abundance of phospholipids in their natural diets, whether from yolk sac lipids prior to first feeding or from natural prey at and after first feeding. The same situation may well hold for *de novo* biosyntheses of cholesterol and sphingolipids, which have been scarcely studied in fish.

5.1 Early studies

During studies in the early 1980s investigating microparticulate diets as replacements for live feeds for rearing larval marine fish, Kanazawa and coworkers observed that dietary phospholipids were essential for normal growth and survival of fish larvae such as ayu (*Plecoglossus altivelis*) and red sea bream (*Pagrus major*) (Kanazawa, 1985). Improved growth and survival rates were obtained with 10-100 day-old larval ayu when the diets were supplemented with 3% chicken-egg lecithin, bonito-egg lecithin fraction (PC) or soybean lecithin (Kanazawa et al., 1981, 1983a). Bonito-egg cephalin fraction (PE) was not as effective as the PC fraction (Kanazawa et al., 1981, 1983a). Similar results were found with larval red sea bream (Kanazawa et al., 1983b). These workers also reported that the addition of some phospholipid to the microparticulate diet formulation reduced the incidence of malformations (Kanazawa et al., 1981, 1983a,b). These initial reports stimulated a significant amount of further research into phospholipid requirements of larvae and juveniles of a number of other fish species (see Coutteau et al., 1997). The following sections summarise these studies.

5.2 Limitations in studies

Studies investigating phospholipid requirements in fish have some limitations based on the requirement for a deficient basal diet. For juvenile fish, deficient diets can be formulated by careful choice of the raw ingredients. However pelleted diets are not suitable for marine fish larvae, and the traditional diets, live feeds, are unsuitable as they contain phospholipids and further enrichment can be difficult (Nordgreen et al., 2007). Therefore, microdiets are required, but they have been difficult to produce and were not available for all the studies (Baskerville-Bridges and Kling, 2000; Yufera et al., 2000; Cahu and Zambonino Infante, 2001; Koven et al.,

2001; Robin and Vincent, 2003; Kvale et al., 2006). Irrespective of whether pelleted feeds or microdiets are used, they can both have problems of low palatability as the formulations must be either fish meal-free or use defatted fish meal. As a result the majority of studies have been performed with casein-based diets and the resultant impact on attractant and palatability factors can lead to acceptance problems.

Supplementation of phospholipids to basal deficient diets can pose problems as, although commercial phospholipid preparations (lecithins) are available, they are not pure products. Lecithin is, strictly speaking, the trivial name for PC, but has been used commercially to describe a range of phospholipid-enriched products usually obtained either from chicken eggs or as a by-product of vegetable oil processing (see section 7.2.1). The commercially available lecithins can vary greatly in purity, and have highly variable phospholipid contents and class compositions, as well as very different fatty acid compositions. An alternative is to use pure phospholipid species (classes), but these have limited availability and are costly, although it is possible to purify individual phospholipid classes in the laboratory (Geurden et al., 1998a). Irrespective of the specific phospholipid supplemented, replacement of one lipid class with another, i.e. triacylglycerol with phospholipid, will invariably alter the fatty acid composition of the diets and this can be difficult to fully control. This is also the case in comparing different phospholipid preparations. Marine phospholipids were superior to soybean phospholipids in promoting growth in turbot larvae, but fatty acid compositions were not controlled and so the marine phospholipid diets also had far superior fatty acid profiles in terms of n-3 HUFA content (MacQueen Leifson et al., 2003).

Finally, another major limitation over all the studies is that older fish have not been investigated. The rationale presumably being that a requirement for intact phospholipids would only manifest itself in fish receiving diets having very low levels of intact phospholipids. Once being fed extruded diets based on fish meal, the diets will contain a luxury of intact phospholipids of all classes and thus a requirement, if any, would be satisfied.

5.3 Requirement criteria

The criteria used to measure requirements for phospholipids have generally been gross performance indicators including growth, survival and development. Thus, in most studies, the requirement for intact phospholipid has been increased growth, improved survival and reduced malformations or deformities (Coutteau et al., 1997). Effects of nutritional components, including dietary phospholipids, on skeletal development in fish larvae have been reviewed recently (Cahu

et al., 2003a). A limited number of studies have also looked at stress resistance and showed that this could be increased by phospholipid supplementation (Takeuchi et al., 1992; Kanazawa, 1993; Weirich and Reigh, 2001). Few other physiological, metabolic, biochemical or molecular parameters have been investigated (Fontagne et al., 2000; Koven, 2003; Zambonino-Infante and Cahu, 2007; Hamza et al., 2008).

5.4 Range of species and developmental stages

A relatively defined quantitative requirement for dietary phospholipid has been determined in a number of fish including the marine species red sea bream, Japanese flounder (*Paralichthys olivaceus*), rock bream/knife jaw (*Oplegnathus fasciatus*), European sea bass (*Dicentrarchus labrax*), striped jack (*Pseudocaranx dentex*), gilthead sea bream (*Sparus aurata*) and turbot, and the freshwater species ayu, common carp, rainbow trout and white sturgeon (*Acipenser transmontanus*) (see Table 1 for references). A semi-quantitative requirement has also been determined in Atlantic salmon at the freshwater stage. The requirement for phospholipid has been demonstrated at both larval and juvenile stages for ayu, Japanese flounder, rock bream/knife jaw and European sea bass, in larvae for carp, red and gilthead sea bream, and in juveniles for striped jack, turbot, rainbow trout, Atlantic salmon and white sturgeon. Based on studies comparing a neutral lipid-rich commercial diet and a phospholipid-rich zooplankton diet, Olsen et al. (1991) suggested that polar lipids may promote growth in juvenile Atlantic cod, but further work is required to establish if there is a genuine requirement in this species. As mentioned previously, few studies have been conducted but a requirement for phospholipid has not been conclusively demonstrated in adults of any fish species (Olsen et al., 1999).

5.5 Quantitative requirements

Quantitative requirements have generally been reported in terms of phospholipid levels as a percentage of the diet by weight. However, some data should be regarded as semi-quantitative only, as the actual phospholipid content of the phospholipid preparations used in different studies, which varies as described above, is not always taken into account and so the absolute phospholipid level required may be lower than the value quoted in some cases. This also means that it is not always appropriate to directly compare values in different trials using different phospholipid preparations. Differences in the initial size of larvae or juveniles used, and the durations of the feeding trials also contribute to the semi-quantitative nature of the data. Coutteau et al. (1997) summarised the differences between the trials in all studies published up until the

date of that review. Acknowledging these caveats, values quoted for the quantitative requirements of phospholipids for larval fish range from about 2 - 12 % of diet (Table 1). Requirements were lower in the freshwater fish, carp (2 %) and ayu (3-5 %), and higher in the marine fish red sea bream (5 %), rock bream/knife jaw (5-7 %) and Japanese flounder (7%), with the highest reported value being for European sea bass at 12 % of diet. In juveniles, the values ranged from around 1.5 - 7 % of diet, including 1.5 % for striped jack, 2-3 % for European sea bass and around 4-6 % for Atlantic salmon, and 7 % for Japanese flounder. In Atlantic salmon, studies showed that the phospholipid requirement was 4 - 6 % of diet in fish of initial size 180 mg, 4 % in fish of 1-1.7 g initial weight, whereas in fish of 7.5 g initial weight, no requirement was observed. Similarly, in white sturgeon of 5-10 g, no requirement for dietary phospholipid was detected. Therefore, the data appear to suggest a possible trend with quantitative requirements decreasing with development, at least from larvae to small juveniles. No requirement has been observed in fish (at least salmon and white sturgeon) of greater than 5 g initial weight although this may be due to the short-term nature of the studies. Thus, in Atlantic salmon cultured over a 24-month growth cycle low dietary phospholipid might induce growth depression and other effects over this longer period of culture.

5.6 Qualitative requirements

There are few data available with which to compare the efficacy of different phospholipid classes in satisfying phospholipid requirements. First of all, few phospholipid preparations have been investigated, principally soybean lecithin (SL), egg lecithin (EL), but occasionally also some marine phospholipids (Table 1). As mentioned above, all of these preparations are mixtures of different phospholipid classes, but mainly PC, PE, PI and PS although the proportions vary. For instance, SL can contain 50 - 86 % phospholipid, with typically around 20 - 25 % each of PC, PE and PI. Therefore, it is not possible to draw any conclusions on the efficacy of individual phospholipid classes in trials using commercial crude phospholipid preparations. However, some studies have utilized semi-purified phospholipids, including soybean PC (80 - 98 % PC) or egg PC (95 % PC), soybean PE (99 % PE) and soybean PI (Geurden et al., 1998a). Thus, in the few studies in which comparisons are possible, both PC and PI were effective in meeting phospholipid requirements in larval ayu whereas PE was less effective (Kanazawa, 1983; Kanazawa et al., 1985). In Japanese flounder larvae, growth improved with PC, but not with PI or PE (Kanazawa, 1993). When carp larvae were fed diets containing either purified phospholipid classes (PC, PE, PS and PI), PC gave higher initial growth but induced deformities, whereas PI

gave optimal survival and minimal deformities (Geurden et al., 1998a). PE and PS gave intermediate results, and hydrogenated PC or lyso-PC was not as effective as native PC. Therefore, there are no clear patterns other than PC and PI may generally be more effective than PE and PS, with PC perhaps being more important for growth and PI being more important for normal development and prevention of malformations, at least in carp larvae. However, it is possible that different phospholipid classes may play slightly different roles and thus have different effects in different species as PI may have been more effective than PC in improving growth in larval ayu (Kanazawa, 1983).

6. Mechanisms of requirement for intact phospholipids in larvae and juveniles

6.1 Improved diet qualities

It has been suggested that phospholipids may help to reduced leaching of water-soluble micronutrients (minerals and vitamins) from semi-purified diets (see Coutteau et al., 1997). Addition of phospholipid influenced the loss of dry matter in casein-based microdiets for post-larval shrimp (Camara, 1994). In a study investigating phospholipid requirements for European sea bass and turbot using extruded diets with or without 2 % EL (69% pure), a diet water stability test showed no effect of phospholipid on dietary fatty acid contents but micronutrients were not investigated (Geurden et al., 1997b). This possible effect of dietary phospholipid requires further study.

Although phospholipids contain oxidatively sensitive PUFAs, there is some evidence that they can express an antioxidative effect on various oils and fats (Ishihara, 1997). Satoh and Ishihara (1997) tested various compounds, representative of the major functional groups in phospholipids, for antioxidant activity in a sardine oil system and found choline and ethanolamine strongly inhibited increases in peroxide values in a sardine oil mixture during storage, whereas PA derivatives and glycerol did not. Antioxidant effect was thus attributable not only to the side-chain amino groups but also to the cooperative effect of the hydroxy group in the side chain. The precise mechanism of the antioxidant effect is unknown and various hypotheses include chelation of transition metals, a synergistic effect with vitamin E, and hydroperoxide decomposing activity (Ishihara, 1997).

PC enhanced feeding activity and diet ingestion rate of microdiets fed to sea bream larvae up to day 30 post-hatch, which may suggest a role for dietary phospholipids as an age-dependent feed attractant (Koven et al., 1998, 2001). An attractant property is likely to be dependent upon

the phospholipid head group as lecithin and certain L-amino acids had previously been tested for attractant activity for aquatic organisms including weatherfish (*Misgurnus anguillicaudatus*) and juvenile yellowtail (*Seriola quinqueradiata*) (Harada, 1987). The attraction activity of amino acids and their derivatives was ascertained to largely depend on both alpha -carboxyl and alpha -amino groups, but especially the former.

6.2 Improved digestion

Phospholipids are surface-active agents, or surfactants, and the main industrial use of lecithins is as lipid emulsifiers particularly in processed foods. Fish diets, of course, do not require the presence of phospholipids as emulsifiers per se, but their presence in the formulation may improve lipid emulsification and aid digestion in the intestine of the fish after consumption. Improved lipid digestibility in salmon fed diets containing soybean lecithin has been attributed to emulsification properties of the phospholipid (Hung et al., 1997), and other studies have shown that dietary phospholipids increased digestibility in juvenile fish (Craig and Gatlin, 1997; Kasper and Brown, 2003). However, supplementation with bile acids did not mimic the effect of dietary phospholipid in promoting growth in larval ayu (Kanazawa, 1983). In a study in gilthead sea bream larvae, a lecithin-supplemented diet substantially increased the uptake of labelled lipids from the diet and this was initially suggested to be via improved emulsification (Koven et al., 1993). However, this result was not substantiated in subsequent studies on gilthead sea bream fed with phospholipid-supplemented diets (Hadas et al., 2003). Phospholipid supplements also aided the absorption of dietary neutral lipids in European sea bass post-larvae, but this was not via a mechanism involving increased emulsification as diets containing other emulsifiers or lyso-phospholipid, which has greater emulsification properties than phospholipid, did not replicate the effect (Geurden et al., 1997c, 1998a). Therefore, there is little evidence to support a role for dietary phospholipids as aids to digestion.

6.3 Provision of essential components

6.3.1 Fatty acids (energy and essential fatty acids)

Phospholipids can be an important source of energy (fatty acids) in fish, particularly during embryonic and early larval development in species that produce phospholipid-rich eggs (Tocher, 1995). Larval fish at first feeding may be predisposed to digestion and metabolism of phospholipids and the use of fatty acids from phospholipids for energy (Sargent et al., 1997).

Dietary lipid is also important as a source of essential fatty acids (EFA), and phospholipids tend to be a richer source of EFA than neutral lipids such as triacylglycerols (Tocher, 1995). In addition, phospholipids may be superior to neutral lipids as a source of EFA in larval fish due to improved digestibility (Sargent et al., 1997, 1999b). Recently it was shown that phospholipids were the more efficient mode of supply for dietary EPA and DHA to sea bass larvae (Gisbert et al., 2005). However, in many experiments on phospholipid requirement, diets have been carefully formulated to discriminate a phospholipid effect from a fatty acid effect with, for instance, diets formulated to be EFA sufficient from neutral lipid alone (see Coutteau et al., 1997). In any case, most of the studies on phospholipid requirement have been performed with soybean lecithin and egg lecithin that are relatively rich in 18:2n-6 (and may have some 18:3n-3), but are deficient in the n-3 HUFA, EPA and DHA, required in abundance by larval fish (Coutteau et al., 1997; Sargent et al., 2002). In addition, further evidence that the beneficial effects of dietary phospholipids are not due to the provision of EFA is that soybean lecithin, lacking n-3 HUFA is as effective or more effective as lecithins from marine sources including fish eggs which are rich in n-3HUFA (Kanazawa et al., 1981, 1983a; Geurden et al., 1995b). Therefore, there is good evidence that the effects of dietary phospholipids on growth-promotion, survival and the prevention of malformations are not due to the provision of EFA. Indeed, diets containing very high levels of marine phospholipids and n-3HUFA actually induced skeletal malformations in sea bass larvae (Villeneuve et al., 2005, 2006).

6.3.2 Phosphorous

Phosphorous is a nutritionally important mineral due to its requirement for growth, bone mineralization, reproduction, nucleic acid synthesis, and energy metabolism (Lall, 2002). Deficiency signs include reduced growth and skeletal deformities and quantitative requirements have been determined for several fish species (Lall, 2002). As phosphate is low in most aquatic environments (non-polluted), feed is the main source of phosphorous, mainly from meals or premixes, with meat/bone meal > fish meal > plant meals. Bioavailability varies, but generally the inorganic (calcium and potassium salts) and organic (phospholipids) forms found in fish meals are more readily available to fish than the phytates (phytic acid salts) found in plant meals (Lall, 2002). Many studies on phospholipid requirements have used casein- or soy protein-based diets, but it is unlikely that this would be a problem with mineral premixes (Coutteau et al., 1997). In the only study to date, the effects of dietary phosphorus and phospholipid level on growth and phosphorus deficiency signs were investigated in juvenile (1 g) Japanese flounder

(Uyan et al., 2007). The results showed there was no interaction between dietary phosphorous and phospholipid suggesting that supply of phosphorous was not a mechanism for the growth promoting effects of dietary phospholipids.

6.3.3 Choline or inositol

It is known that choline cannot be synthesised in animals (Lykidis, 2007), but there may be evidence that some inositol can be synthesised in carp intestine and channel catfish (see Halver, 2002) and may not be required by juvenile sunshine bass (Deng et al., 2002). However, both choline and inositol are regarded as vitamins for fish, as there are known deficiency signs for both, and quantitative requirements for growth have been defined, at least for choline (Halver, 2002). Both choline and inositol are quite ubiquitous in feed ingredients including wheat germ, fish and plant (bean) meals, but they are often also supplemented in vitamin premixes. In studies with carp larvae, the growth-promoting and prevention of malformation effects of PC and PI were not mimicked by choline and inositol (Geurden et al., 1995b).

6.4 Provision of intact phospholipid

Over twenty years ago, based on his own series of studies with larval ayu, red sea bream and Japanese flounder, Kanazawa speculated that phospholipid biosynthesis itself may be limiting in larval fish. He reasoned that, as larval fish require an abundance of phospholipid for new tissue growth, “biosynthesis of phospholipids may not take place at a sufficiently fast rate to meet phospholipid requirement” and “requirement for dietary phospholipids may diminish with increasing age” (Kanazawa, 1985). However, it was another 10-15 years before there was a sufficient body of evidence to support the hypothesis of limited phospholipid biosynthetic ability in early life stages of fish.

Several studies showed that larval diets deficient in phospholipid could lead to the accumulation of lipid vacuoles (droplets) in the intestinal enterocytes (Fontagné et al., 1998; Olsen et al., 1999; Salhi et al., 1999; Liu et al., 2002). For example, intestinal steatosis (fat drop accumulation) was induced in carp larvae fed phospholipid-deficient diets and PC and, to a lesser extent, PI, prevented this (Fontagné et al., 1998). These studies led to the suggestion that intact dietary phospholipids may be required for the efficient exportation of dietary lipids from the enterocytes (Fontagné et al., 1998; Geurden et al., 1998b; Olsen et al., 1999; Salhi et al., 1999). Supporting this, in sea bream larvae fed phospholipid-deficient diets containing radiolabelled 18:1n-9, increased transport of ¹⁴C-18:1n-9 free fatty acid from enterocytes to tissues was

observed upon supplementation with PC (Hadas et al., 2003). As described in section 4.3, dietary lipids are transported from the intestine in the form of the large, neutral lipid-rich lipoproteins, chylomicrons and, to a lesser extent, VLDL and so it appeared that, in fish fed phospholipid-deficient diets, there may be insufficient lipoprotein synthesis. Lipoprotein production in the enterocyte involves a series of biosynthetic processes whereby neutral lipids, triacylglycerol and cholesteryl esters, form a large hydrophobic core surrounded by a thin coat of protein (apoproteins), phospholipid (particularly PC) and cholesterol (Vernier and Sire, 1986). Clearly, intact phospholipid is required for lipoprotein assembly, and so it was proposed that the stimulating effects of phospholipids in larval fish growth were due to the fish larvae having a limited ability to biosynthesise phospholipids *de novo* (Geurden et al., 1995b, 1999; Coutteau et al., 1997; Fontagné et al., 1998). However, the Golgi are intimately involved in the secretion of lipoproteins from the enterocytes, which may implicate another role for phospholipids in this mechanism. *De novo* PC synthesis has been shown to be required to replenish the Golgi-associated PC pool and to support Golgi-mediated secretion of tumour necrosis factor α in mouse macrophages (Tian and Jackowski, 2007).

Although the limiting biosynthesis hypothesis is quite widely accepted now, there is actually very little biochemical or enzymological data to support it. The major question is, what enzymatic step or steps are responsible for the limitation in phospholipid biosynthesis in the enterocyte? There is plenty of evidence that fish larvae readily reacylate dietary glycerides including lyso-phospholipids and so there is no limitation in acylation reactions (see Sargent et al., 2002). Thus, Sargent et al. (2002) speculated that the limitation in phospholipid biosynthesis in larvae is in forming the glycerophosphobase backbone (e.g. glycerophosphocholine or glycerophosphoethanolamine) of the phospholipid molecule. Unfortunately, due to the complexity of the phospholipid synthesis pathways (Lykidis, 2007), this does not narrow the search down significantly as there are many possible enzymatic steps that could be involved. Possibly the most obvious candidates are the CDP-choline and CDP-ethanolamine phosphotransferases (CPT and EPT) involved in the conversion of DAG to PC and PE, respectively. Further enzymes are required to produce the CDP-choline and CDP-ethanolamine required by the phosphotransferases, but synthesis of CDP-choline is not limiting as it was shown that CDP-choline could not mimic the effect of PC on the improvement of survival rates in larval ayu (Kanazawa, 1983). However, limitations in CPT and EPT cannot account for the observed effects of PI, as it is not formed via the CPT/EPT pathways. This suggests that the

limiting step would have to be further back in the phospholipid synthesis pathway, but there is no single enzyme that could be responsible for limiting both PC and PI synthesis. PA is the common intermediate in the production of PC and PI but different enzymes, phosphatidate phosphatase and CDP-DAG synthase, convert PA to DAG and CDP-DAG, the precursors of PC and PI, respectively (see section 3.1). There is no direct evidence and no studies investigating the pathways of phospholipid biosynthesis in larval fish and so the enzymic location(s) of any limitation in these pathways in larval and early juvenile fish is still unknown.

Sargent et al. (2002) also speculated that such apparent limitations in the ability of fish larvae to synthesise phospholipids *de novo* may not be surprising since, in their natural environment, larvae ingest live feed whose lipid is predominantly phospholipid. Thus, the larvae will seldom if ever be required to biosynthesise phospholipids extensively *de novo*. It is possible that early developing fish larvae have limited lipid biosynthetic capabilities in general, whether for fatty acids or lipid classes, including cholesterol.

7. Dietary phospholipids and sustainability

7.1 Sustainability

Exploitation and, indeed, over-exploitation of wild fisheries has meant that an increasing proportion of fish for human consumption is provided by aquaculture, which has been expanding at around 10% per year over the last decade (Tidwell and Allan, 2002). Paradoxically, diets for aquaculture have been based traditionally on fishmeals and oils, themselves derived from industrial feed-grade or reduction fisheries, as the predominant protein and lipid sources (Sargent and Tacon, 1999; Pike, 2005). This is certainly the case in Europe where intensive farming is based largely on carnivorous species that feed high up in the food chain, particularly salmonid and marine species. The demand for these marine products is rapidly increasing and current estimates suggest requirements for aquaculture feeds could exceed global supplies of fish oil and fish meal within the next two to eight years, respectively (FAO, 2006). Therefore, if aquaculture is to continue to expand and supply more of the global demand for fish, alternatives to fish oil and meal must be found (Barlow, 2000; Tacon, 2004). Under the present regulations, other animal-based meals such as feather, meat, bone and blood meals are not permitted for use in fish diet formulations in the European Union. The only obvious, and sustainable, alternative to marine meals are plant meals, but these contain very little, if any, phospholipid in comparison with fish meals (Sargent et al., 2002). Therefore, replacing the fish meal component of diets with plant

meals will result in greatly reduced phospholipid concentrations in diets. This means that possible phospholipid requirements, if any, of adult and on-growing fish, may become an important and significant issue in the future as plant meals comprise an increasing proportion of diet formulations.

As mentioned previously, there have been no studies investigating phospholipids requirements in adult fish, or individuals of much more than 5 g. However, interestingly, accumulation of lipid droplets was reported in the hindgut of juvenile gilthead sea bream fed diets based on plant proteins and fish oil (Sitjà-Bobadilla et al., 2005). Similar histological alterations in gilthead sea bream intestine had been reported earlier in studies investigating replacement of fish oil with vegetable oils (Caballero et al., 2003). Similarly, substantial accumulation of lipid droplets was observed in the gastrointestinal tract of salmonids fed with vegetable oils, and this condition was reversed by phospholipid supplementation (Olsen et al., 1999, 2003). Liver steatosis was also observed in gilthead sea bream fed plant meals (Sitjà-Bobadilla et al., 2005) or vegetable oils (Caballero et al., 2004; Wassef et al., 2007). Perhaps related, there is some evidence that soybean PC may alleviate similar signs of liver disease in human and animal studies (Canty and Zeisel, 1994; Ipatova et al., 2004). Therefore, there is circumstantial evidence that perhaps may indicate that dietary phospholipid supplementation may be beneficial in larger (on-growing juvenile or adult) fish particularly when fed diets high in plant products.

7.2 Dietary sources of phospholipids

7.2.1 Oils

The phospholipid contents of refined oils, both fish oils and vegetable oils, are very low as they are removed during the normal refining processes. Degumming separates the phospholipids as a gum. Crude soybean oil contains around 1.5 - 3.1 % total phospholipid that is removed to become the by-product, soybean lecithin (Daniel, 2004). As mentioned above, soybean lecithin can have very variable phospholipid contents and class compositions but most commonly contains around 50 – 60 % total phospholipid, and around 13 - 18% PC, 10 - 15% PE, 10 – 15 % PI and 5 – 12 % PA (Daniel, 2004). Crude sunflower oil can have a phospholipid content of 0.5 – 1 % and is also used to produce lecithin having similar phospholipid class profile to soybean lecithin. Most other crude vegetable oils have less phospholipid, up to about 0.5 % total phospholipid before degumming, but due to the large production, rapeseed/canola lecithin is also commercially available (Daniel, 2004).

7.2.2 Meals

Unless rather crude oils are used, the vast majority of phospholipids in fish diets will be provided by meals and other essentially “protein” components. Residual lipid contents in fish meals can be reasonably high, varying from 5 – 13 % of weight, with a triacylglycerol/phospholipid ratio of around 2:1 (de Koning, 2005). Phospholipid content of meals has been suggested as a quality index as fresh fish generally contain a rather constant amount of phospholipids that decrease rapidly upon storage (de Koning, 2005). In fish feeds, phospholipid can account from 5 – 25 % of the total lipid depending on lipid content and formulation of the feed, and analytical method used (Johnson and Barnett, 2003). Individual phospholipid class compositions for fish meal are difficult to find, but should generally reflect the phospholipid composition of fish (PC, PE, PS, PI, PA and lyso-phospholipids) (see Tocher, 1995). Generally, most plant meals have much lower levels of residual phospholipids, mainly due to the original products (seeds/beans etc.) having much lower phospholipid contents. However, soybean has the highest levels with full-fat soybean meal (FFSM or soy flour) having about 20 – 25 % lipid with only around 0.3 - 0.6 % phospholipid, primarily PC, PE, PI and PA. Defatted soybean meal, prepared by solvent extraction, has < 1 % lipid with soybean cake (produced by pressing) possibly having slightly higher lipid content. Very little information is available about the lipid and phospholipid contents and compositions of other plant meals.

7.3 Analysis of phospholipid content and composition in fish diets and ingredients

Although not an exhaustive list, Table 2 indicates that the phospholipid content of commercially available fish meals can vary considerably. In this sample of four meals, the phospholipid content varied from around 17 to 27 % of total lipids. The phospholipid content appears to be largely independent of lipid content of the meals, which varied from around 6 to 10 % of the wet weight giving phospholipids contents of between 1.2% and almost 2.4% of wet weight of the fish meals. The predominant lipid classes were PC and PE in a roughly 2:1 ratio, followed by around 1 - 3 % of the other phospholipids classes PI, PS and sphingomyelin. Standard, commercial feeds formulated predominantly with fish meal will therefore show a range of phospholipid contents from around 1 % to around 4 % depending upon lipid content of the diet (Table 2). Thus, diets with lower lipid contents (lower levels of added oil) have a higher phospholipid content that is predominantly PC, with PE also detected in the diets with higher phospholipid content.

The phospholipid contents and compositions of a variety of feed ingredients including alternative protein sources are presented in Table 3. The two fish oils analysed (northern and southern hemisphere oils) did not contain any detectable phospholipids. However, the rapeseed oil contained around 1.7% phospholipids, specifically PE and PC. The cereal products, which had reasonably high lipid contents (~ 6 to 10%), showed markedly different phospholipid contents with corn gluten having very low levels (0.5%) whereas the phospholipid content of wheat gluten was almost 7% of total lipids, although it also contained high levels of galactolipids and other glycolipids (20%). Sunflower cake, which also had a relatively high lipid content at 8 %, contained phospholipids at 9 % of total lipid and a lower amount of galactolipid. The high protein soy products analysed (HP soya and SPC 60) contained only 1 – 2 % lipid, but it was relatively rich in phospholipids amounting to 28 – 40 % of total lipid. Peas were not lipid-rich (1.8%) but almost 30% was present as phospholipids. The animal-derived protein alternatives showed markedly different phospholipids contents. The haemoglobin and blood meals, although relatively low in lipids contained phospholipids at between 22 and 31% of total lipid, whereas the feather meals contained 6-7% lipid, but had very low or no phospholipids. Binders (field beans and kidney beans) were also low fat products with 12-28% of the total lipid being phospholipids. The rank order for the phospholipid classes in the soy and legume products was generally PC > PE > PS > PI. The legumes and soy products also contained galacto- and other glycolipids at around about 10% of total lipid. The lecithin analysed (BergaPur) was 75 % lipid of which two-thirds was phospholipids (PC, PE, PS and PI in roughly similar proportions), along with around 15 % glycolipids and 17% other neutral lipids, mainly triacylglycerol.

7.4 Dietary cholesterol

Although not the subject of the present review it is pertinent to mention that, in addition to reducing the level of dietary phospholipids, the increasing proportion of plant meals in dietary formulations will also reduce the level of dietary cholesterol. This may not be obvious from the lipid class compositions presented in Tables 2 and 3, as these data were generated by thin-layer chromatography (TLC) followed by staining and densitometric analysis (Henderson and Tocher, 1992), and this method cannot distinguish between cholesterol and other sterols. However, cholesterol is the predominant sterol in animals, whereas other sterols are present in plant tissues (Padley et al., 1986). The TLC/densitometric method also has other confounding factors such as co-eluting lipids (e.g. diacylglycerols) and the fact that sterols are generally overestimated as a result of their different staining compared to fatty acid-containing lipids (Henderson and Tocher,

1992). Indeed, quantitative cholesterol determination remains a difficult analysis (Christie, 1989). The cholesterol (as opposed to total sterol) content of traditional salmonid feeds and a variety of alternative feed ingredients as determined by gas chromatography (see Christie, 1989) is listed in Table 4. This clearly shows that plant products, both oils and meals, contain virtually no cholesterol. Feeds traditionally formulated with fish meal and fish oil will provide at least 1 g of cholesterol per kg feed. Substitution of fish meal and/or fish oil will greatly reduce the level of cholesterol supplied by the feed. The only alternatives to fish meal and oils that could supply dietary cholesterol are the other animal-derived products, the blood and feather meals (Table 4).

Although there have been a few early studies investigating dietary supplementation of cholesterol in salmonids, primarily in relation to the development of coronary arteriosclerotic lesions (Farrell and Munt, 1981; Farrell et al., 1986), there are few reports on the effects of dietary cholesterol on growth or metabolism in fish. However, the effects of dietary cholesterol supplementation on growth, organ indices, digestibility of cholesterol and macronutrients, and tissue fatty acid compositions have been investigated in Atlantic salmon (Bjerkeng et al., 1999). This study showed that dietary cholesterol level had no significant effect on specific growth rate (SGR), mortality, apparent digestibility coefficients of macronutrients, and total lipid content. However, both hepatic cholesterol concentration and hepatosomatic index were increased by the dietary cholesterol supplement. In a very recent study, it was shown that genes of the cholesterol biosynthesis pathway in liver were up-regulated in salmon fed a vegetable oil blend compared to fish fed a diet containing fish oil (Taggart et al., 2008). The effects on cholesterol metabolism suggest this may also be an area requiring consideration in the future as dietary cholesterol concentrations decline as the proportion of plant meals and oils in dietary formulations increases.

8. Conclusions

Intact phospholipids are required for optimal growth, survival, prevention of skeletal deformities and, possibly, stress resistance in larval and early juvenile fish, of both marine and freshwater species. The quantitative requirement appears to diminish with age, and no requirement has been demonstrated in fish greater than 5 g, although there are very few studies and most are of short duration. Qualitative requirements suggest that PC may be more important for growth and PI for development. The mechanism of apparent essentiality of phospholipids in larval and early juvenile fish is not clear. Most potential mechanisms can be discounted, leaving “limiting biosynthesis” as the leading hypothesis. It is suggested that phospholipid biosynthesis pathways

are inefficient or not fully developed in larvae of a number of (possibly all) fish species. In larvae fed diets rich in triacylglycerol, lack of sufficient dietary phospholipid limits lipoprotein synthesis in enterocytes, leading to impaired transport of lipid (energy supply) nutrients to tissues. Dietary phospholipid may provide intact glycerophosphobase backbones, which bypasses this limitation. Thus, growth stimulation is due to improved transport, assimilation and utilization of dietary lipid.

References

- Arias, H.R., Barrantes, F.J., 1990. Phosphoinositides and inositol phosphates in *Discopyge tschudii* electrocyte membranes. Int. J. Biochem. 22, 1387-1392.
- Barlow, S., 2000. Fishmeal and oil: sustainable feed ingredients for aquafeeds. Global Aquacult. Advocate 4, 85-88.
- Bell, M.V., Dick, J.R., 1993. Molecular species composition of the major glycerophospholipids from the muscle, liver, retina and brain of cod (*Gadus morhua*). Lipids 26, 565-573.
- Bell, M. V., Sargent, J.R., 1987. Protein kinase C activity in the spleen of trout (*Salmo gairdneri*) and the rectal gland of dogfish (*Scyliorhinus canicula*), and the effects of phosphatidylserine and diacylglycerol containing (n-3) polyunsaturated fatty acids. Comp. Biochem. Physiol. 87B, 875-880.
- Bell, M.V., Tocher, D.R., 1989. Molecular species composition of the major phosphoglycerides in brain and retina from trout : Occurrence of high levels of di-(n-3) polyunsaturated fatty acid species. Biochem. J. 264, 909-914.
- Berridge, M.J., 1987. Inositol trisphosphate and diacylglycerol: Two interacting second messengers. Ann. Rev. Biochem. 56, 159-194.
- Bjerkeng, B., Storebakken, T., Wathne, E., 1999. Cholesterol and short-chain fatty acids in diets for Atlantic salmon *Salmo salar* (L.): effects on growth, organ indices, macronutrient digestibility, and fatty acid composition. Aquacult. Nutr. 5, 181-191.
- Caballero, M.J., Izquierdo, M.S., Kjørsvik, E., Montero, D., Socorro, J., Fernández, A.J. and Rosenlund, G., 2003. Morphological aspects of intestinal cells from gilthead seabream (*Sparus aurata*) fed diets containing different lipid sources. Aquaculture 225, 325-340.
- Caballero, M.J., Izquierdo, M.S., Kjørsvik, E., Fernández, A.J., Rosenlund, G., 2004. Histological alterations in the liver of sea bream, *Sparus aurata* L., caused by short- or long-

914 term feeding with vegetable oils. Recovery of normal morphology after feeding fish oil as the
 915 sole lipid source. J. Fish Dis. 27, 531-541.

916 Cabot, M.C., Faulkner, L.A., Lackey, B.J., Snyder, F., 1984. Vertebrate class distribution of 1-
 917 alkyl-2-acetyl-*sn*-glycero-3-phosphocholine acetylhydrolase in serum. Comp. Biochem.
 918 Physiol. 78B, 37-40.

919 Cahu, C., Zambonino-Infante, J., 2001. Substitution of live food by formulated diets in marine
 920 fish larvae. Aquaculture 200, 161-180.

921 Cahu, C., Zambonino Infante, J., Takeuchi, T., 2003a. Nutritional components affecting skeletal
 922 development in fish larvae. Aquaculture 227, 245-258.

923 Cahu, C., Zambonino Infante, J., Barbosa, V., 2003b. Effect of dietary phospholipid level and
 924 phospholipid:neutral lipid value on the development of sea bass (*Dicentrarchus labrax*) larvae
 925 fed a compound diet. Br. J. Nutr. 90, 21-28.

926 Camara, M.R., 1994. Dietary phosphatidylcholine requirements of *Penaeus japonicus* Bate and
 927 *Penaeus vannamei* Boone (Crustacea, Decapoda, Penaeidae). PhD Thesis, Univ. Gent,
 928 Belgium, 173 pp.

929 Canty, D.J., Zeisel, S.H., 1994. Lecithin and choline in human health and disease. Nutr. Rev. 52,
 930 327-339.

931 Chapman, M.J., 1980. Animal lipoproteins: chemistry, structure, and comparative aspects. J.
 932 Lipid Res. 21, 789-853.

933 Chapman, M.J., Goldstein, S., Mills, G.L., Leger, C., 1978. Distribution and characterization of
 934 the serum lipoproteins and their apoproteins in the rainbow trout (*Salmo gairdneri*).
 935 Biochemistry 17, 4455-4464.

936 Christie, W.W., 1989. Gas Chromatography and Lipids. The Oily Press, Dundee, UK.

937 Coutteau, P., Geurden, I., Camara, M.R., Bergot, P., Sorgeloos, P., 1997. Review on the dietary
 938 effects of phospholipids in fish and crustacean larviculture. Aquaculture 155, 149-164.

939 Cowey, C.B., Bell, J.G., Knox, D., Fraser, A., Youngson, A., 1985. Lipids and antioxidant
 940 systems in developing eggs of salmon (*Salmo salar*). Lipids 20, 567-572.

941 Craig, S.R., Gatlin, D.M., 1997. Growth and body composition of juvenile red drum (*Sciaenops*
 942 *ocellatus*) fed diets containing lecithin and supplemental choline. Aquaculture 151, 259-267.

943 Daniel, K.T., 2004. Soy lecithin: From sludge to profit.
 944 (<http://www.westonaprice.org/soy/lecithin.html>)

945 Deng, D., Hemre, G., Wilson, R.P., 2002. Juvenile sunshine bass (*Morone chrysops* *Morone*
 946 *saxatilis*) do not require dietary *myo*-inositol. Aquaculture 213, 387-393.

947 De Koning, A.J., 2005. Properties of South African fish meal: a review. S. African J. Sci. 101,
948 21-25.

949 Farber, S.A., Olson, E.S., Clark, J.D., Halpern, M.E., 1999. Characterization of Ca²⁺ dependent
950 phospholipase A₂ activity during zebrafish embryogenesis. J. Biol. Chem. 274, 19338-19346.

951 Farrell, A.P., Munt, B., 1981. Dietary induced hypercholesterolemia and arteriosclerosis in brook
952 trout (*Salvelinus fontinalis*). Physiologist 24, 43.

953 Farrell, A.P., Saunders, R.L., Freeman, H.C., Mommsen, T.P., 1986. Arteriosclerosis in Atlantic
954 salmon. Effects of dietary cholesterol and maturation. Arteriosclerosis 6, 453-461.

955 Finn, R.N., Henderson, R.J., Fyhn, H.J. (1995. Physiological energetics of developing embryos
956 and yolk-sac larvae of Atlantic cod (*Gadus morhua*). 2. Lipid metabolism and enthalpy
957 balance. Mar. Biol. 124, 371-379.

958 Fontagné, S., Geurden, I., Escaffre, A.-M., Bergot, P., 1998. Histological changes induced by
959 dietary phospholipids in intestine and liver of common carp (*Cyprinus carpio* L.) larvae.
960 Aquaculture 161, 213-223.

961 Fontagné, S., Burtairé, L., Corraze, G., Bergot, P., 2000. Effects of dietary medium-chain
962 triacylglycerols (tricaprylin and tricaproin) and phospholipid supply on survival, growth and
963 lipid metabolism in common carp (*Cyprinus carpio* L.) larvae. Aquaculture 190, 289-303.

964 Food and Agricultural Organisation (FAO), 2006. State of world aquaculture 2006. FAO
965 Fisheries Technical paper No. 500, FAO, Rome, 134 pp.

966 Fraser, A. J., Gamble, J.C., Sargent, J.R., 1988. Changes in lipid content, lipid class composition
967 and fatty acid composition of developing eggs and unfed larvae of cod (*Gadus morhua*).
968 Mar. Biol. 99, 307-313.

969 Fremont, L., Leger, C., 1981. The transport of plasma lipids. In: Fontaine, M., (Ed.), Nutrition
970 des Poissons, Actes du Colloque CNERNA, Paris, 1979. Centre National de la Recherche
971 Scientifique, Paris, pp. 263-282.

972 Geurden, I., Coutteau, P., Sorgeloos, P., 1995a. Dietary phospholipids for European sea bass
973 (*Dicentrarchus labrax* L.) during first ongrowing. In: Lavens, P., Jaspers, E., Roelants, I.
974 (Eds.), Larvi '95 – Fish and Shellfish symposium, Gent, Belgium. EAS Spec. Publ. 24.
975 European Aquaculture Society, 175-178.

976 Geurden, I., Radünz-Neto, J., Bergot, P., 1995b. Essentiality of dietary phospholipids for carp
977 (*Cyprinus carpio* L.) larvae. Aquaculture 131, 303-314.

978 Geurden, I., Charlon, N., Marion, D., Bergot, P., 1997a. Influence of purified soybean
979 phospholipids on early development of common carp. Aquacult. Int. 5, 137-149.

980 Geurden, I., Coutteau, P., Sorgeloos, P., 1997b. Effect of dietary phospholipid supplementation
981 on growth and fatty acid composition of European sea bass (*Dicentrarchus labrax* L.) and
982 turbot (*Scophthalmus maximus* L.) juveniles from weaning onwards. Fish Physiol. Biochem.
983 16, 259-272.

984 Geurden, I., Coutteau, P., Sorgeloos, P., 1997c. Increased docosahexaenoic acid levels in total
985 and polar lipid of European sea bass (*Dicentrarchus labrax* L.) postlarvae fed vegetable or
986 animal phospholipids. Mar. Biol. 129, 689-698.

987 Geurden, I., Marion, D., Charlon, N., Coutteau, P., Bergot, P., 1998a. Comparison of different
988 soybean phospholipidic fractions as dietary supplements for common carp, *Cyprinus carpio*,
989 larvae. Aquaculture 161, 225-235.

990 Geurden, I., Bergot, P., Schwarz, L., Sorgeloos, P., 1998b. Relationship between dietary
991 phospholipids class composition and neutral lipid absorption in postlarval turbot. Fish
992 Physiol. Biochem. 19, 217-228.

993 Geurden, I., Bergot, P., Van Ryckeghem, K., Sorgeloos, P., 1999. Phospholipid composition of
994 common carp (*Cyprinus carpio*) larvae starved or fed different phospholipid classes.
995 Aquaculture 171, 93-107.

996 Gisbert, E., Villeneuve, L., Zambonino-Infante, J.L, Quazuguel, P., Cahu, C.L., 2005. Dietary
997 phospholipids are more efficient than neutral lipids for long-chain polyunsaturated fatty acid
998 supply in European sea bass *Dicentrarchus labrax* larval development. Lipids 40, 609-618.

999 Hadas, E., Koven, W., Sklan, D., Tandler, A., 2003. The effect of dietary phosphatidylcholine on
1000 the assimilation and distribution of ingested free oleic acid (18:1n-9) in gilthead sea bream
1001 (*Sparus aurata*) larvae. Aquaculture 217, 577-588.

1002 Halver, J.E., 2002. The vitamins. In: Halver, J.E., Hardy, R.W. (Eds.), Fish Nutrition, 3rd Edition,
1003 Academic Press, San Diego, pp. 61-141.

1004 Hamza, N., Mhetli, M., Khemis, I.B., Cahu, C., Kestemont, P., 2008. Effect of dietary
1005 phospholipids levels on performance, enzyme activities and fatty acid composition of
1006 pikeperch (*Sander lucioperca*) larvae. Aquaculture 275, 274-282.

1007 Hao, F., Tan, M., Xu, X., Han, J., Miller, D.D., Tigyi, G., Cui, M.-Z., 2007. Lysophosphatidic
1008 acid induces prostate cancer PC3 cell migration via activation of LPA₁, p42 and p38 α .
1009 Biochim. Biophys. Acta 1771, 883-892.

1010 Harada, K., 1987. Relationships between structure and feeding attraction activity of certain
1011 L-amino acids and lecithin in aquatic animals. Nippon Suisan Gakkaishi 53, 2243-2247.

1012 Hazel, J. R., 1990. Adaptation to temperature: phospholipid synthesis in hepatocytes of rainbow

1013 trout. Am. J. Physiol. 258 (Regulatory Integrative Comp. Physiol. 27) R1495-1501.

1014 Hazel, J.R., Williams, E.E., 1990. The role of alterations in membrane lipid composition in
 1015 enabling physiological adaptation of organisms to their physical environment. Prog. Lipid Res.
 1016 29, 167-227.

1017 Henderson, R.J., Tocher, D.R. 1987. Lipid composition and biochemistry of freshwater fish.
 1018 Prog. Lipid Res. 26, 281-347.

1019 Henderson, R.J., Tocher, D.R., 1992. Thin-layer chromatography. In: Hamilton, R.J., Hamilton,
 1020 S. (Eds.), Lipid Analysis : A Practical Approach. IRL Press, Oxford, pp. 65-111.

1021 Hochachka, P.W., Mommsen, T.P. (Eds.), 1995. Biochemistry and Molecular Biology of Fishes,
 1022 Environmental and Ecological Biochemistry, Vol. 5. Elsevier Press, Amsterdam.

1023 Holub, B. J., Connor, J.T.H., Slinger, S.J., 1975a. Incorporation of glycerol-3-phosphate into
 1024 hepatic lipids of rainbow trout, *Salmo gairdneri*. J. Fish. Res. Board Can. 32, 61-64.

1025 Holub, B. J., Nilsson, K., Piekarski, J., Slinger, S.J., 1975b. Biosynthesis of lecithin by the
 1026 CDP-choline pathway in liver microsomes of rainbow trout, *Salmo gairdneri*. J. Fish. Res.
 1027 Board Can. 32, 1633-1637.

1028 Holub, B.J., Piekarski, J., Cho, C.Y., Slinger, S.J., 1976. Incorporation of fatty acids into
 1029 phosphatidylcholine by acyl-CoA: 1-acyl-*sn*-glycero-3-phosphorylcholine acyltransferase in
 1030 liver of rainbow trout, *Salmo gairdneri*. J. Fish. Res. Board Can. 33, 2821-2826.

1031 Hung, S.S., Lutes, P.B., 1988. A preliminary study on the non-essentiality of lecithin for
 1032 hatchery-produced juvenile white sturgeon (*Acipenser transmontanus*) Aquaculture 68, 353-
 1033 360.

1034 Hung, S.S.O., Berge, G.M., Storebakken, T., 1997. Growth and digestibility of soya lecithin and
 1035 choline chloride in juvenile Atlantic salmon. Aquacult. Nutr. 3, 141-144.

1036 Iijima, N., Zama, K., Kayama, M., 1983. Effect of oxidized lipids on the metabolic pathway of
 1037 lipid biosynthesis in the intestine of the carp. Bull. Jpn. Soc. Sci. Fish. 49, 1465-1470.

1038 Iijima, N., Aihara, M., Kayama, M., Okazaki, M., Hara, I., 1989. Comparison of carp plasma
 1039 lipoproteins under starved and fed conditions. Nippon Suissan Gakkaishi 55, 2001-2007.

1040 Iijima, N., Aida, S. Mankura, M., Kayama, M., 1990. Intestinal absorption and plasma transport
 1041 of dietary triglyceride and phosphatidylcholine in the carp (*Cyprinus carpio*). Comp.
 1042 Biochem. Physiol. 96A, 45-56.

1043 Ipatova, O.M., Prozorovskaia, N.N., Torkhovskaia, T.I., Baranova, V.S., Guseva, D.A., 2004.
 1044 Biological effects of the soybean phospholipids. Biomed. Khim. 50, 436-450.

1045 Iritani, N., Ikeda, Y., Fukuda, H., Katsurada, A., 1984. Comparative study of lipogenic enzymes
1046 in several vertebrates. *Lipids* 19, 825-835.

1047 Ishihara, K., 1997. Antioxidant mechanisms of phospholipid. *Bull. Natl. Res. Inst. Fish. Sci.*
1048 (Japan) 8, 139-146.

1049 Johnson, R.B., Barnett, H.J., 2003. Determination of fat content in fish feed by supercritical fluid
1050 extraction and subsequent lipid classification of extract by thin layer chromatography-flame
1051 ionization detection. *Aquaculture* 216, 263-282.

1052 Kagan, V. E., Tyurin, V.A., Gorbunov, N.V., Prilipko, L.L., Chelomin, V.P., 1984. Are changes
1053 in the microviscosity and an asymmetrical distribution of phospholipids in the membrane
1054 necessary conditions for signal transmission? A comparison of the mechanisms of signal
1055 transmission in plasma membranes of brain synaptosomes and photoreceptor membranes of
1056 the retina. *J. Evol. Biochem. Physiol.* 20, 6-11.

1057 Kanazawa, A., 1983. Effect of phospholipids on aquatic animals. *Feed Oil Abst.* 18, 1-5.

1058 Kanazawa, A., 1993. Essential phospholipids of fish and crustaceans. In: Kaushik, S.J., Luquet,
1059 P. (Eds.), *Fish Nutrition in Practice. IV International Symposium on Fish Nutrition and*
1060 *Feeding*, INRA, France, National Institute for Agricultural Research, pp. 519-530.

1061 Kanazawa, A., Teshima, S., Inamori, S., Iwashita, T., Nagao, A., 1981. Effects of phospholipids
1062 on survival rate and incidence of malformation in the larval ayu. *Mem. Fac. Fish., Kagoshima*
1063 *Univ.* 30, 301-309.

1064 Kanazawa, A., Teshima, S., Kobayashi, T., Takae, M., Iwashita, T., Uehara, R., 1983a. Necessity
1065 of phospholipids for growth of the larval ayu. *Mem. Fac. Fish. Kagoshima Univ.* 32, 115-120.

1066 Kanazawa, A., Teshima, S., Inamori, S., Matsubara, H., 1983b. Effects of dietary phospholipids
1067 on growth of the larval red sea bream and knife jaw. *Mem. Fac. Fish., Kagoshima Univ.* 32,
1068 109-114.

1069 Kanazawa, A., Teshima, S. and Sakamoto, M., 1985. Effects of dietary bonito-egg phospholipids
1070 and some phospholipids on growth and survival of the larval ayu, *Plecoglossus altivelis*. *Z.*
1071 *Angew. Ichthyologie* 4, 165-170.

1072 Kasper, C.S., Brown, P.B., 2003. Growth improved in juvenile Nile tilapia fed
1073 phosphatidylcholine. *N. Am. J. Aquacult.* 65, 39-43.

1074 Koven, W.M., 2003. Key factors influencing juvenile quality in mariculture: A review. *Israel J.*
1075 *Aquacult.* 55, 283-297.

1076 Koven, W.M., Kolkovski, S., Tandler, A., Kissil, G.W., Sklan, D., 1993. The effect of dietary
1077 lecithin and lipase, as a function of age, on n-9 fatty acid incorporation in the tissue lipids of
1078 *Sparus aurata* larvae. Fish Physiol. Biochem. 10, 357-364.

1079 Koven, W.M., Henderson, R.J., Sargent, J.R., 1994. Lipid digestion in turbot (*Scophthalmus*
1080 *maximus*). 1: Lipid class and fatty acid composition of digesta from different segments of the
1081 digestive tract. Fish Physiol. Biochem. 13, 69-79.

1082 Koven, W.M., Parra, G., Kolkovski, S., Tandler, A., 1998. The effect of dietary
1083 phosphatidylcholine and its constituent fatty acids on microdiet ingestion and fatty acid
1084 absorption rate in gilthead seabream, *Sparus aurata*, larvae. Aquacult. Nutr. 4, 39-45.

1085 Koven, W.M., Kolkovski, S., Hadas, E., Gamsiz, K., Tandler, A., 2001. Advances in the
1086 development of microdiets for gilthead seabream, *Sparus aurata*: a review. Aquaculture 194,
1087 107-121.

1088 Kuo, J.F., Anderson, R.G.G., Wise, B.C., Mackerlova, L., Salomonsson, I., Brackett, N.L.,
1089 Katoh, N., Shoji, M., Wrenn, R.W., 1980. Calcium-dependent protein kinase: widespread
1090 occurrence in various tissues and phyla of the animal kingdom and comparison of effects of
1091 phospholipid, calmodulin, and trifluoperazine. Proc. Natl. Acad. Sci. USA 77, 7039-7043.

1092 Kvale, A., Yufera, M., Nygard, E., Aursland, K., Harboe, T., Hamre, K., 2006. Leaching
1093 properties of three different microparticulate diets and preference of the diets in cod (*Gadus*
1094 *morhua* L.) larvae. Aquaculture 251, 402-415.

1095 Lall, S.P., 2002. The minerals. In: Halver, J.E., Hardy, R.W. (Eds.), Fish Nutrition 3rd Edition,
1096 Academic Press, San Diego, pp. 259-308.

1097 Leslie, J. M., Buckley, J.T., 1975. Phospholipid composition of goldfish (*Carassius auratus* L.)
1098 liver and brain and temperature-dependence of phosphatidylcholine synthesis. Comp.
1099 Biochem. Physiol. 55B, 335-337.

1100 Liu, J., Caballero, M.J., Izquierdo, M.S., El-Sayed Ali, T., Hernández-Cruz, C.M., Valencia, A.,
1101 Fernández-Palacios, H., 2002. Necessity of dietary lecithin and eicosapentaenoic acid for
1102 growth, survival, stress resistance and lipoprotein formation in gilthead sea bream *Sparus*
1103 *aurata*. Fisheries Sci. 68, 1165-1172.

1104 Lykidis, A., 2007. Comparative genomics and evolution of eukaryotic phospholipid biosynthesis.
1105 Prog. Lipid Res. 46, 171-199.

1106 MacQueen Leifson, R., Homme, J.M., Jøstensen, J.P., Lie, Ø., Myklebust, R., Strøm, T., 2003.
 1107 Phospholipids in formulated start-feeds – Effect on turbot (*Scophthalmus maximus* L.) larval
 1108 growth and mitochondrial alteration in enterocytes. *Aquacult. Nutr.* 9, 43-54.

1109 Moschetta, A., Xu, F., Hagey, L.R., van Berge-Henegouwen, G.P., van Erpecum, K.J., Browsers,
 1110 J.F., Cohen, J.C., Bierman, M., Hobbs, H.H., Steinbach, J.H., Hofmann, A.F., 2005. A
 1111 phylogenetic survey of biliary lipids in vertebrates. *J. Lipid Res.* 46, 2221-2232.

1112 Mourente, G., Vazquez, R., 1996. Changes in the content of total lipid, lipid classes and fatty
 1113 acids of developing eggs and unfed larvae of the Senegal sole, *Solea senegalensis* Kaup. *Fish*
 1114 *Physiol. Biochem.* 15, 221-235.

1115 Mourente, G., Rodriguez, A., Grau, A., Pastor, E., 1999. Utilization of lipids by *Dentex dentex* L.
 1116 (Osteichthyes, Sparidae) larvae during lecithotrophia and subsequent starvation. *Fish Physiol.*
 1117 *Biochem.* 21, 45-58.

1118 Neas, N.P., Hazel, J.R., 1985. Phospholipase A₂ from liver microsomal membrane of thermally
 1119 acclimated rainbow trout. *J. Exp. Zool.* 233, 51-60.

1120 Nelson, G.J., Shore, V.G., 1974. Characterization of the serum high-density lipoproteins and
 1121 apolipoproteins of pink salmon. *J. Biol. Chem.* 249, 536-542.

1122 Noaillac-Depeyre, J., Gas, N., 1974. Fat absorption by the enterocytes of the carp (*Cyprinus*
 1123 *carpio* L.). *Cell Tissue Res.* 155, 353-365.

1124 Nordgreen, A., Hamre, K., Langdon, C., 2007. Development of lipid microbeads for delivery of
 1125 lipid and water-soluble materials to *Artemia*. *Aquaculture* 273, 614-623.

1126 Olsen, R.E., Ringø, E., 1997. Lipid digestibility in fish: a review. *Recent Res. Devel. Lipids Res.*,
 1127 1, 199-265.

1128 Olsen, R., Henderson, R.J., Pedersen, T., 1991. The influence of dietary lipid classes on the fatty
 1129 acid composition of small cod (*Gadus morhua* L.) juveniles reared in an enclosure in northern
 1130 Norway. *J. Exp. Mar. Biol. Ecol.* 148, 59-76.

1131 Olsen, R.E., Myklebust, R., Kaino, T., Ringø, E., 1999. Lipid digestibility and ultrastructural
 1132 changes in the enterocytes of Arctic char (*Salvelinus alpinus* L.) fed linseed oil and soybean
 1133 lecithin. *Fish Physiol. Biochem.* 21, 35-44.

1134 Olsen, R.E., Dragnes, B.T., Myklebust, R., Ringø, E., 2003. Effect of soybean oil and soybean
 1135 lecithin on intestinal lipid composition and lipid droplet accumulation of rainbow trout,
 1136 *Oncorhynchus mykiss* Walbaum. *Fish Physiol. Biochem.* 29, 181-192.

1137 Ono, H., Iijima, N., 1998. Purification and characterization of phospholipase A₂ isoforms from
 1138 the hepatopancreas of red sea bream, *Pagrus major*. Fish Physiol. Biochem. 18, 135-147.

1139 Oxley, A., Torstensen, B.E., Rustan, A.C., Olsen, R.E., 2005. Enzyme activities of intestinal
 1140 triacylglycerol and phosphatidylcholine biosynthesis in Atlantic salmon (*Salmo salar* L.).
 1141 Comp. Biochem. Physiol. 141B, 77-87.

1142 Oxley, A., Jutfelt, F., Sundell, K., Olsen, R.E., 2007. *Sn*-2-monoacylglycerol, not glycerol, is
 1143 preferentially utilised for triacylglycerol and phosphatidylcholine biosynthesis in Atlantic
 1144 salmon (*Salmo salar* L.) intestine. Comp. Biochem. Physiol. 146B, 115-123.

1145 Padley, F.B., Gunstone, F.D., Harwood, J.L., 1986. Occurrence and characteristics of oils and
 1146 fats. In: Gunstone, F.D., Harwood, J.L., Padley, F.B. (Eds.), The Lipid Handbook. Chapman
 1147 Hall, London, pp. 49-170.

1148 Perez, J.A., Rodriguez, C., Henderson, R.J., 1999. The uptake and esterification of radiolabelled
 1149 fatty acids by enterocytes isolated from rainbow trout (*Oncorhynchus mykiss*). Fish Physiol.
 1150 Biochem. 20, 125-134.

1151 Pike I., 2005. Eco-efficiency in aquaculture: global catch of wild fish used in aquaculture.
 1152 Internat. Aquafeed 8, 38-40.

1153 Poston, H.A., 1990a. Performance of rainbow trout fed supplemental soybean lecithin and
 1154 choline. Prog. Fish-Cult. 52, 218-225.

1155 Poston, H.A., 1990b. Effect of body size on growth, survival and chemical composition of
 1156 Atlantic salmon fed soy lecithin and choline. Prog. Fish-Cult. 52, 226-230.

1157 Poston, H.A., 1991. Response of Atlantic salmon fry to feed-grade lecithin and choline. Prog.
 1158 Fish-Cult. 53, 224-228.

1159 Rainuzzo, J. R., Reitan, K.I., Jorgensen, L., 1992. Comparative study on the fatty acid and lipid
 1160 composition of four marine fish larvae. Comp. Biochem. Physiol. 103B, 21-26.

1161 Rinchar, J., Czesny, S., Dabrowski, K., 2007. Influence of lipid class and fatty acid deficiency
 1162 on survival, growth, and fatty acid composition in rainbow trout juveniles. Aquaculture 264,
 1163 363-371.

1164 Robin, J.H., Vincent, B., 2003. Microparticulate diets as first food for gilthead sea bream larvae
 1165 (*Sparus aurata*): study of fatty acid incorporation. Aquaculture 225, 463-474.

1166 Ronnestad, I., Finn, R.N., Lein, I., Lie, Ø., 1995. Compartmental changes in the contents of total
 1167 lipid, lipid classes and their associated fatty acids in developing yolk-sac larvae of Atlantic
 1168 halibut, *Hippoglossus hippoglossus* (L.). Aquacult. Nutr. 1, 119-130.

1169 Saitoh, H., Ishihara, K., 1997. Antioxidant activity and active sites of phospholipids as
 1170 antioxidants. J. Am. Oil Chem. Soc. 74, 1531-1536.

1171 Salhi, M., Hernández-Cruz, C.M., Bessonart, M., Izquierdo, M.S., Fernández-Palacios, H., 1999.
 1172 Effect of different dietary polar lipid levels and different n-3 HUFA content in polar lipids on
 1173 gut and liver histological structure of gilthead sea bream (*Sparus aurata*) larvae. Aquaculture
 1174 179, 253-263.

1175 Salze, G., Tocher, D.R., Roy, W.J., Robertson, D.A., 2005. Egg quality determinants in cod
 1176 (*Gadus morhua* L.): Egg performance and lipids in eggs from farmed and wild broodstock.
 1177 Aquacult. Res. 36, 1488-1499.

1178 Sargent J.R., Tacon, A., 1999. Development of farmed fish: a nutritionally necessary alternative
 1179 to meat. Proc. Nutr. Soc. 58, 377-383.

1180 Sargent, J.R., Henderson, R.J., Tocher, D.R., 1989. The lipids. In: Halver, J.E. (Ed.), Fish
 1181 Nutrition, 2nd Edition. Academic Press, New York, pp. 153-218.

1182 Sargent, J.R., McEvoy, L.A., Bell, J.G., 1997. Requirements, presentation and sources of
 1183 polyunsaturated fatty acids in marine fish larval feeds. Aquaculture, 155, 119-129.

1184 Sargent, J.R., Bell, J.G., McEvoy, L., Tocher, D.R., Estevez, A., 1999a. Recent developments in
 1185 the essential fatty acid nutrition of fish. Aquaculture 177, 191-200.

1186 Sargent, J., McEvoy, L., Estevez, A., Bell, G., Bell, M., Henderson, J., Tocher, D.R., 1999b.
 1187 Lipid nutrition of marine fish during early development: current status and future directions.
 1188 Aquaculture 179, 217-229

1189 Sargent, J.R., Tocher, D.R., Bell, J.G., 2002. The lipids. In: Halver, J.E., Hardy, R.W. (Eds.),
 1190 Fish Nutrition, 3rd Edition. Academic Press, San Diego, pp. 181-257.

1191 Seiliez, I., Bruant, J.S., Zambonino-Infante, J., Kaushik, S., Bergot, P., 2006. Effect of dietary
 1192 phospholipid level on the development of gilthead sea bream (*Sparus aurata*) larvae fed a
 1193 compound diet. Aquacult. Nutr. 12, 372-378.

1194 Shewfelt, R. L., 1981. Fish muscle lipolysis - a review. J. Food Biochem. 5, 79-100.

1195 Sire, M. F., Lutton, C., Vernier, J.M. 1981. New views on intestinal absorption of lipids in
 1196 teleostan fishes: an ultrastructural and biochemical study in the rainbow trout. J. Lipid Res.
 1197 22, 81-94.

1198 Sitjà-Bobadilla, A., Peña-Llopis, S., Gómez-Requeni, P., Médale, F., Kaushik, S., Pérez-Sánchez,
 1199 J., 2005. Effect of fish meal replacement by plant protein sources on non-specific defence
 1200 mechanisms and oxidative stress in gilthead sea bream (*Sparus aurata*). Aquaculture 249,
 1201 387-400.

1202 Snyder, F. (Ed), 1987. Platelet-Activating Factor and Related Lipid Mediators. Plenum Press,
 1203 New York.

1204 Tacon A.G.J., 2004. Use of fish meal and fish oil in aquaculture: a global perspective. Aquatic
 1205 Resources, Cult. Develop. 1, 3-14.

1206 Taggart, J.B., Bron, J.E., Martin, S.A.M., Seear, P.J., Hoyheim, B., Talbot, R., Villeneuve, L.,
 1207 Sweeney, G.E., Houlihan, D.F., Secombes, C.J., Tocher, D.R., Teale, A.J., 2008. A
 1208 description of the origins, design and performance of the TRAITS/SGP Atlantic salmon
 1209 (*Salmo salar* L.) cDNA microarray. J. Fish Biol., in press.

1210 Takeuchi, T., Arakawa, T., Satoh, S., Watanabe, T., 1992. Supplemental effect of phospholipids
 1211 and requirement of eicosapentaenoic acid and docosahexaenoic acid of juvenile striped jack.
 1212 Nippon Suisan Gakkaishi 58, 707-713.

1213 Tian, Y., Jackowski, S., 2007. Golgi-mediated secretion requires de novo phospholipids
 1214 synthesis. Chem. Physics Lipids 149S, S11-S22.

1215 Tidwell, J.H., Allan, G.L., 2002. Fish as food: aquaculture's contribution. World Aquacult. 33,
 1216 44-48.

1217 Tocher, D.R., 1995. Glycerophospholipid metabolism. In: Hochachka, P.W., Mommsen, T.P.
 1218 (Eds.), Biochemistry and Molecular Biology of Fishes, Vol. 4. Metabolic and Adaptational
 1219 Biochemistry. Elsevier Press, Amsterdam, pp. 119-157.

1220 Tocher, D.R., 2003. Metabolism and functions of lipids and fatty acids in teleost fish. Rev.
 1221 Fisheries Sci. 11, 107-184.

1222 Tocher, D.R., Fraser, A.J., Sargent, J.R., Gamble, J.C., 1985a. Lipid class composition during
 1223 embryonic and early larval development in Atlantic herring (*Clupea harengus*, L.). Lipids
 1224 20, 84-89.

1225 Tocher, D.R., Fraser, A.J., Sargent, J.R., Gamble, J.C., 1985b. Fatty acid composition of
 1226 phospholipids and neutral lipids during embryonic and early larval development in Atlantic
 1227 herring (*Clupea harengus*, L.). Lipids 20, 69-74.

1228 Turner, M.R., Lumb, R.H., 1989. Synthesis of platelet activating factor by tissues from the
 1229 rainbow trout, *Salmo gairdneri*. Biochim. Biophys. Acta 1004, 49-52.

1230 Uyan, O., Koshio, S., Ishikawa, M., Uyan, S., Ren, T., Yokoyama, S., Komilus, C.F., Michael,
 1231 F.R., 2007. Effects of dietary phosphorus and phospholipid level on growth, and phosphorus
 1232 deficiency signs in juvenile Japanese flounder, *Paralichthys olivaceus*. Aquaculture 267, 44-
 1233 54.

1234 Van Der Kraak, G., 1990. The influence of calcium ionophore and activators of protein kinase C
 1235 on steroid production by preovulatory ovarian follicles of the goldfish. Biol. Reprod. 42, 231-
 1236 238.

1237 Vernier, J.-M., Sire, M.-F., 1986. Is the Golgi apparatus the obligatory final step for lipoprotein
 1238 secretion by intestinal cells? Tissue Cell 18, 447-460.

1239 Villeneuve, L., Gisbert, E., Zambonino Infante, J.L., Quazuguel, P., Cahu, C.L., 2005. Effect of
 1240 nature of dietary lipids on European sea bass morphogenesis: implication of retinoid receptors.
 1241 Br. J. Nutr. 94, 877-884.

1242 Villeneuve, L., Gisbert, E., Moriceau, J., Cahu, C.L., Zambonino Infante, J.L., 2006. Intake of
 1243 high levels of vitamin A and polyunsaturated fatty acids during different developmental
 1244 periods modifies the expression of morphogenesis genes in European sea bass (*Dicentrarchus*
 1245 *labrax*). Br. J. Nutr. 95, 677-687.

1246 Wade, M. G., Van Der Kraak, G., 1991. The control of testicular androgen production in the
 1247 goldfish: Effects of activators of different intracellular signalling pathways. Gen. Comp.
 1248 Endocrinol. 83, 337-344.

1249 Wassef, E.A., Wahby, O.M., Sakr, E.M., 2007. Effect of dietary vegetable oils on health and liver
 1250 histology of gilthead seabream (*Sparus aurata*) growers. Aquacult. Res. 38, 852-861.

1251 Wiegand, M.D., 1996. Composition, accumulation and utilization of yolk lipids in teleost fish.
 1252 Rev. Fish Biol. Fish. 6, 259-286.

1253 Weirich, C.R., Reigh, R.C., 2001. Dietary lipids and stress tolerance of larval fish. In: Lim, C.,
 1254 Webster, C.D. (Eds.), Nutrition and Fish Health. Food Products Press, NY, pp. 301-312.

1255 Yufera, M., Fernandez-Diaz, C., Pascual, E., Sarasquete, M., Moyano, F., Diaz, M., Alarcon, F.,
 1256 Garcia-Gallego, M., Parra, G., 2000. Towards an inert diet for first-feeding gilthead sea bream
 1257 *Sparus aurata* larvae. Aquacult. Nutr. 6, 143-152.

1258 Zambonino Infante, J.L., Cahu, C.L., 2001. Ontogeny of the gastrointestinal tract of marine fish
 1259 larvae. Comp. Biochem. Physiol. 130C, 477-487.

1260 Zambonino Infante, J.L., Cahu, C.L., 2007. Dietary modulation of some digestive enzymes and
 1261 metabolic processes in developing marine fish: applications to diet formulation. Aquaculture
 1262 268, 98-105.

1263




Fig.1. Biosynthetic pathways for the major phospholipid classes. CDP-Cho, CDP-choline; CDP-DAG, CDP-diacylglycerol; CDP-Etn, CDP-ethanolamine; DAG, diacylglycerol; PA, phosphatidic acid; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PS, phosphatidylserine; SM, sphingomyelin. The enzymes are;- 1, CDP-DAG synthases; 2, PA-phosphatases; 3, PI synthase; 4, PS synthase; 5, CDP-choline:DAG phosphotransferase; 6, CDP-ethanolamine:DAG phosphotransferase; 7, PE methyltransferase; 8, PS decarboxylase; 9, SM synthase; 10 and 11, PS synthases via base exchange.

Table 1. Quantitative and qualitative phospholipid requirements of teleost fish

Species	Developmental stage	Phospholipid supplement ^a and levels studied ^b	Optimal requirement and criteria used ^c	Feeding period	Reference
Ayu (<i>Plecoglossus altivelis</i>)	Larvae	0 and 3 % SL or EL	3 % (G,S,M)	20 days	Kanazawa et al. 1981
	Larvae	0, 1, 3 and 5 % SL	3 % (M), 5 % (G,S)	50 days	Kanazawa et al. 1983a
	Larvae	0 and 3 % EL or BPL	3 % (G,S,M)	50 days	Kanazawa et al. 1983a
	Juvenile	0 and 3 % SL or BPL	3 % (G)	33 days	Kanazawa et al. 1981
	Juvenile	0, 1, 3 and 5 % EL	3 % (G)	33 days	Kanazawa et al. 1981
Japanese flounder (<i>Paralichthys olivaceus</i>)	Larvae	0, 3, 5 and 7 % SL	7 % (G,S)	30 days	Kanazawa 1993
	Juvenile	0, 3, 5 and 7 % SL	7 % (G)	30 days	Kanazawa 1993
Knife jaw (<i>Oplegnathus fasciatus</i>)	Larvae	0, 2.5, 5 and 7.4 % SL	7.4 % (G,S)	22 days	Kanazawa et al. 1983b
	Larvae	0, 3, 5 and 7 % SL	5 % (G,S,R)	28 days	Kanazawa 1993
	Juvenile	0, 3, 5 and 7 % SL	3 % (G)	60 days	Kanazawa 1993
European sea bass (<i>Dicentrarchus labrax</i>)	Larvae	3, 6, 9 and 12 % SL	12 % (G,S,M)	40 days	Cahu et al. 2003b
	Juvenile	0 and 3 % SL	3 % (G)	40 days	Geurden et al. 1995a
	Juvenile	0 and 2 % EPC or SPC	2 % (G)	40 days	Geurden et al. 1995a
Red sea bream (<i>Pagrus major</i>)	Larvae	0 and 5 % SL	5 % (G,S)	20 days	Kanazawa et al., 1983b
Common carp (<i>Cyprinus carpio</i>)	Larvae	0 and 2 % EL	2 % (G,S)	25 days	Geurden et al. 1995b
	Larvae	0 and 2 % PL	2 % (G,S)	21 days	Geurden et al. 1995b
	Larvae	0 and 2 % SPC, SPI or EL	2 % (G,S,M except EL)	25 days	Geurden et al. 1997a
	Larvae				Geurden et al. 1998
Pikeperch (<i>Sander lucioperca</i>)	Larvae	1, 5 and 9 % SL	9 % (G)	24 days	Hamza et al. 2008
Gilthead sea bream (<i>Sparus aurata</i>)	Larvae	9, 11 and 15 % SL	> 9 % (G,S)	23 days	Seiliez et al. 2006
Striped jack (<i>Pseudocaranx dentex</i>)	Juvenile	0, 0.5, 1, 1.5 and 2 % SPC	1.5 % (G,S,R)	6 weeks	Takeuchi et al. 1992
	Juvenile	0 and 1.5 % SPE	1.5 % (G)	6 weeks	Takeuchi et al. 1992
Rainbow trout (<i>Oncorhynchus mykiss</i>)	Juvenile	0, 2, 4 and 8 % SL	4 % (G)	20 weeks	Poston 1990a
	Juvenile	0 and 14 %	14% (G)	8 weeks	Rinchard et al. 2007
Turbot (<i>Psetta maximus</i>)	Juvenile	0 and 2 % EL	2% (G)		Geurden et al. 1997b
Atlantic salmon (<i>Salmo salar</i>)	Juvenile (180 mg)	0, 2, 4, 6 and 8 % SL/CPL	6 % (G)	14 weeks	Poston 1991
	Juvenile (180 mg)	0 and 4 % SL	4 % (G)	16 weeks	Poston 1990b
	Juvenile (1.0 g)	0 and 4 % SL	4 % (G)	12 weeks	Poston 1990b
	Juvenile (1.7 g)	0 and 4 % SL	4 % (G)	12 weeks	Poston 1990b
	Juvenile (7.5 g)	0 and 4 % SL	0 % (no requirement)	12 weeks	Poston 1990b
White sturgeon (<i>Acipenser transmontanus</i>)	Juvenile (5-10 g)	0 and 8% SL	0% (no requirement)	6 weeks	Hung and Lutes 1988

^a BPL, bonito egg polar lipid; CPL, corn polar lipid; EL, chicken egg lecithin; EPC, purified egg phosphatidylcholine ;

PL, various phospholipid sources supplemented to supply 2 % dietary phospholipids including EL, SL, sunflower, rapeseed and marine phospholipids: SL, soybean lecithin; SPC, purified soybean phosphatidylcholine; SPE, purified soybean phosphatidylethanolamine;

SPI, purified soybean phosphatidylinositol; ^b Percentage of diet weight; ^c G, growth; S, survival; M, malformations; R, stress resistance.

Table 2. Lipid content (percentage of wet weight) and lipid class composition (% total lipid) of commercially available fishmeals and traditional feeds for salmonids based on high levels of fishmeal

Lipid class	Fishmeals				Feeds	
	1	2	3	4	A	B
Lipid content	10.4	10.0	8.5	5.9	16.5	34.4
Phosphatidylcholine (PC)	7.5	12.1	11.8	9.3	2.5	1.1
Phosphatidylethanolamine	4.1	5.8	6.7	3.9	1.4	0.0
Phosphatidylserine	0.9	1.4	1.6	0.7	0.0	0.0
Phosphatidylinositol	1.2	1.8	2.1	1.9	0.0	0.0
CL/PG	0.6	1.6	1.5	2.7	0.0	0.0
Sphingomyelin	1.6	1.3	1.5	1.5	0.0	0.0
Lyso-PC	1.0	1.3	1.5	0.5	0.0	0.0
Total polar lipids	16.9	23.4	26.7	20.4	3.9	1.1
Total neutral lipids	83.2	76.6	73.3	79.6	96.1	98.9
Triacylglycerol	55.0	44.3	47.7	66.5	69.2	78.6
Free fatty acids	15.3	14.2	15.1	5.6	11.9	9.3
Sterols	9.3	8.9	9.9	7.5	10.6	8.8
Steryl esters	3.6	9.1	0.7	0.0	4.5	2.2

Values are means of duplicate analyses. CL, cardiolipin; PG, phosphatidylglycerol.

1273

1274

Table 3. Lipid content (percentage of wet weight) and lipid class composition (% total lipid) of feed ingredients

Lipid class/diet	Oils				Meals										Binders			Lecithin	
	RO	SO	NFO	SFO	Corn gluten	Wheat gluten	SF cake	Soy SPC60	HP soya	Peas	Rice protein	Blood meal	Hb meal	German FeaM	French FeaM	Field beans	Raw KB	Cooked KB	BergaPur
Lipid content	~100	~100	~100	~100	5.6	9.9	8.0	0.8	1.9	1.8	9.6	1.4	1.0	6.8	6.6	1.5	1.7	1.4	74.5
Phosphatidylcholine (PC)	0.5	0.0	0.0	0.0	0.5	2.2	4.7	9.6	14.6	13.6	0.5	7.8	9.8	0.0	0.6	12.4	3.8	7.9	19.8
Phosphatidylethanolamine	1.2	0.0	0.0	0.0	0.0	2.8	1.4	6.7	7.8	7.6	0.3	3.7	5.5	0.0	0.0	7.2	1.5	5.9	17.6
Phosphatidylserine	0.0	0.0	0.0	0.0	0.0	0.0	2.0	6.3	9.4	4.7	1.0	0.6	1.3	0.0	0.0	5.2	2.9	6.1	15.0
Phosphatidylinositol	0.0	0.0	0.0	0.0	0.0	1.8	0.6	4.8	5.5	1.5	0.7	1.7	2.6	0.0	0.0	1.6	2.2	0.8	12.6
PG/PA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9	0.3	0.0	0.0	0.0	0.0	0.8	0.0	0.0	0.0
Sphingomyelin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	6.9	11.8	0.0	0.3	0.0	0.0	0.0	0.0
Lyso-PC	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.8	1.6	0.5	4.9	1.1	0.0	0.0	0.2	0.7	1.1	1.0	1.0
Total phospholipid	1.7	0.0	0.0	0.0	0.5	6.8	9.0	28.2	38.9	28.9	7.7	21.8	31.0	0.0	1.1	27.8	11.5	21.7	66.1
MGDG	0.0	0.0	0.0	0.0	0.0	2.4	3.3	4.1	4.2	3.1	1.7	0.0	0.0	0.0	0.0	2.7	4.0	3.9	9.8
DGDG	0.0	0.0	0.0	0.0	0.0	9.7	0.0	0.9	2.0	2.0	0.4	0.9	0.0	0.0	0.0	2.1	1.8	0.8	2.7
Other glycolipids	0.0	0.0	0.0	0.0	0.0	7.8	0.0	2.2	4.3	3.6	0.9	0.2	1.2	0.4	0.0	3.6	6.9	4.9	4.3
Total polar lipids	1.7	0.0	0.0	0.0	0.5	26.7	12.4	35.4	49.4	37.6	10.7	22.8	32.1	0.4	1.1	36.1	24.2	31.3	82.9
Total neutral lipids	98.3	100.0	100.0	100.0	99.5	73.3	87.6	64.6	50.6	62.4	89.3	77.2	67.9	99.6	98.9	63.9	75.8	68.7	17.1
Triacylglycerol	74.0	83.3	83.9	73.3	52.1	46.3	64.8	47.6	33.0	38.6	39.0	15.5	6.6	31.7	34.0	37.2	30.0	40.5	12.7
Free fatty acid	9.6	8.0	9.3	12.9	17.0	12.4	7.1	6.2	4.2	6.2	36.1	13.5	10.6	39.1	31.8	9.8	32.9	14.1	1.6
Total sterols	8.2	7.9	6.8	10.0	19.3	7.4	5.7	4.0	8.1	10.2	12.0	29.4	35.4	17.6	17.9	8.2	11.8	10.9	2.8
Steryl ester	6.5	0.8	0.0	3.9	11.1	2.1	3.4	3.9	0.5	3.0	0.0	18.9	15.2	11.1	15.2	3.0	1.0	0.5	0.0
Unknown neutral lipid	0.0	0.0	0.0	0.0	0.0	5.2	6.6	2.9	4.8	4.4	0.0	0.0	0.0	0.0	0.0	5.7	0.0	2.7	0.0

Values are means of duplicate analyses. BergaPur is a deoiled lecithin (98%) (Berg+Schmidt, Hamburg, Germany); DGDG, digalactosyldiacylglycerols; FeaM, feather meal; Hb, Hemoglobin; KB, kidney beans; MGDG, monogalactosyldiacylglycerols; NFO, North Atlantic fish oil; PA, phosphatidic acid; PG, phosphatidylglycerol; RO, rapeseed oil; SF, sunflower; SFO, South American fish oil; SO, soybean oil.

1274

1275

Table 4. Cholesterol content (mg/100g product) of traditional feeds and alternative feed ingredients

	Product	Cholesterol content
<u>Feeds</u>	Feed A	100.2
	Feed B	112.5
<u>Oils</u>	Rapeseed oil	5.9
	Soybean oil	4.8
	North Atlantic fish oil	359.9
	South American fish oil	324.3
<u>Meals</u>	Corn gluten	1.0
	Wheat gluten	8.1
	Sunflower cake	1.0
	SPC 60	0.2
	High protein soya	0.1
	Peas	0.4
	Rice protein	6.8
	Blood meal	218.7
	Haemoglobin meal	251.8
	German feather meal	134.3
	French feather meal	142.9
<u>Binders</u>	Field beans	4.3
	Raw kidney beans	1.1
	Cooked kidney beans	0.4
<u>Lecithin</u>	Bergapur	7.4

Values are means of duplicate analyses.

1275

1276

1277