

753 **Table 3**754 Growth and performance parameters in three strains of Atlantic salmon fed the experimental fish oil (FO) and vegetable oil (VO) diets for 55
755 weeks

								Two-way ANOVA (P value)		
757	Parameter/Diet and strain	Lean FO	Lean VO	Fat FO	Fat VO	CAL FO	CAL VO	Diet	Strain	Interaction
758	Initial weight (g)	80.6 ± 11.6	89.5 ± 12.9	92.1 ± 4.9	84.3 ± 8.7	52.8 ± 1.4	51.6 ± 2.1	0.995	0.002	0.409
759	Final weight (kg)	3.12 ± 0.13	3.03 ± 0.09	3.18 ± 0.08	2.84 ± 0.04	2.75 ± 0.13	2.89 ± 0.04	0.118	0.019	0.027
760	Final length (cm)	65.3 ± 0.5	63.7 ± 1.0	63.9 ± 1.0	62.5 ± 0.7	63.1 ± 1.6	62.8 ± 0.9	0.101	0.139	0.650
761	SGR	0.92 ± 0.02	0.91 ± 0.01	0.93 ± 0.01	0.89 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	0.109	<0.0001	0.065
762	FCR	1.23 ± 0.03	1.39 ± 0.22	1.08 ± 0.09	1.19 ± 0.03	1.08 ± 0.06	1.06 ± 0.03	0.211	0.041	0.495
763	TGC	3.55 ± 0.14	3.50 ± 0.05	3.57 ± 0.05	3.38 ± 0.03	3.57 ± 0.08	3.66 ± 0.02	0.247	0.072	0.087
764	CF (%)	1.17 ± 0.03	1.19 ± 0.02	1.18 ± 0.03	1.22 ± 0.04	1.07 ± 0.06	1.12 ± 0.03	0.129	0.013	0.742
765	HSI (%)	0.71 ± 0.00	0.70 ± 0.07	0.59 ± 0.03	0.61 ± 0.00	0.62 ± 0.04	0.61 ± 0.01	0.896	0.010	0.809
766	VSI (%)	4.92 ± 0.32	4.82 ± 0.01	5.17 ± 0.09	5.51 ± 0.18	4.97 ± 0.32	4.82 ± 0.01	0.730	0.034	0.291

767 All values are mean ± SD (n = 2). Fish weights and lengths were obtained by weighing 50 fish/pen. SGR, specific growth rate; FCR, feed
 768 conversion ratio; TGC, thermal growth coefficient; CF, condition factor; HIS, hepato-somatic index; VSI, viscero-somatic index. P values
 769 shown in bold are significantly different (P < 0.05).

770 **Table 4**

771 Lipid content (%) of flesh, viscera and liver and protein content of flesh in three strains of Atlantic salmon fed the experimental fish oil (FO) and
772 vegetable oil (VO) diets for 55 weeks

773								Two-way ANOVA (P value)		
774	Parameter/Diet and strain	Lean FO	Lean VO	Fat FO	Fat VO	CAL FO	CAL VO	Diet	Strain	Interaction
775	Flesh lipid (%)	11.6 ± 0.2	12.8 ± 0.3	13.2 ± 0.2	12.9 ± 0.2	12.3 ± 0.3	11.3 ± 0.2	0.816	0.0008	0.0002
776	Viscera lipid (%)	28.0 ± 0.4	29.6 ± 0.3	31.7 ± 0.7	29.4 ± 0.3	25.8 ± 0.4	26.8 ± 0.3	0.702	<0.0001	<0.0001
777	Liver lipid (%)	4.0 ± 0.3	4.1 ± 0.1	3.4 ± 0.1	4.8 ± 0.4	3.2 ± 0.1	3.6 ± 0.2	0.003	0.009	0.017
778	Flesh protein (%)	19.4 ± 0.1	19.9 ± 0.3	19.1 ± 0.2	19.2 ± 0.1	19.3 ± 0.1	19.5 ± 0.1	0.034	0.016	0.291

779 All values are mean ± SD, n = 2. P values shown in bold are significantly different (P < 0.05).

780

781

782

783

784

785 **Table 5**

786 Fatty acid compositions (% by weight of total fatty acids) of flesh from three strains of Atlantic salmon fed the experimental fish oil (FO) and
 787 vegetable oil (VO) diets for 55 weeks

							Two-way ANOVA (P value)		
Fatty acid/Diet and strain	Lean FO	Lean VO	Fat FO	Fat VO	CAL FO	CAL VO	Diet	Strain	Interaction
14:0	4.5 ± 0.0	1.4 ± 0.0	4.6 ± 0.1	1.5 ± 0.1	4.5 ± 0.2	1.5 ± 0.2	<0.0001	0.471	0.761
16:0	15.0 ± 0.3	13.1 ± 0.1	15.0 ± 0.3	13.7 ± 0.2	15.4 ± 0.3	13.8 ± 0.2	<0.0001	0.044	0.271
18:0	2.6 ± 0.1	2.9 ± 0.1	2.7 ± 0.1	3.0 ± 0.1	2.8 ± 0.0	3.2 ± 0.1	0.007	0.023	0.687
Total saturated ¹	22.6 ± 0.4	17.9 ± 0.3	22.8 ± 0.5	18.8 ± 0.4	23.1 ± 0.5	18.9 ± 0.3	<0.0001	0.071	0.449
16:1n-7	5.5 ± 0.1	1.7 ± 0.1	5.6 ± 0.1	2.0 ± 0.1	5.6 ± 0.1	1.9 ± 0.2	<0.0001	0.132	0.548
18:1n-9	17.7 ± 0.5	37.7 ± 0.3	17.8 ± 0.7	37.3 ± 0.4	18.5 ± 0.3	37.1 ± 0.6	<0.0001	0.765	0.204
18:1n-7	3.2 ± 0.2	2.4 ± 0.2	3.2 ± 0.1	2.6 ± 0.5	3.5 ± 0.2	2.7 ± 0.2	0.003	0.332	0.831
20:1n-9	7.6 ± 0.1	5.2 ± 0.1	7.5 ± 0.1	5.1 ± 0.1	7.3 ± 0.1	5.1 ± 0.1	<0.0001	0.079	0.332
22:1n-11	7.6 ± 0.2	1.3 ± 0.1	7.6 ± 0.2	1.5 ± 0.1	7.2 ± 0.4	1.3 ± 0.2	<0.0001	0.234	0.492
22:1n-9	0.9 ± 0.2	0.7 ± 0.0	0.8 ± 0.1	0.7 ± 0.0	0.9 ± 0.4	0.7 ± 0.0	0.174	0.911	0.911
Total monounstaurates ²	43.7 ± 0.5	49.9 ± 0.3	43.9 ± 0.3	49.8 ± 0.4	44.3 ± 0.6	49.6 ± 0.3	<0.0001	0.877	0.361
18:2n-6	4.9 ± 0.1	14.0 ± 0.2	4.9 ± 0.1	13.3 ± 0.2	5.0 ± 0.1	13.3 ± 0.3	<0.0001	0.069	0.041
20:2n-6	0.6 ± 0.0	1.6 ± 0.0	0.6 ± 0.0	1.4 ± 0.0	0.6 ± 0.0	1.5 ± 0.0	<0.0001	<0.0001	<0.0001
20:3n-6	0.3 ± 0.0	0.6 ± 0.0	0.3 ± 0.0	0.6 ± 0.0	0.3 ± 0.0	0.6 ± 0.0	<0.0001	1.000	1.000

804	20:4n-6	0.4 ± 0.0	0.2 ± 0.0	0.4 ± 0.0	0.2 ± 0.0	0.4 ± 0.0	0.2 ± 0.0	<0.0001	1.000	1.000
805	Total n-6 ³	6.5 ± 0.2	16.5 ± 0.2	6.4 ± 0.1	15.7 ± 0.2	6.6 ± 0.1	15.8 ± 0.3	0.043	0.995	0.955
806	18:3n-3	1.8 ± 0.1	6.7 ± 0.1	1.9 ± 0.0	6.3 ± 0.1	1.8 ± 0.1	6.2 ± 0.2	<0.0001	0.058	0.341
807	18:4n-3	1.7 ± 0.1	0.7 ± 0.0	1.8 ± 0.1	0.8 ± 0.0	1.5 ± 0.1	0.7 ± 0.0	<0.0001	0.020	0.148
808	20:4n-3	1.7 ± 0.1	0.9 ± 0.0	1.7 ± 0.0	0.9 ± 0.0	1.6 ± 0.1	0.8 ± 0.0	<0.0001	0.079	0.079
809	20:5n-3	5.8 ± 0.1	1.7 ± 0.1	5.7 ± 0.2	1.9 ± 0.1	5.5 ± 0.2	1.9 ± 0.1	<0.0001	0.630	0.115
810	22:5n-3	2.9 ± 0.1	0.9 ± 0.1	2.8 ± 0.1	0.9 ± 0.0	2.8 ± 0.1	0.9 ± 0.1	<0.0001	0.687	0.687
811	22:6n-3	13.1 ± 0.4	4.0 ± 0.1	12.8 ± 0.4	4.2 ± 0.2	12.7 ± 0.6	4.5 ± 0.2	<0.0001	0.926	0.280
812	Total n-3 ⁴	27.1 ± 0.7	15.7 ± 0.3	26.9 ± 0.7	15.7 ± 0.4	26.0 ± 1.0	15.7 ± 0.2	<0.0001	0.453	0.453
813	Total PUFA	33.6 ± 0.8	32.2 ± 0.3	33.3 ± 0.8	31.4 ± 0.6	32.6 ± 1.1	31.5 ± 0.4	0.012	0.309	0.740
814	Total n-3 LC-PUFA	23.6 ± 0.6	8.3 ± 0.3	23.2 ± 0.5	8.6 ± 0.3	22.7 ± 0.8	8.8 ± 0.2	<0.0001	0.842	0.216
815	n-3/n-6	4.2 ± 0.4	1.0 ± 0.2	4.2 ± 0.4	1.0 ± 0.3	3.9 ± 0.5	1.0 ± 0.2	<0.0001	0.792	0.792

816

817 Values are mean ± SD, n = 2. ¹Includes 15:0, 17:0, 20:0 and 22:0. ²Includes 16:1n-9, 20:1n-7 and 20:1n-11. ³Includes 18:3n-6 and 22:5n-6.

818 ⁴Includes 20:3n-3. P values shown in bold are significantly different (P < 0.05).